

SOME COMPOSITION FORMULAS OF JACOBI TYPE ORTHOGONAL POLYNOMIALS

PRADEEP MALIK AND SAIFUL R. MONDAL

ABSTRACT. The composition of Jacobi type finite classes of the classical orthogonal polynomials with two generalized Riemann-Liouville fractional derivatives are considered. The outcomes are expressed in terms of generalized Wright function or generalized hypergeometric function. Similar composition formulas are also obtained by considering the generalized Riemann-Liouville and Erdélyi-Kober fractional integral operators.

1. introduction

It is evident from various research articles [3, 4, 8, 17, 22, 23, 25–27, 32] that the fractional calculus have wide applications in solving various integral equations, ordinary differential equations and partial differential equations in applied sciences like as turbulence and fluid dynamics, stochastic dynamical system, nonlinear control theory, nonlinear biological systems. As per the requirement, the fractional integration and differential operators have gone through several improvement and extensions, which can be seen in [1, 2, 9–14, 30, 31].

In this article, we will study the composition formula of fractional integration given in [15, 17, 29, 32] and the fractional derivatives given in [10] and [2]. Outcomes of the composition are expressed in term of generalized hypergeometric or Wright functions [28, 33, 34]. Similar type of work can be seen in [20, 24] and references therein.

For this purpose, first we will give a brief introductions about the fractional integral operators and its generalization. Then two type generalized fractional derivatives and its properties are discussed. At the end of this section, we introduced the finite classes require in sequel.

Received September 18, 2016.

2010 *Mathematics Subject Classification.* 26A33, 33C45, 33C05, 33C20, 33C60, 26A09, 33E20.

Key words and phrases. fractional integral transform, classical orthogonal polynomials, generalized Wright function, generalized hypergeometric function.

©2017 Korean Mathematical Society

1.1. Riemann-Liouville fractional integrals and its generalization

The left and right-sided generalized integral transforms defined for $x > 0$ and complex $\alpha, \beta, \eta \in \mathbb{C}$, and $\text{Re}(\alpha) > 0$ are given in [15, 17, 29, 32], respectively as

$$(1.1) \quad \left(I_{0+}^{\alpha, \beta, \eta} f\right)(x) = \frac{x^{-\alpha-\beta}}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} {}_2F_1\left(\alpha + \beta, -\eta; \alpha; 1 - \frac{t}{x}\right) f(t) dt$$

and

$$(1.2) \quad \left(I_-^{\alpha, \beta, \eta} f\right)(x) = \frac{1}{\Gamma(\alpha)} \int_x^\infty (t-x)^{\alpha-1} t^{-\alpha-\beta} {}_2F_1\left(\alpha + \beta, -\eta; \alpha; 1 - \frac{x}{t}\right) f(t) dt,$$

where $\Gamma(\alpha)$ represents the gamma function [28] and $\text{Re}(\alpha)$ denotes the real part of α .

The generalized hypergeometric function ${}_pF_q(a_1 \cdots a_p; c_1 \cdots c_q; x)$ is given by the representation

$$(1.3) \quad {}_pF_q(a_1 \cdots a_p; c_1 \cdots c_q; x) = \sum_{k=0}^{\infty} \frac{(a_1)_k \cdots (a_p)_k}{(c_1)_k \cdots (c_q)_k (1)_k} x^k,$$

where none of the denominator parameters can be zero or a negative integer and $(a)_n$ is the well-known Pochhammer symbol given by $(\lambda)_n = \lambda(\lambda+1)\cdots(\lambda+n-1)$; $(\lambda)_0 = 1$. Results related to the generalized hypergeometric functions are abundant in the literature. For example, see [28]. In particular, for $p = 2$ and $q = 1$, we get the Gaussian hypergeometric function $F(a, b; c; x) = {}_2F_1(a, b; c; x)$ is defined by

$${}_2F_1(a, b; c; x) = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{(c)_k (1)_k} x^k,$$

where a, b are complex numbers and $c \in \mathbb{C} \setminus \{0, -1, -2, \dots\}$. The series in (1.3) is absolutely convergent for $|x| < 1$ and $|x| = 1$, when $\text{Re}(c - a - b) > 0$.

For $\beta = -\alpha$, (1.1) and (1.2) coincide respectively, with the classical left and right-hand sided Riemann-Liouville fractional integrals [12, 32] for $x > 0$ of order $\alpha \in \mathbb{C}$, $\text{Re}(\alpha) > 0$,

$$(1.4) \quad \left(I_{0+}^{\alpha, -\alpha, \eta} f\right)(x) = \left(I_{0+}^{\alpha} f\right)(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} f(t) dt, \quad (x > 0)$$

and

$$(1.5) \quad \left(I_-^{\alpha, -\alpha, \eta} f\right)(x) = \left(I_-^{\alpha} f\right)(x) = \frac{1}{\Gamma(\alpha)} \int_x^\infty (t-x)^{\alpha-1} f(t) dt, \quad (x > 0).$$

If $\beta = 0$, (1.1) and (1.2) respectively, are the so-called Erdélyi-Kober fractional integrals [32] defined for complex $\alpha, \eta \in \mathbb{C}$, $\text{Re}(\alpha) > 0$

$$(1.6) \quad \left(I_{0+}^{\alpha, 0, \eta} f\right)(x) = \left(I_{\alpha, \eta}^+ f\right)(x) = \frac{x^{-\alpha-\eta}}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} t^\eta f(t) dt, \quad (x > 0)$$

and

$$(1.7) \quad \left(I_{-}^{\alpha,0,\eta} f \right) (x) = \left(K_{\eta,\alpha}^{-} f \right) (x) = \frac{x^{\eta}}{\Gamma(\alpha)} \int_x^{\infty} (t-x)^{\alpha-1} t^{-\alpha-\eta} f(t) dt \quad (x > 0).$$

1.2. Riemann-Liouville fractional derivatives and its generalization

The classical Riemann-Liouville fractional derivative of f of order μ is defined by

$$(1.8) \quad D_x^{\alpha} f(x) := \begin{cases} \frac{1}{\Gamma(-\alpha)} \int_0^x \frac{f(t)}{(x-t)^{\alpha+1}} dt & \text{if } \operatorname{Re}(\alpha) < 0, \\ \frac{d^m}{dx^m} D_{0+}^{\alpha-m} f(x) & \text{if } m-1 < \operatorname{Re}(\alpha) < m. \end{cases}$$

In recent years, many authors have developed various extended fractional derivative formulas of Riemann-Liouville type. Here, we consider two generalized fractional derivative.

First, we consider the extension defined in [2] by

$$(1.9) \quad D_x^{\mu,\sigma} f(x) := \frac{1}{\Gamma(-\mu)} \int_0^x f(t) (z-t)^{-\mu-1} \exp\left(-\frac{\sigma x^2}{t(x-t)}\right) dt,$$

with $\operatorname{Re}(\mu) < 0$ and $\operatorname{Re}(\sigma) > 0$. For $f(x) = x^{\nu}$, it follows that

$$(1.10) \quad D_x^{\mu,\sigma} (x^{\nu}) = \frac{B_{\sigma}(\nu+1, -\mu)}{\Gamma(-\mu)} x^{\nu-\mu}$$

for $\operatorname{Re}(\nu) > -1$ and $\operatorname{Re}(\mu) < 0$. Further, if f is analytic in the disc $|x| < \rho$ and have the power series $f(x) = \sum_{n=0}^{\infty} a_n x^n$, then

$$(1.11) \quad D_x^{\mu,\sigma} (x^{\lambda-1} f(x)) = \frac{x^{\lambda-\mu}}{\Gamma(-\mu)} \sum_{n=0}^{\infty} a_n B_{\sigma}(\lambda+n, -\mu) x^n,$$

provided $\operatorname{Re}(\lambda) > -1$ and $\operatorname{Re}(\mu) < 0$. Here the function B_{σ} is the extended beta function, which was introduced by Chaudhry et al. [5] and defined by

$$(1.12) \quad B_{\sigma}(a, b) := \int_0^1 t^{a-1} (1-t)^{b-1} \exp\left(-\frac{\sigma}{t(1-t)}\right) dt,$$

when $\operatorname{Re}(a) > 0$, $\operatorname{Re}(b) > 0$, $\operatorname{Re}(\sigma) > 0$. This extended beta functions have close association with the Macdonald, error and Whittaker functions, and using this, an extension of the hypergeometric functions can be found in [6].

Recently, Kataugampola [10] introduced a generalized fractional derivatives operator ${}^{\rho}D_{0+}^{\alpha}$ defined by

$$(1.13) \quad {}^{\rho}D_{0+}^{\alpha} f(x) := \begin{cases} \frac{\rho^{\alpha+1}}{\Gamma(-\alpha)} \int_0^x \frac{f(t) t^{\rho-1}}{(x^{\rho}-t^{\rho})^{\alpha+1}} dt & \text{if } \operatorname{Re}(\alpha) < 0, \\ \left(x^{-\rho-1} \frac{d}{dx}\right)^m {}^{\rho}D_{0+}^{\alpha-m} f(x) & \text{if } m-1 < \operatorname{Re}(\alpha) < m, \end{cases}$$

where $m \in \mathbb{N}$ and $\rho > 0$.

For the function $f(x) = x^\nu$, $\nu \in \mathbb{R}$, the generalized fractional derivative operator with $\text{Re}(\alpha) < 0$ defined in (1.13) yields the formula

$$(1.14) \quad {}^\rho D_{0+}^\alpha x^\nu = \frac{\rho^{\alpha-1} \Gamma\left(\frac{\nu}{\rho} + 1\right)}{\Gamma\left(\frac{\nu}{\rho} + 1 - \alpha\right)} x^{\nu-\alpha\rho}.$$

Replacing $\rho = 1$ in (1.14) gives the classical Riemann-Liouville fractional integral for the function x^ν , which is,

$$(1.15) \quad D_x^\alpha x^\nu = \frac{\Gamma(\nu + 1)}{\Gamma(\nu + 1 - \alpha)} x^{\nu-\alpha}.$$

1.3. Finite class of classical orthogonal polynomials

The solution of the differential equation

$$x(x+1)y_n''(x) + ((2-p)x + (1+q))y_n'(x) - n(n-1+p)y_n(x) = 0,$$

is the polynomial

$$(1.16) \quad M_n^{(p,q)}(x) = (-1)^n n! \sum_{k=0}^n \binom{p-n-1}{k} \binom{q+n}{n-k} (-x)^k.$$

With respect to the weight function $w_{p,q}(x) = x^q(1+x)^{-(p+q)}$, the polynomials defined in (1.16) are orthogonal on $[0, \infty)$ if and only if $p > 2n+1$ and $q > -1$. It is also known that the polynomials $M_n^{(p,q)}$ are related with hypergeometric functions as

$$(1.17) \quad M_n^{(p,q)}(x) = (-1)^n n! \binom{q+n}{n} {}_2F_1(-n, n+1-p; q+1; -x).$$

The Jacobi polynomials $P_n^{(\alpha,\beta)}$ and $M_n^{(p,q)}$ can be related by

$$M_n^{(p,q)}(x) = (-1)^n n! P_n^{(q, -p-q)}(2x+1) \Leftrightarrow P_n^{(p,q)}(x) = \frac{(-1)^n}{n!} M_n^{(-p-q,p)}\left(\frac{x-1}{2}\right).$$

Details related to this finite class of classical orthogonal polynomials can be found [18, 19, 21].

1.4. Outline of this work

The aim of this work is to investigate compositions of integral transforms (1.1) and (1.2) for Jacobi type finite class of classical orthogonal polynomials (1.16). In the sequel, the compositions are expressed in terms of the generalized Wright hypergeometric function ${}_p\psi_q(x)$ complex $a_i, b_j \in \mathbb{C}$, and real $\alpha_i, \beta_j \in \mathbb{R}$ ($i = 1, 2, \dots, p; j = 1, 2, \dots, q$) and defined by the series

$$(1.18) \quad {}_p\psi_q(x) = {}_p\psi_q \left[\begin{matrix} (a_i, \alpha_i)_{1,p} \\ (b_j, \beta_j)_{1,q} \end{matrix} \middle| x \right] = \sum_{k=0}^{\infty} \frac{\prod_{i=1}^p \Gamma(a_i + \alpha_i k)}{\prod_{j=1}^q \Gamma(b_j + \beta_j k)} \frac{x^k}{k!}.$$

Asymptotic behavior of this function for large values of argument of $z \in \mathbb{C}$ was studied in [7, 33, 34] and under the condition

$$(1.19) \quad \sum_{j=1}^q \beta_j - \sum_{i=1}^p \alpha_i > -1,$$

in [33, 34].

2. Composition with generalized fractional integrations

The following lemmas proved in [16], which are required to prove the result in this section.

Lemma 2.1 ([16]). *Let $\alpha, \beta, \eta \in \mathbb{C}$ be such that*

$$\operatorname{Re}(\alpha) > 0, \operatorname{Re}(\sigma) > \max[0, \operatorname{Re}(\beta - \eta)].$$

Then there exists the relation

$$\left(I_{0+}^{\alpha, \beta, \eta} t^{\sigma-1} \right) (x) = \frac{\Gamma(\sigma)\Gamma(\sigma + \eta - \beta)}{\Gamma(\sigma - \beta)\Gamma(\sigma + \alpha + \eta)} x^{\sigma - \beta - 1}.$$

In particular,

$$(2.1) \quad \left(I_{0+}^{\alpha} t^{\sigma-1} \right) (x) = \frac{\Gamma(\sigma)}{\Gamma(\sigma + \alpha)} x^{\sigma + \alpha - 1}, \quad \operatorname{Re}(\alpha) > 0, \operatorname{Re}(\sigma) > 0,$$

$$(2.2) \quad \left(I_{\eta, \alpha}^{+} t^{\sigma-1} \right) (x) = \frac{\Gamma(\sigma + \eta)}{\Gamma(\sigma + \alpha + \eta)} x^{\sigma + \alpha - 1}, \quad \operatorname{Re}(\alpha) > 0, \operatorname{Re}(\sigma) > -\operatorname{Re}(\eta).$$

Lemma 2.2 ([16]). *Let $\alpha, \beta, \eta \in \mathbb{C}$ be such that*

$$\operatorname{Re}(\alpha) > 0, \operatorname{Re}(\sigma) < 1 + \min[\operatorname{Re}(\beta), \operatorname{Re}(\eta)].$$

Then

$$\left(I_{-}^{\alpha, \beta, \eta} t^{\sigma-1} \right) (x) = \frac{\Gamma(\beta - \sigma + 1)\Gamma(\eta - \sigma + 1)}{\Gamma(1 - \sigma)\Gamma(\alpha + \beta + \eta - \sigma + 1)} x^{\sigma - \beta - 1}.$$

In particular,

$$(2.3) \quad \left(I_{-}^{\alpha} t^{\sigma-1} \right) (x) = \frac{\Gamma(1 - \alpha - \sigma)}{\Gamma(1 - \sigma)} x^{\sigma + \alpha - 1}, \quad 0 < \operatorname{Re}(\alpha) < 1 - \operatorname{Re}(\sigma)$$

$$(2.4) \quad \left(K_{\eta, \alpha}^{-} t^{\sigma-1} \right) (x) = \frac{\Gamma(\eta - \sigma + 1)}{\Gamma(\alpha + \eta - \sigma + 1)} x^{\sigma - 1}, \quad \operatorname{Re}(\sigma) < 1 + \operatorname{Re}(\eta)$$

Theorem 2.1. *Let $\alpha, \beta, \eta, \sigma \in \mathbb{C}$ be such that*

$$\operatorname{Re}(\alpha) > 0, \quad \text{and} \quad \operatorname{Re}(\sigma) > \max[0, \operatorname{Re}(\beta - \eta)].$$

Then,

$$\left(I_{0+}^{\alpha, \beta, \eta} t^{\sigma-1} M_n^{(p, q)}(t) \right) (x) = \frac{(-1)^n \Gamma(\sigma)\Gamma(\sigma + \eta - \beta)\Gamma(1 + q + n)x^{\sigma - \beta - 1}}{\Gamma(1 + q)\Gamma(\sigma - \beta)\Gamma(\sigma + \alpha + \eta)}$$

$$(2.5) \quad \times {}_4F_3 \left[\begin{matrix} 1+n-p, -n, \sigma, \sigma+\eta-\beta, \\ 1+q, \sigma-\beta, \sigma+\eta+\alpha, \end{matrix} \middle| -x \right].$$

Proof. First of all note that ${}_4F_3$ given in (2.5) exists because this series absolutely convergent and moreover, it is an entire function if $z \in \mathbb{C}$. Now using (1.1) and changing the order of the integration, the summation yields

$$\begin{aligned} & \left(I_{0+}^{\alpha, \beta, \eta} t^{\sigma-1} M_n^{(p, q)}(t) \right) (x) \\ &= \sum_{k=0}^{\infty} (-1)^k \binom{p-(n+1)}{k} \binom{q+n}{n-k} \left(I_{0+}^{\alpha, \beta, \eta} t^{\sigma+k-1} \right) (x) \end{aligned}$$

for any $k = 0, 1, 2, \dots, n$. Since

$$\operatorname{Re}(\sigma+k) > \operatorname{Re}(\sigma) > \max[0, \operatorname{Re}(\beta-\eta)],$$

applying Lemma 2.1 with σ replaced by $\sigma+k$ gives

$$(2.6) \quad \begin{aligned} & \left(I_{0+}^{\alpha, \beta, \eta} t^{\sigma-1} M_n^{(p, q)}(t) \right) (x) \\ &= \frac{(-1)^n \Gamma(1+q+n) x^{\sigma-\beta-1}}{\Gamma(1+n-p) \Gamma(-n)} \\ & \times \sum_{k=0}^{\infty} \frac{\Gamma(1+n-p+k) \Gamma(-n+k) \Gamma(\sigma+k) \Gamma(\sigma+\eta-\beta+k)}{\Gamma(1+q+k) \Gamma(\sigma-\beta+2k) \Gamma(\sigma+\alpha+\eta+k) \Gamma(1+k)} (-x)^k. \end{aligned}$$

Finally using the well-known identity $\Gamma(x+k) = (x)_k \Gamma(x)$, the right hand side of (2.6) can be rewrite as

$$\begin{aligned} \left(I_{0+}^{\alpha, \beta, \eta} t^{\sigma-1} M_n^{(p, q)}(t) \right) (x) &= \frac{(-1)^n \Gamma(1+q+n) \Gamma(\sigma) \Gamma(\sigma+\eta-\beta)}{\Gamma(1+q) \Gamma(\sigma-\beta) \Gamma(\sigma+\alpha+\eta)} x^{\sigma-\beta-1} \\ & \times \sum_{k=0}^{\infty} \frac{(-n)_k (1+n-p)_k (\sigma)_k (\sigma+\eta-\beta)_k}{(1+q)_k (\sigma-\beta)_k (\sigma+\alpha+\eta)_k k!} (-x)^k. \end{aligned}$$

This establish (2.5). \square

Substituting $\beta = -\alpha$ in Theorem 2.1, and using (2.1) gives the following result.

Corollary 2.1. *Let $\alpha, \sigma, \in \mathbb{C}$ be such that $\operatorname{Re}(\alpha) > 0, \operatorname{Re}(\sigma) > 0$. Then*

$$\begin{aligned} & \left(I_{0+}^{\alpha} t^{\sigma-1} M_n^{(p, q)}(t) \right) (x) \\ &= \frac{(-1)^n \Gamma(1+q+n) \Gamma(\sigma) x^{\sigma+\alpha-1}}{\Gamma(1+q) \Gamma(\sigma+\alpha)} {}_3F_2 \left[\begin{matrix} 1+n-p, -n, \sigma \\ 1+q, \sigma+\alpha \end{matrix} \middle| -x \right]. \end{aligned}$$

Substituting $\beta = 0$ in Theorem 2.1, and using (2.2) implies the following result.

Corollary 2.2. Let $\alpha, \eta, \sigma \in \mathbb{C}$ be such that $\operatorname{Re}(\alpha) > 0$, $\operatorname{Re}(\sigma) > -\operatorname{Re}(\eta)$. Then

$$\begin{aligned} & \left(I_{\eta, \alpha}^+ t^{\sigma-1} M_n^{(p, q)}(t) \right) (x) \\ &= \frac{(-1)^n \Gamma(1+q+n) \Gamma(\sigma+\eta) x^{\sigma-1}}{\Gamma(1+q) \Gamma(\sigma+\alpha+\eta)} {}_3F_2 \left[\begin{matrix} 1+n-p, -n, \sigma+\eta \\ 1+q, \sigma+\alpha+\eta \end{matrix} \middle| -x \right]. \end{aligned}$$

Next the generalized right-hand sided fractional integration (1.2) of the Jacobi type finite class of classical orthogonal polynomials related to Jacobi polynomials and the relation with generalized Wright functions are considered to get the following result.

Theorem 2.2. Let $\alpha, \eta, \sigma \in \mathbb{C}$ be such that

$$\operatorname{Re}(\alpha) > 0 \quad \text{and} \quad \operatorname{Re}(\sigma) < 1 + \min[\operatorname{Re}(\beta), \operatorname{Re}(\eta)].$$

Then

$$\begin{aligned} & \left(I_-^{\alpha, \beta, \eta} t^{\sigma-1} M_n^{(p, q)} \left(\frac{1}{t} \right) \right) (x) \\ &= \frac{\Gamma(1+q+n) \Gamma(\beta-\sigma+1) \Gamma(\eta-\sigma+1) x^{\sigma-\beta-1}}{\Gamma(1+q) \Gamma(1-\sigma) \Gamma(\alpha+\beta+\eta-\sigma+1)} \\ & \quad \times {}_4F_3 \left[\begin{matrix} 1+n-p, -n, \beta-\sigma+1, \eta-\sigma+1 \\ 1+q, 1-\sigma, \alpha+\beta+\eta-\sigma+1 \end{matrix} \middle| -\frac{1}{x} \right]. \end{aligned}$$

Proof. Now using (1.2) and changing the order of the integration, the summation yields

$$\begin{aligned} \left(I_-^{\alpha, \beta, \eta} t^{\sigma-1} M_n^{(p, q)} \left(\frac{1}{t} \right) \right) (x) &= (-1)^n \Gamma(n+1) \sum_{k=0}^{\infty} \binom{p-(n-1)}{k} \binom{q+n}{n-k} \\ & \quad \times \left((-1)^k I_-^{\alpha, \beta, \eta} t^{\sigma-k-1} \right) (x) \end{aligned}$$

for any $k = 0, 1, 2, \dots$

Since

$$\operatorname{Re}(\sigma - k - 1) \leq 1 + \operatorname{Re}(\sigma - 1) < 1 + \min[\operatorname{Re}(\beta), \operatorname{Re}(\eta)],$$

applying Lemma 2.2 with σ replaced by $\sigma - k$, it is easy to see that

$$\begin{aligned} & \left(I_-^{\alpha, \beta, \eta} t^{\sigma-1} M_n^{(p, q)} \left(\frac{1}{t} \right) \right) (x) \\ &= \frac{(-1)^n \Gamma(1+q+n) x^{\sigma-\beta-1}}{\Gamma(1+n-p) \Gamma(-n)} \\ & \quad \times \sum_{k=0}^{\infty} \frac{\Gamma(1+n-p+k) \Gamma(-n+k) \Gamma(\beta-\sigma+1+k) \Gamma(\eta-\sigma+1+k)}{\Gamma(1+q+k) \Gamma(1-\sigma+k) \Gamma(\alpha+\beta+\eta-\sigma+1+k) \Gamma(1+k)} (-x)^{-k}. \end{aligned}$$

Again the use of identity $\Gamma(x+k) = (x)_k \Gamma(x)$ in the above equation will give the required result. \square

Corollary 2.3. Let $\alpha, \eta, \sigma \in \mathbb{C}$ be such that

$$\operatorname{Re}(\alpha) > 0 \quad \text{and} \quad \operatorname{Re}(\sigma) < 1 + \min[\operatorname{Re}(-\alpha), \operatorname{Re}(\eta)]$$

and let $\sigma + \alpha \neq 1, 2, \dots$. Then

$$\begin{aligned} & \left(I_{-}^{\alpha} t^{\sigma-1} M_n^{(p,q)} \left(\frac{1}{t} \right) \right) (x) \\ &= \frac{\Gamma(1+q+n)\Gamma(-\alpha-\sigma+1)x^{\sigma+\alpha-1}}{\Gamma(1+q)\Gamma(1-\sigma)} {}_3F_2 \left[\begin{matrix} 1+n-p, -n, -\alpha-\sigma+1 \\ 1+q, 1-\sigma \end{matrix} \middle| -\frac{1}{x} \right]. \end{aligned}$$

Corollary 2.4. Let $\alpha, \eta, \sigma \in \mathbb{C}$ be such that

$$\operatorname{Re}(\alpha) > 0 \quad \text{and} \quad \operatorname{Re}(\sigma) < 1 + \min[0, \operatorname{Re}(\eta)]$$

and let $\sigma - \eta \neq 1, 2, \dots$. Then

$$\begin{aligned} \left(K_{\eta, \alpha}^{-} t^{\sigma-1} M_n^{(p,q)} \left(\frac{1}{t} \right) \right) (x) &= \frac{(-1)^n \Gamma(1+q+n)\Gamma(\eta-\sigma+1)x^{\sigma-1}}{\Gamma(1+q)\Gamma(\alpha+\eta-\sigma+1)} \\ &\quad \times {}_3F_2 \left[\begin{matrix} 1+n-p, -n, \eta-\sigma+1 \\ 1+q, \alpha+\eta-\sigma+1 \end{matrix} \middle| -\frac{1}{x} \right]. \end{aligned}$$

3. Composition with the generalized fractional derivatives

In this section, first we consider the composition of the finite class of classical orthogonal polynomials with the generalized fractional derivatives defined in (1.13) and the outcomes are represented in the closed form of the generalized Wright functions defined in (1.18).

Theorem 3.1. Let $\rho > 0$ and $\operatorname{Re}(\alpha) < 0$. Then for $p \geq 2n+1$ and $q > -1$,

(3.1)

$$\begin{aligned} & {}^{\rho}D_{0+}^{\alpha} \left(M_n^{(p,q)}(x) \right) \\ &= \frac{(-1)^n \rho^{\alpha-1} \Gamma(q+1+n)}{x^{\rho\alpha} \Gamma(-n) \Gamma(n+1-p)} {}_3\Psi_2 \left(\begin{matrix} (-n, 1) & (n+1-p, 1) & \left(1, \frac{1}{\rho}\right) \\ (q+1, 1) & \left(1-\alpha, \frac{1}{\rho}\right) \end{matrix} \middle| -x \right). \end{aligned}$$

In particular,

$$(3.2) \quad D_x^{\alpha} \left(M_n^{(p,q)}(x) \right) = \frac{(-1)^n (q+1)_n}{\Gamma(1-\alpha) x^{\alpha}} {}_3F_2(-n, n+1-p, 1; q+1, 1-\alpha; -x).$$

Proof. From (1.17) it follows that

$$M_n^{(p,q)}(x) = (-1)^n (q+1)_n \sum_{k=0}^{\infty} \frac{(-n)_k (n+1-p)_k}{(q+1)_k k!} (-x)^k.$$

Thus, an application of the operator ${}^{\rho}\mathcal{D}_{0+}^{\alpha}$ on both side of the above equation along with the relation (1.14) gives

$$\begin{aligned} {}^{\rho}\mathcal{D}_{0+}^{\alpha} \left(\mathbb{M}_n^{(p,q)}(x) \right) &= (-1)^n (q+1)_n \sum_{k=0}^{\infty} \frac{(-1)^k (-n)_k (n+1-p)_k}{(q+1)_k k!} {}^{\rho}\mathcal{D}_{0+}^{\alpha} (x^k) \\ (3.3) \quad &= (-1)^n (q+1)_n \rho^{\alpha-1} \sum_{k=0}^{\infty} \frac{(-1)^k (-n)_k (n+1-p)_k \Gamma\left(1 + \frac{k}{\rho}\right)}{(q+1)_k \Gamma\left(1 - \alpha + \frac{k}{\rho}\right) k!} x^{k-\alpha\rho}. \end{aligned}$$

Next the identity $\Gamma(x+k) = (x)_k \Gamma(x)$ yields

$$\begin{aligned} &{}^{\rho}\mathcal{D}_{0+}^{\alpha} \left(\mathbb{M}_n^{(p,q)}(x) \right) \\ &= \frac{(-1)^n \Gamma(n+1+q) \rho^{\alpha-1} x^{-\rho\alpha}}{\Gamma(-n) \Gamma(n+1-p)} \sum_{k=0}^{\infty} \frac{\Gamma(-n+k) \Gamma(n+1-p+k) \Gamma\left(1 + \frac{k}{\rho}\right)}{\Gamma(q+1+k) \Gamma\left(1 - \alpha + \frac{k}{\rho}\right) k!} (-x)^k \\ &= \frac{(-1)^n \Gamma(n+1+q) \rho^{\alpha-1} x^{-\rho\alpha}}{\Gamma(-n) \Gamma(n+1-p)} {}_3\Psi_2 \left(\begin{matrix} (-n, 1) & (n+1-p, 1) & \left(1, \frac{1}{\rho}\right) \\ (q+1, 1) & \left(1 - \alpha, \frac{1}{\rho}\right) \end{matrix} \middle| -x \right). \end{aligned}$$

Since ${}^1\mathcal{D}_{0+}^{\alpha} = \mathcal{D}_x^{\alpha}$, the relation (3.3) reduces

$$\mathcal{D}_x^{\alpha} \left(\mathbb{M}_n^{(p,q)}(x) \right) = (-1)^n (q+1)_n x^{-\alpha} \sum_{k=0}^{\infty} \frac{(-n)_k (n+1-p)_k \Gamma(1+k)}{(q+1)_k \Gamma(1-\alpha+k) k!} (-x)^k.$$

Finally, the result follows by the use of identity $\Gamma(x+k) = (x)_k \Gamma(x)$. \square

Corollary 3.1. For $\alpha = 1$, equations (3.2) becomes

$$(3.4) \quad \mathcal{D} \left(\mathbb{M}_n^{(p,q)}(x) \right) = \frac{(-1)^n n(1+n-p)(q+1)_n}{(1+q)} {}_2F_1[-(n-1), n+1-p; q+2; -x],$$

which is verified by [18].

In the next result, we will give an inequality for the composition of the finite class $\mathbb{M}_n^{(p,q)}$ with the generalized fractional derivatives defined in (1.9).

Theorem 3.2. Let $\mu, \nu, \sigma, x \in \mathbb{R}$ such that $\nu, \sigma, x > 0$ and $\mu < 0$. Then

$$(3.5) \quad \left| D_x^{\mu, \sigma} \left(x^{\nu-1} \mathbb{M}_n^{(p,q)}(x) \right) \right| \leq \frac{e^{-4\sigma} \Gamma(\nu) x^{\nu-\mu-1}}{\Gamma(\nu-\mu)} {}_3F_2(-n, 1+n-p, \nu; q+1, \nu-\mu; x).$$

Proof. From (1.11) and (1.17), it follows that

$$(3.6) \quad \begin{aligned} &D_x^{\mu, \sigma} \left(x^{\nu-1} \mathbb{M}_n^{(p,q)}(x) \right) \\ &= \frac{(-1)^n (q+1)_n x^{\nu+\mu-1}}{\Gamma(-\mu)} \sum_{n=0}^{\infty} \frac{(-n)_k (1+n-p)_k}{(q+1)_k k!} \mathbb{B}_{\sigma}(\nu+k, -\mu) x^k. \end{aligned}$$

It is proved in [5] that $|\mathbf{B}_\sigma(x, y)| \leq e^{-4\sigma} \mathbf{B}(x, y)$ for $x, y, \sigma \geq 0$ and $\mathbf{B}(x, y)$ is the classical beta functions. Using this to (3.6), it follows that

$$(3.7) \quad \left| D_x^{\mu, \sigma} \left(x^{\nu-1} \mathbf{M}_n^{(p, q)}(x) \right) \right| \leq \frac{e^{-4\sigma} (q+1)_n x^{\nu+\mu-1}}{\Gamma(-\mu)} \sum_{n=0}^{\infty} \left| \frac{(-n)_k (1+n-p)_k}{(q+1)_k k!} \right| \mathbf{B}(\nu+k, -\mu) x^k.$$

Now $p \geq 2n+1$ implies $1+n-p \leq -n$ and thus, $(-n)_k (1+n-p)_k$ is non-negative for all k . Thus, we can rewrite (3.7) as

$$\left| D_x^{\mu, \sigma} \left(x^{\nu-1} \mathbf{M}_n^{(p, q)}(x) \right) \right| \leq \frac{e^{-4\sigma} (q+1)_n x^{\nu+\mu-1}}{\Gamma(-\mu)} \sum_{n=0}^{\infty} \frac{(-n)_k (1+n-p)_k}{(q+1)_k k!} \mathbf{B}(\nu+k, -\mu) x^k,$$

which is equivalent to (3.5) after using the identity

$$\mathbf{B}(\nu+k, -\mu) = \frac{\Gamma(\nu+k)\Gamma(-\mu)}{\Gamma(\nu-\mu+k)} = \frac{\Gamma(\nu)(\nu)_k}{\Gamma(\nu-\mu)(\nu-\mu)_k}.$$

This prove the inequality. \square

Remark 3.1. For $\sigma = 0$ and $\nu = 1$, we have from Theorem 3.2 that

$$\left| D_x^\mu \left(\mathbf{M}_n^{(p, q)}(x) \right) \right| \leq \frac{x^{-\mu}}{\Gamma(1-\mu)} {}_3F_2(-n, 1+n-p, 1; q+1, 1-\mu; x).$$

References

- [1] P. Agarwal and J. Choi, *Fractional calculus operators and their image formulas*, J. Korean Math. Soc. **53** (2016), no. 5, 1183–1210.
- [2] M. Ali Özarslan and E. Özergin, *Some generating relations for extended hypergeometric functions via generalized fractional derivative operator*, Math. Comput. Modelling **52** (2010), no. 9-10, 1825–1833.
- [3] R. Caponetto, *Fractional Order Systems: Modeling and Control Applications*, World Scientific, 2010.
- [4] M. Caputo, *Elasticitae dissipazione Zanichelli*, Bologna, 1969.
- [5] M. A. Chaudhry, A. Qadir, M. Rafique, and S. M. Zubair, *Extension of Euler's beta function*, J. Comput. Appl. Math. **78** (1997), no. 1, 19–32.
- [6] M. A. Chaudhry, A. Qadir, H. M. Srivastave, and R. B. Paris, *Extended hypergeometric and confluent hypergeometric functions*, Appl. Math. Comput. **159** (2004), no. 2, 589–602.
- [7] C. Fox, *The asymptotic expansion of generalized hypergeometric functions*, Proc. London. Math. Soc. **27** (1938), no. 1, 389–400.
- [8] R. Hilfer (ed.), *Applications of Fractional Calculus in Physics*, World Scientific Publishing Co., Singapore, New York, 2000.
- [9] S. L. Kalla and R. K. Saxena, *Integral operators involving hypergeometric functions*, Math. Z. **108** (1969), 231–234.
- [10] U. N. Katugampola, *New approach to a generalized fractional integral*, Appl. Math. Comput. **218** (2011), no. 3, 860–865.

- [11] V. S. Kiryakova, *Generalized Fractional Calculus and Applications*, Pitman Res Notes Math. 301, Longman Scientific & Technical; Harlow, Co-published with John Wiley, New York, 1994.
- [12] ———, *All the special functions are fractional differintegrals of elementary functions*, J. Phys. A **30** (1997), no. 14, 5083–5103.
- [13] ———, *Multiple (multi-index) Mittag-Leffler functions and relations to generalized fractional calculus*, J. Comput. Appl. Math. **118** (2000), no. 1-2, 241–259.
- [14] H. Kober, *On fractional integrals and derivatives*, Quart. J. Math., Oxford Ser. **11** (1940), 193–212.
- [15] A. A. Kilbas and M. Saigo, *H-transforms*, Chapman & Hall/CRC, Boca Raton, FL, 2004.
- [16] A. A. Kilbas and N. Sebastian, *Generalized fractional integration of Bessel function of the first kind*, Integral Transforms Spec. Funct. **19** (2008), no. 11-12, 869–883.
- [17] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies 204, Elsevier, Amsterdam, 2006.
- [18] P. Malik and A. Swaminathan, *Derivatives of a finite class of orthogonal polynomials defined on positive real line related to F-Distribution*, Comput. Math. Appl. **61** (2011), no. 4, 1180–1189.
- [19] ———, *Derivatives of a finite class of orthogonal polynomials related to inverse gamma distribution*, Appl. Math. Comput. **218** (2012), no. 11, 6251–6262.
- [20] P. Malik, S. R. Mondal, and A. Swaminathan, *Riemann-Liouville Fractional calculus for generalized Bessel functions*, Proceedings of the ASME 2011, International Design Engineering Technical Conferences and Computers and Information in Engineering Conference **3** (2011), 409–418.
- [21] M. Masjedjamei, *Three finite classes of hypergeometric orthogonal polynomials and their application in functions approximation*, Integral Transforms Spec. Funct. **13** (2002), no. 2, 169–191.
- [22] A. C. McBride and G. F. Roach (Editors), *Fractional Calculus*, (University of Strathclyde, Glasgow, Scotland, August 5–11, 1984) Research Notes in Mathematics 138, Pitman Publishing Limited, London, 1985.
- [23] K. S. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, Wiley and Sons, New York, 1993.
- [24] Saiful. R. Mondal and K. S. Nisar, *Marichev-Saigo-Maeda fractional integration operators of generalized Bessel functions*, Mathematical Problems in Engineering **2014** (2014), Article ID 274093, 11 pages.
- [25] K. Nishimoto, *Fractional Calculus 1, 2, 3, 4, 5*, Descartes Press, Koriyama, Japan, 1984, 1987, 1989, 1991, 1996.
- [26] K. Nishimoto (Editor), *Fractional Calculus and Its Applications*, (May 29-June 1, 1989), Nihon University (College of Engineering), Koriyama, 1990.
- [27] K. B. Oldham and J. Spanier, *The Fractional Calculus: Theory and Applications of Differentiation and Integration of Arbitrary Order*, Academic Press, New York, 1974.
- [28] E. D. Rainville, *Special Functions*, Macmillan, New York, 1960.
- [29] M. Saigo, *A remark on integral operators involving the Gauss hypergeometric functions*, Math. Rep. Kyushu Univ. **11** (1977/78), no. 2, 135–143.
- [30] M. Saigo and A. A. Kilbas, *Generalized fractional calculus of the H function*, Fukuoka Univ. Sci. Rep. **29** (1999), no. 1, 31–45.
- [31] M. Saigo and N. Maeda, *More generalization of fractional calculus*, Transform methods & special functions, Varna '96, 386–400, Bulgarian Acad. Sci., Sofia, 1998.
- [32] S. G. Samko, A. A. Kilbas, and O. I. Marichev, *Fractional integrals and derivatives*, translated from the 1987 Russian original, Gordon and Breach, Yverdon, 1993.

- [33] E. M. Wright, *The asymptotic expansion of integral functions defined by Taylor series*, Philos. Trans. Roy. Soc. London, Ser. A. **238** (1940), 423–451.
- [34] ———, *The asymptotic expansion of the generalized hypergeometric function*, Proc. London Math. Soc. (2) **46** (1940), 389–408.

PRADEEP MALIK
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF PETROLEUM AND ENERGY STUDIES
DEHRADUN, UTTARAKHAND, INDIA
E-mail address: `pmalik@ddn.upes.ac.in`

SAIFUL R. MONDAL
DEPARTMENT OF MATHEMATICS AND STATISTICS
COLLEGE OF SCIENCE
KING FAISAL UNIVERSITY
AL-HASA, HOFUF 31982, SAUDI ARABIA
E-mail address: `smondal@kfu.edu.sa`