

THE HULL NUMBER OF POWERS OF CYCLES

HASAN AL-EZEH, MANAL GHANEM, AND JAMEEL RWALAH

ABSTRACT. Let C_n be the cycle graph of order n on the vertices v_0, v_1, \dots, v_n and C_n^k be the k -th power of C_n . In this article we determine the hull-number of C_n^k .

1. Introduction

Given a finite simple connected graph G , let u and v be two vertices of G . The distance between u and v is the length of a shortest path between u and v , we denote it by $d_G(u, v)$. A shortest path between u and v is called a $u - v$ geodesic. The set of all vertices in G that lie on a $u - v$ geodesic is denoted by $I[u, v]$. The closed interval $I[u, v]$ consists of all vertices that lie on a uv geodesic of G . For $A \subseteq V(G)$, let the closed interval $I[A]$ be the union of all sets $I[u, v]$ for $u, v \in A$, then A is called a convex set if $I[A] = A$. The convex hull of A , denoted by $[A]$, is the smallest convex set containing A . If $[A] = V(G)$, then A is called a hull set of G . The cardinality of a minimum hull set of G is called the hull number of G , and it is denoted by $h(G)$. If $I[A] = V(G)$, then A is called a geodetic set of G . The minimum cardinality of a geodetic set in G is named the geodetic number of G and it is denoted by $g(G)$. Certainly, $h(G) \leq g(G)$.

The process of rebuilding a network modelled by a connected graph is a discrete optimization problem, consisting in finding a subset of vertices of cardinality as small as possible, which would allow us to store and retrieve the whole graph. One way to approach this problem is by using a certain convex operator. This procedure has attracted much attention since it was shown in [9] that every convex subset in a graph is the convex hull of its extreme vertices if and only if the graph is chordal and contains no induced 3-fan. The hull number of a graph was introduced by [8]. They characterized graphs having some particular hull numbers and they obtained a number of bounds for the hull numbers of graphs. Dourado et al. [7] proved that the hull number of

Received December 6, 2016; Accepted June 14, 2017.

2010 *Mathematics Subject Classification.* 05C12, 05C38.

Key words and phrases. geodetic number, hull number, paths, powers of cycles graphs.

This paper constitutes a part of PhD Thesis at the University of Jordan written by the third author and supervised by the first and the second authors.

unit interval graphs, cographs and split graphs can be computed in polynomial time. The hull number of an oriented graph was studied in [5] and [6]. The hull number of the power of paths was determined in [1]. For more results on the subject, see [3], [4] and [5]. For any positive integer k and a connected graph G the k -th power graph G^k of G has $V(G^k) = V(G)$ and the distinct vertices x and y are adjacent in G^k if $d(x, y) \leq k$. Circulant graphs have been extensively studied and have a vast number of applications to multicomputer networks and distributed computation (see [2] and [10]). One type of the circulant graphs is the k -th power of the n -cycle C_n^k . Our aim in this paper is to find the hull number of the graph C_n^k .

2. The hull number of C_n^k

For positive integers n and k , we denote by C_n^k the graph with vertex set $\{v_0, v_1, \dots, v_{n-1}\}$ and edge set $\{v_i v_j : i - j \equiv \pm m \pmod{n}, 1 \leq m \leq k\}$. The graph C_n^k is the k -th power of the n -cycle C_n .

In this section, we will determine the hull number of the k -th power of the n -cycle C_n^k . The hull number of a connected graph G of order n is n if and only if G is the complete graph of order n , [8]. It is clear that C_n^k is the complete graph of order n when $k \geq \lfloor \frac{n}{2} \rfloor$ and hence its hull number equal n . So next we will only consider C_n^k when $1 < k < \lfloor \frac{n}{2} \rfloor$.

Let G be a graph. Given a vertex v , denote by $N(v)$ the set of neighbors of v . And denote, the subgraph of G induced by the set $B, B \subseteq V(G)$ by $G[B]$. We say that v is a simplicial vertex of G if $N(v)$ induces a complete subgraph. It is clear that every hull set of a graph G contains the set of all simplicial vertices of G . In this section, we characterize the hull number of the graph C_n^k in the following sequences of lemmas.

First, we start by the following lemma that determines the geodetic number of the k -th power of a path with $qk + 2$ vertices.

Lemma 1 ([1]). *Let P_{n+1} be the path of order $n+1$ and P_{n+1}^k be the k -th power of P_{n+1} . Suppose that $n = qk + r$ where q is a positive integer and $0 \leq r < k$, then $g(P_{n+1}^k) = 2$ if and only if $n = qk + 1$.*

In the following lemma, we show that 3 is an upper bound for $h(C_n^k)$.

Lemma 2. *The hull number of any power of cycle graph is at most 3.*

Proof. Let C_n^k be the k -th power of the cycle graph with n vertices. Then there are two cases of n .

Case 1: n is even. Use division algorithm, to write $n = 2qk + 2r$, where q is a positive integer, $0 \leq 2r < 2k$, and $V(C_n^k) = \{v_0, v_1, \dots, v_k, \dots, v_{qk}, \dots, v_{qk+r}, \dots, v_{qk+2r}, \dots, v_{(q+1)k+2r}, \dots, v_{(2q-1)k+2r}, \dots, v_{2qk+2r-1}\}$. Now, assume that the hull number of C_n^k is not equal 2 and let $A = \{v_0, v_1, v_{qk+r}\}$, we claim that A is a hull-set of C_n^k . To show this, first note that $\{v_0, v_{qk+r}\} \subseteq A$ gives $A_1 = \{v_0, v_k, v_{2k}, \dots, v_{qk}, v_{qk+r}\}$ and $A_2 = \{v_{qk+r}, v_{qk+2r}, v_{(q+1)k+2r}, \dots$

$v_{(2q-1)k+2r}, v_0\}$ are subsets of $[A]$, since their elements lie on geodesics between v_0 and v_{qk+r} . And observe that, $\{v_1, v_{qk}, v_{qk+2r}\} \subseteq [A]$ and the induced subgraph of C_n^k with the set of vertices $A_3 = \{v_1, v_0, v_{2qk+2r-1}, \dots, v_{(q+1)k+2r}, \dots, v_{qk+2r}\}$ is isomorphic to P_{m+1}^k , where $m = qk + 1$. Use Lemma 1, to get $A_3 \subseteq [A]$. Therefore, $v_{2qk+2r-1} \in [A]$. Take $A_4 = \{v_{2qk+2r-1}, v_0, \dots, v_k, \dots, v_{qk}\}$, then $C_n^k[A_4] \cong P_{qk+2}^k$, and hence $A_4 \subseteq [A]$. Moreover,

$$A_5 = \{v_{(q+1)k+r+1}, v_{(q+1)k+r}, \dots, v_{qk+2r}, \dots, v_{qk+r}\} \subseteq [A] \text{ and}$$

$$A_6 = \{v_{qk+r}, v_{qk+r-1}, \dots, v_{qk}, \dots, v_{(q-1)k+r}, v_{(q-1)k+r-1}\} \subseteq [A],$$

since $\{v_{(q+1)k+r+1}, v_{qk+r}, v_{(q-1)k+r-1}\} \subseteq [A]$ and $C_n^k[A_i] \cong P_{k+2}^k$ for $i = 5, 6$. So, $V(C_n^k) = \bigcup_{i=1}^6 A_i = [A]$ and hence A is a hull set of C_n^k .

Case 2: n is odd. By division algorithm $n - 1 = 2qk + 2r$, where q is a positive integer, $0 \leq 2r < 2k$ and $V(C_n^k) = \{v_0, v_1, \dots, v_k, \dots, v_{qk}, \dots, v_{qk+r}, \dots, v_{qk+2r+1}, \dots, v_{(q+1)k+2r+1}, \dots, v_{(2q-1)k+2r+1}, \dots, v_{2qk+2r}\}$. If $r = 0$, then $A = \{v_0, v_{(q-1)k}, v_{qk+1}\}$ is a hull set of C_n^k . To show this set $A_1 = \{v_{(q-1)k}, v_{(q-1)k+1}, \dots, v_{qk+1}\}$, $A_2 = \{v_0, v_1, \dots, v_{(q-1)k+1}\}$, $A_3 = \{v_{qk+1}, v_{(q+1)k+1}, \dots, v_{(2q-1)k+1}, v_0\}$, $A_4 = \{v_{qk}, v_{qk+1}, \dots, v_{(q+1)k+1}\}$, $A_5 = \{v_{(q+1)k+1}, \dots, v_{(2q-1)k+1}, \dots, v_0, v_1\}$. By using Lemma 1 and noting that all vertices of A_3 lie on a $v_0 - v_{qk+1}$ geodesic, we can prove respectively that $A_i \subseteq [A]$ for all i . Hence, $V(C_n^k) = \bigcup_{i=1}^5 A_i = [A]$ and thus $h(C_n^k) = 3$. Now, suppose that $r \neq 0$. We claim that $A = \{v_0, v_1, v_{qk+r}\}$ is a hull set of C_n^k , to prove this claim we mimic the proof of Case 1. First, observe that, there are two paths between v_0 and v_{qk+r} , the first one is $v_0 - v_1 - \dots - v_k - \dots - v_{qk} - \dots - v_{qk+r}$ and the second is $v_{qk+r} - \dots - v_{qk+2r+1} - \dots - v_{(q+1)k+2r+1} - \dots - v_{(2q-1)k+2r+1} - \dots - v_0$. Since the length of the first path is $qk + r$ and the length of the second path is $qk + r + 1$ and $r + 1 \leq k$, we have $v_0 - v_k - \dots - v_{qk} - v_{qk+r}$ and $v_{qk+r} - v_{qk+2r+1} - v_{(q+1)k+2r+1} - \dots - v_{(2q-1)k+2r+1} - v_0$ are geodesics between v_0 and v_{qk+r} . Thus, $A_1 = \{v_0, v_k, \dots, v_{qk}, v_{qk+r}, v_{qk+2r+1}, v_{(q+1)k+2r+1}, \dots, v_{(2q-1)k+2r+1}\} \subseteq [A]$. Now, take

$$A_2 = \{v_{qk+2r+1}, v_{qk+2r+2}, \dots, v_{(q+1)k+r+1}, \dots, v_{2qk+2r}, v_0, v_1\},$$

$$A_3 = \{v_{2qk+2r}, v_0, \dots, v_k, \dots, v_{qk}\},$$

$$A_4 = \{v_{qk+r}, v_{qk+r+1}, \dots, v_{qk+2r}, \dots, v_{(q+1)k+r}, v_{(q+1)k+r+1}\},$$

$$A_5 = \{v_{(q-1)k+r-1}, v_{(q-1)k+r}, \dots, v_{qk}, v_{qk+1}, \dots, v_{qk+r}\}.$$

Then $C_n^k[A_i] \cong P_{qk+2}^k$ for $i = 2, 3$ and $C_n^k[A_i] \cong P_{k+2}^k$ for $i = 4, 5$. By Lemma 1, we get $\bigcup_{i=2}^5 A_i \subseteq [A]$. So, $V(C_n^k) = [A]$ and hence A is a hull set of C_n^k . \square

In the following lemma, we show that 3 is a sharp upper bound of C_n^k .

Lemma 3. *Suppose that $n = 2qk$ where q is a positive integer, then $h(C_n^k) = 3$.*

Proof. Suppose that $A = \{v_0, v_{qk}\}$, we will show that A is not a hull set of C_n^k . Observe that, there are exactly two $v_0 - v_{qk}$ geodesics, the first one is

$v_0 - v_k - \dots - v_{qk}$ and the second is $v_{qk} - v_{(q+1)k} - v_{(q+2)k} - \dots - v_{(2q-1)k} - v_0$. So, $[A] \neq V(C_n^k)$ and hence A is not a hull set of C_n^k . Similarly, if we replace v_{qk} in A by any other vertex of C_n^k we can easily show that A is not a hull set. So $h(C_n^k) > 2$. By Lemma 2, we get the result \square

Lemma 4 ([1]). *Suppose that $n = qk + r$ where $0 < r < k$, then*

$$h(P_{n+1}^k) = \begin{cases} 2, & \text{if } q > 1, r \neq k; \\ 3, & \text{if } q = 1, r \neq 1; \\ 2, & \text{if } q = 1, r = 1. \end{cases}$$

Lemma 5. *Suppose that $n = 2qk + 2r$ where q is a positive integer and $0 < r < k$, then $h(C_n^k) = 2$.*

Proof. Let

$$A_1 = \{v_0, v_1, \dots, v_k, \dots, v_{qk}, \dots, v_{qk+r}\} \text{ and}$$

$$A_2 = \{v_{qk+r}, \dots, v_{qk+2r}, \dots, v_{(q+1)k+2r}, \dots, v_{2qk+2r-1}, v_0\}.$$

Then $C_n^k[A_i] \cong P_{qk+r+1}^k$ for $i = 1, 2$. By using Lemma 4, we have the following three cases:

Case 1: $q = 1$ and $r = 1$. Since v_0 and v_{qk+r} are simplicial vertices of $C_n^k[A_i]$, the hull set of $C_n^k[A_i]$ is $A = \{v_0, v_{qk+r}\}$ for $i = 1, 2$. But $A_1 \cup A_2 = V(C_n^k)$, so A is a hull set of C_n^k .

Case 2: $q > 1$ and $r \neq k$, then the hull set of $C_n^k[A_i]$ is $A = \{v_0, v_{qk+r}\}$ for $i = 1, 2$. Thus A is a hull set of C_n^k .

Case 3: $q = 1$ and $r \neq 1$. Then the hull number $h(C_n^k[A_i]) = 3$ for $i = 1, 2$. In this case, $A = \{v_0, v_{k+r}\}$ is a hull set of C_n^k . To show this, observe that $v_0 - v_k - v_{k+r}$ and $v_{k+r} - v_{2k+r} - v_0$ are $v_0 - v_{k+r}$ geodesics. Therefore, $\{v_{2k+r}, v_k\} \subseteq [A]$. Since $r < k$, the path $v_{2k+r} - v_1 - v_k$ is a $v_{2k+r} - v_k$ geodesic. So, v_1 belongs to $[A]$ and hence $\{v_0, v_1, v_{k+r}\} \subseteq [A]$. By using the proof of Lemma 2, we have A is a hull set of C_n^k . \square

Lemma 6. *Suppose that $n-1 = 2qk$ where q is a positive integer, then $h(C_n^k) = 3$.*

Proof. Assume that $n-1 = 2qk$, where q is a positive integer, that means the number of the vertices of the graph C_n^k is odd. Set $A = \{v_0, v_{qk}\}$. Clearly, there exists unique $v_0 - v_{qk}$ geodesic which is $v_0 - v_k - \dots - v_{qk}$. So, A is not a hull set of C_n^k . Similarly, if we replace v_{qk} by any other vertex, we get the same result. By Lemma 2, we conclude that $h(C_n^k) = 3$. \square

Lemma 7. *Suppose that $n-1 = 2qk + 2r$ where q is a positive integer and $0 < r < k$, then $h(C_n^k) = 2$.*

Proof. Let $A = \{v_0, v_{qk+r}\}$. Then A is a hull set of C_n^k . To show this it is enough to show that v_1 belongs to $[A]$ (see the proof of Lemma 2). Since $v_0 - v_k - v_{k+r} - v_{2k+r} - \dots - v_{qk+r}, v_0 - v_r - v_{k+r} - v_{2k+r} - \dots - v_{qk+r}$ and $v_0 - v_{(2q-1)k+2r+1} - v_{(2q-2)k+2r+1} - \dots - v_{qk+2r+1} - v_{qk+r}$ are $v_0 -$

v_{qk+r} geodesics, we have $\{v_{(2q-1)k+2r+1}, v_r, v_k\} \subseteq [A]$. But $v_{(2q-1)k+2r+1} - v_{2qk+2r} - v_r$ is a $v_{(2q-1)k+2r+1} - v_r$ geodesic, so $v_{2qk+2r} \in [A]$. Now, let $B = \{v_{2qk+2r}, v_0, \dots, v_k\}$, then $C_n^k[B] \cong P_{k+2}^k$. By Lemma 1, we get $v_1 \in [A]$ and hence the result holds. \square

We can summarize the above in the following theorem.

Theorem 8. *If $n = 2qk + 2r$ or $n - 1 = 2qk + 2r$ where q is a positive integer and $0 \leq r < k$, then*

$$h(C_n^k) = \begin{cases} 2, & \text{if } 0 < r < k; \\ 3, & \text{if } r = 0. \end{cases}$$

For interested readers one might try to find the hull number of some other types of circulant graphs.

References

- [1] O. AbuGhneim, B. Al-Khamaiseh, and H. Al-Ezeh, *The geodetic, hull, and Steiner numbers of powers of paths*, Util. Math. **95** (2014), 289–294.
- [2] J. Bermond, F. Cormellas, and D. F. Hsu, *Distributed loop computer networks: a survey*, J. Parallel Distributed Computing **24** (1995), 2–10.
- [3] J. Cáceresa, C. Hernandob, M. Morab, I. M. Pelayob, and M. L. Puertasa, *On the geodetic and the hull numbers in strong product graphs*, Comput. Math. Appl. **60** (2010), no. 11, 3002–3031.
- [4] S. R. Canoy, G. B. Cagaanan, and S. V. Gervacio, *Convexity, geodetic and hull numbers of the join of graphs*, Utilitas Math. **71** (2006), 143–159.
- [5] G. Chartrand, J. F. Fink, and P. Zhang, *Convexity in oriented graphs*, Discrete Appl. Math. **116** (2002), 115–126.
- [6] ———, *The hull number of an oriented graph*, International J. Math. Math. Sci. **36** (2003), 2265–2275.
- [7] M. C. Durado, J. G. Gimbel, J. Kratochvíl, F. Protti, and J. L. Szwarcfiter, *On the computation of the hull number of a graph*, Discrete Math. **309** (2009), no. 18, 5668–5674.
- [8] M. G. Everett and S. B. Seidman, *The hull number of a graph*, Discrete Math. **57** (1985), no. 3, 217–223.
- [9] M. Farber and R. E. Jamison, *Convexity in graphs and hypergraphs*, SIAM J. Algebraic Discrete Methods **7** (1986), no. 3, 433–444.
- [10] D. E. Knuth, *The Art of Computer Programming. Vol. 3*, Addison-Wesley, Reading, MA, 1975.

HASAN AL-EZEH
 DEPARTMENT OF MATHEMATICS
 SCHOOL OF SCIENCE
 THE UNIVERSITY OF JORDAN AMMAN 11942, JORDAN
 E-mail address: alezehh@ju.edu.jo

MANAL GHANEM
DEPARTMENT OF MATHEMATICS
SCHOOL OF SCIENCE
THE UNIVERSITY OF JORDAN AMMAN 11942, JORDAN
E-mail address: m.ghanem@ju.edu.jo

JAMEEL RWALAH
DEPARTMENT OF MATHEMATICS
SCHOOL OF SCIENCE
THE UNIVERSITY OF JORDAN AMMAN 11942, JORDAN
E-mail address: jameel.31@yahoo.com

Ahead of Print