

AN EXTENSION OF RANDOM SUMMATIONS OF INDEPENDENT AND IDENTICALLY DISTRIBUTED RANDOM VARIABLES

LE TRUONG GIANG AND TRAN LOC HUNG

ABSTRACT. The main goal of this paper is to study an extension of random summations of independent and identically distributed random variables when the number of summands in random summation is a partial sum of n independent, identically distributed, non-negative integer-valued random variables. Some characterizations of random summations are considered. The central limit theorems and weak law of large numbers for extended random summations are established. Some weak limit theorems related to geometric random sums, binomial random sums and negative-binomial random sums are also investigated as asymptotic behaviors of extended random summations.

1. Introduction

Let X, X_1, X_2, \dots be a sequence of independent, identically distributed (i.i.d.) random variables, having common distribution F_X , mean $E(X) = \mu$, and finite variance $D(X) = \sigma^2 < +\infty$. Suppose that Y, Y_1, Y_2, \dots, Y_n are independent, identically distributed, non-negative, integer-valued random variables, with mean $E(Y) = \alpha$, and finite variance $D(Y) = \tau^2 < +\infty$. Additionally, suppose that the random variables X, X_1, X_2, \dots and Y, Y_1, Y_2, \dots, Y_n are independent. Set, for $n \geq 1$, $N_n := Y_1 + Y_2 + \dots + Y_n$ and define $S_{N_n} := \sum_{j=1}^{N_n} X_j$. Then, the random summation S_{N_n} is a remarkable random summation because of the number of summands in random summation is a partial sum of i.i.d. random variables. It is obvious that when $P(N_1 = Y_1) = 1$, the random summation will be returned to the classical random sum $S_{Y_1} = \sum_{j=1}^{Y_1} X_j$.

Since the appearance of the Robbin's results in 1948 (see [21] for more details), the random summations have been investigated in the theory probability, statistics, stochastic processes and various related fields for quite some time by Robbins (1948), Feller (1971), Gnedenko (1972), Gnedenko and Korolev (1996), Renyi (1970), Korolev and Kruglov (1990), Gut (2005), Hung

Received March 19, 2017; Revised August 27, 2017; Accepted February 1, 2018.

2010 *Mathematics Subject Classification.* Primary 60E10; Secondary 60F05.

Key words and phrases. random summation, central limit theorem, weak law of large numbers, gamma distribution, Laplace distribution.

and Thanh (2010), Hung et al. (2008), Durrett (1977), Hu and Cheng (2012), Chen and Shao (2007), Chen and Goldstein (2011), Omev and Vesilob (2015), . . . (see [2–9, 15, 17, 19–21], and the references therein).

Recently, a number of results related to the specific cases of random summations like geometric random sums, compound Poisson sums, binomial random sums, negative-binomial random sums, etc. and their applications have been investigated by many authors as Gnedenko (1971), Gnedenko and Korolev (1996), Kruglov and Korolev (1990), Kalashnikov (1997), Vellaisamy and Chaudhuri (1996), Hung et al. (2008, 2010), Sunklodas (2009, 2014, 2015), . . . Results of this nature may be found in [7, 10, 11, 13, 17, 22–25] and the references given there.

It makes sense to consider an extension of random summation when N_n , $n \geq 1$, is being a partial sum of independent, identically distributed non-negative integer-valued random variables. This extension comes from the actual requirements, e.g. a negative-binomial random sum is an extension of geometric random sum. Actually, suppose that Y_j , $j = 1, 2, \dots, n$ are n independent, geometric distributed random variables with parameter $p \in (0, 1)$. Then, the sum $S_{Y_1} = X_1 + X_2 + \dots + X_{Y_1}$ is said to be a geometric random sum. Obviously, the sum $N_n = Y_1 + Y_2 + \dots + Y_n$ will be a negative-binomial random variable with parameters n and p ($n \geq 1, p \in (0, 1)$), and the random sum S_{N_n} will be an extension of the S_{Y_1} , and we will call it by negative-binomial random summation (see Remark 2.2 in next Section and Theorems 3.4, 3.5 and 3.6 in last Section).

As far as we know, up to the present there is just a little number of the results concerning with convergence rates in limit theorems for an extension of random summations have been discussed like by Chen and Shao (2007) in [3] for case of independent and identically distributed random variables and by Islack (2013) in case of m -dependent and identically distributed random variables (see [12] for more details).

The main purpose of this paper is to investigate the characterizations and asymptotic behaviors of the extended random summations of i.i.d. random variables when the numbers of random summation is a partial sum of n i.i.d. non-negative integer-valued random variables. Some limit theorems for extended random summations like central limit theorem and weak law of large numbers are re-established. Moreover, some limit theorems for binomial random sum (Corollary 3.1), geometric random sums (Corollary 3.2 and Corollary 3.4) and negative-binomial random sums (Theorems 3.3, 3.4 and 3.5) also considered. The received results are extensions of the known results (see [1, 4–10, 16–18, 21–24] and [14]).

The organization of this paper is as follows. Section 2 is devoted to the discussion on some characterizations of random summation as mean, variance, generating function and characteristic function. Section 3 gives some results on asymptotic behaviors of normalization of the extended random summations.

2. Characterizations of random summation

Throughout this paper, we shall denote by $\psi(t)$ and $\varphi(t)$ the generating function and characteristic function of random variables, respectively. We start to consider some characterizations of the random summation S_{N_n} with two following propositions.

Proposition 2.1. *Let X, X_1, X_2, \dots be a sequence of i.i.d. random variables with common distribution F_X , mean $E(X) = \mu$, and finite variance $D(X) = \sigma^2 < +\infty$. Suppose that Y, Y_1, Y_2, \dots, Y_n are i.i.d. non-negative integer-valued random variables with mean $E(Y) = \alpha$, and finite variance $D(Y) = \tau^2 < +\infty$. Additionally, assume that random variables X, X_1, X_2, \dots and Y, Y_1, Y_2, \dots, Y_n are independent. Define*

$$(1) \quad S_{N_n} := \sum_{k=1}^{N_n} X_k,$$

where $N_n = Y_1 + Y_2 + \dots + Y_n$. Then,

- (1) *Probability distribution function of random summation S_{N_n} in (1) is defined in form*

$$\begin{aligned} F_{S_{N_n}}(x) &= P(S_{N_n} \leq x) = \sum_{k=1}^{\infty} P(N_n = k) \times P(S_k \leq x) \\ &= \sum_{k=1}^{\infty} P(N_n = k) \times F_X^{*k}(x), \end{aligned}$$

where $F_X^{*k}(x)$ denotes the k -th convolution power of the distribution function $F_X(x)$.

- (2) *The Wald's identity for random summation S_{N_n} in (1) is given by*

$$E(S_{N_n}) = n \times E(Y) \times E(X) = n\alpha\mu.$$

- (3) *Variance of random summation in (1) is defined by*

$$D(S_{N_n}) = n [E(Y) \times D(X) + D(Y) \times (E(X))^2] = n [\alpha\sigma^2 + \tau^2\mu^2].$$

Proof. (1) It is easy to verify that

$$\begin{aligned} F_{S_{N_n}}(x) &= P(S_{N_n} \leq x) = E(\mathbb{I}_{(-\infty, x]}(S_{N_n})) = E(E(\mathbb{I}_{(-\infty, x]}(S_{N_n}) | N_n)) \\ &= \sum_{k=1}^{\infty} P(N_n = k) E(\mathbb{I}_{(-\infty, x]}(S_k) | N_n = k) \\ &= \sum_{k=1}^{\infty} P(N_n = k) E(\mathbb{I}_{(-\infty, x]}(S_k)) \\ &= \sum_{k=1}^{\infty} P(N_n = k) P(S_k \leq x) = \sum_{k=1}^{\infty} P(N_n = k) \times F_X^{*k}(x), \end{aligned}$$

where

$$\mathbb{I}_A(x) = \begin{cases} 1, & \text{if } x \in A \\ 0, & \text{if } x \notin A. \end{cases}$$

(2) Evidently,

$$\begin{aligned} E(S_{N_n}) &= \sum_{k=1}^{\infty} P(N_n = k) \times E(S_k) = \sum_{k=1}^{\infty} kP(N_n = k) \times E(X) \\ &= nE(Y)E(X) = n\alpha\mu. \end{aligned}$$

(3) It is obvious that

$$\begin{aligned} E(S_{N_n}^2) &= E(E(S_{N_n}^2 | N_n)) = \sum_{k=1}^{\infty} P(N_n = k) \times E(S_k^2 | N_n = k) \\ &= \sum_{k=1}^{\infty} P(N_n = k) \times E(S_k^2) \\ &= \sum_{k=1}^{\infty} P(N_n = k) \left(\sum_{j=1}^k EX_j^2 + \sum_{\substack{i \neq j \\ i, j=1}}^k E(X_i) \times E(X_j) \right) \\ &= \sum_{k=1}^{\infty} P(N_n = k) \left[kE(X^2) + (k^2 - k)(E(X))^2 \right] \\ &= \sum_{k=1}^{\infty} P(N_n = k) \left[kD(X) + k^2(E(X))^2 \right] \\ &= E(N_n) \times D(X) + E(N_n^2) \times (E(X))^2. \end{aligned}$$

Therefore,

$$\begin{aligned} D(S_{N_n}) &= E(S_{N_n}^2) - (E(S_{N_n}))^2 \\ &= E(N_n) \times D(X) + E(N_n^2)(E(X))^2 - (E(N_n) \times E(X))^2 \\ &= n [E(Y) \times D(X) + D(Y) \times (E(X))^2] = n [\alpha\sigma^2 + \tau^2\mu^2]. \quad \square \end{aligned}$$

Remark 2.1. For the case of $n = 1$, we have the well-known Wald's identity (see for instance [8], Theorem 9.1, page 194)

$$E(S_{Y_1}) = E(Y_1) \times E(X) = E(Y) \times E(X) = \alpha\mu,$$

and

$$D(S_{Y_1}) = E(Y_1) \times D(X) + (E(X))^2 \times D(Y_1) = \alpha\sigma^2 + \mu^2\tau^2.$$

Remark 2.2. On account of the Proposition 2.1, we consider following example, which will be used in next section. Suppose that Y, Y_1, Y_2, \dots, Y_n are independent, geometric distributed random variables with the success probability $p \in (0, 1)$ (for short, $Y_j \sim \mathcal{Geo}(p), p \in (0, 1)$). We remark that the partial

sum $N_n = \sum_{j=1}^n Y_j$ will be a negative-binomial distributed random variable with two parameters n and p , $N_n \sim \mathcal{NB}(n, p)$, $n = 1, 2, \dots$, with

$$P(N_n = k) = \binom{k-1}{n-1} p^n (1-p)^{k-n}, \quad k = n, n+1, n+2, \dots$$

Additionally, assume that random variables X, X_1, X_2, \dots and Y, Y_1, Y_2, \dots, Y_n are independent. Then, the random summation S_{N_n} will become a negative-binomial random summation of random variables X_1, X_2, \dots with following probability distribution function

$$\begin{aligned} F_{S_{N_n}}(x) &= P(S_{N_n} \leq x) = \sum_{k=1}^{\infty} \binom{k-1}{n-1} p^n (1-p)^{k-n} \times P(S_k \leq x) \\ &= \sum_{k=1}^{\infty} \binom{k-1}{n-1} p^n (1-p)^{k-n} \times F_X^{*k}(x), \quad k = n, n+1, n+2, \dots, \end{aligned}$$

where $F_X^{*k}(x)$ denotes the k -th convolution power of the distribution function $F_X(x)$.

Proposition 2.2. *Assume that the hypotheses of Proposition 2.1 hold. Then*

- (1) *Generating function of the random summation S_{N_n} in (1) is given by*

$$\psi_{S_{N_n}}(t) = E(t^{S_{N_n}}) = (\psi_Y \circ \psi_X(t))^n.$$

- (2) *Characteristic function of the random summation S_{N_n} in (1) is defined by*

$$\varphi_{S_{N_n}}(t) = E(e^{itS_{N_n}}) = (\psi_Y \circ \varphi_X(t))^n.$$

Proof. Write $p_k = P(N_n = k)$, $k = 0, 1, \dots$. Then,

- (1) Direct computation shows that the generating function of the random summation S_{N_n} will be given by

$$\begin{aligned} \psi_{S_{N_n}}(t) &= E(t^{S_{N_n}}) = E(E(t^{S_{N_n}} | N_n)) \\ &= \sum_{k=0}^{\infty} p_k E(t^{S_{N_n}} | N_n = k) = \sum_{k=0}^{\infty} p_k E(t^{S_k}) \\ &= \sum_{k=0}^{\infty} p_k [E(t^X)]^k = \sum_{k=0}^{\infty} p_k [\psi_X(t)]^k = E([\psi_X(t)]^{N_n}) \\ &= E(\psi_X(t)^{Y_1} \times \psi_X(t)^{Y_2} \times \dots \times \psi_X(t)^{Y_n}) = \left(E(\psi_X(t)^Y)\right)^n \\ &= (\psi_Y \circ \psi_X(t))^n. \end{aligned}$$

- (2) Upon simple computation, the characteristic function of the random summation S_{N_n} is defined as follows

$$\varphi_{S_{N_n}}(t) = E(e^{itS_{N_n}}) = E(E(e^{itS_{N_n}} | N_n)) = \sum_{k=0}^{\infty} p_k E(e^{itS_{N_n}} | N_n = k)$$

$$\begin{aligned}
&= \sum_{k=0}^{\infty} p_k E(e^{itS_k}) = \sum_{k=0}^{\infty} p_k [E(e^{itX})]^k = \sum_{k=0}^{\infty} p_k [\varphi_X(t)]^k \\
&= E([\varphi_X(t)]^{N_n}) = E(\varphi_X(t)^{Y_1} \times \varphi_X(t)^{Y_2} \times \cdots \times \varphi_X(t)^{Y_n}) \\
&= \left(E(\varphi_X(t)^Y)\right)^n = (\psi_Y \circ \varphi_X(t))^n. \quad \square
\end{aligned}$$

Remark 2.3. For the case of $n = 1$, when $P(N_1 = Y_1) = 1$, we will return to known results (see for instance [8], Theorem 9.2 and Theorem 9.3, page 194)

- (1) $\psi_{S_{Y_1}}(t) = \psi_Y \circ \psi_X(t)$,
- (2) $\varphi_{S_{Y_1}}(t) = \psi_Y \circ \varphi_X(t)$.

3. Asymptotic behaviors of the extended random summations

Throughout the forthcoming, unless otherwise specified, we shall denote by \xrightarrow{d} and \xrightarrow{P} the convergence in distribution and in probability, respectively. The following limit theorems are extensions of classical results for random summations of i.i.d. random variables.

Theorem 3.1 (Weak law of large numbers). *Let X, X_1, X_2, \dots be a sequence of i.i.d. random variables with finite mean $\mu = E(X)$. Suppose that Y, Y_1, Y_2, \dots, Y_n are i.i.d., non-negative integer-valued random variables with finite mean $\alpha = E(Y)$. Moreover, assume that random variables X, X_1, X_2, \dots and Y, Y_1, Y_2, \dots, Y_n are independent. Write $S_{N_n} := \sum_{i=1}^{N_n} X_i$, where $N_n = Y_1 + Y_2 + \cdots + Y_n$. Then, the weak law of large numbers for random summation states in form*

$$(2) \quad \frac{S_{N_n}}{n} \xrightarrow{P} \alpha\mu \quad \text{as } n \rightarrow \infty.$$

Proof. In view of the continuity theorem for characteristic function (see [8] for more details), it suffices to prove that

$$\varphi_{\frac{S_{N_n}}{n}}(t) \rightarrow \varphi_{\alpha\mu}(t) \quad \text{as } n \rightarrow \infty, \quad \text{for } -\infty < t < +\infty.$$

According to Proposition 2.2, the characteristic function of random summation $\frac{S_{N_n}}{n}$ is given by

$$\varphi_{\frac{S_{N_n}}{n}}(t) = \varphi_{S_{N_n}}\left(\frac{t}{n}\right) = \left[\psi_Y \circ \varphi_X\left(\frac{t}{n}\right)\right]^n.$$

Setting $h(t) = \ln[\psi_Y \circ \varphi_X(t)]$. Direct computation shows that $h(0) = 0$, and the derivative of the function $h(t)$ is calculated by

$$h'(t) = \frac{dh(t)}{dt} = \frac{\frac{d}{dt}[\psi_Y \circ \varphi_X(t)]}{\psi_Y \circ \varphi_X(t)} = \frac{E\left[Y(\varphi_X(t))^{Y-1}\varphi_X'(t)\right]}{E[(\varphi_X(t))^Y]}.$$

It is easily seen that $h'(0) = E[Yi\mu] = iE(Y)\mu = i\alpha\mu$. Then, by letting $n \rightarrow \infty$, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \varphi_{\frac{S_{N_n}}{n}}(t) &= \lim_{n \rightarrow \infty} \exp[nh(t/n)] = \lim_{t/n \rightarrow 0} \exp\left[\frac{h(t/n) - h(0)}{t/n - 0} \times t\right] \\ &= \exp\left(h'(0)\right) = \exp(\alpha\mu t) = \varphi_{\alpha\mu}(t) \quad \text{for all } t. \end{aligned}$$

Consequently,

$$\frac{S_{N_n}}{n} \xrightarrow{P} \alpha\mu \quad \text{as } n \rightarrow \infty.$$

This finishes the proof. \square

Theorem 3.2 (Central limit theorem). *Let X, X_1, X_2, \dots be a sequence of i.i.d. random variables with mean $\mu = E(X)$ and positive, finite variance $D(X) = \sigma^2 < +\infty$. Suppose that Y, Y_1, Y_2, \dots, Y_n is a sequence of i.i.d., positive integer-value random variables with mean $\alpha = E(Y) \in (0, +\infty)$ and positive, finite variance $D(Y) = \tau^2 < +\infty$. Additionally, assume that random variables X, X_1, X_2, \dots and Y, Y_1, Y_2, \dots, Y_n are independent. Write $S_{N_n} := \sum_{j=1}^{N_n} X_j$, where $N_n := Y_1 + Y_2 + \dots + Y_n$. Then,*

$$\frac{S_{N_n} - n\alpha\mu}{\sqrt{n[\alpha\sigma^2 + \mu^2\tau^2]}} \xrightarrow{d} \mathcal{N}(0, 1) \quad \text{as } n \rightarrow \infty.$$

Proof. According to the assumptions on sequence Y, Y_1, Y_2, \dots , it is easily seen that the central limit theorem for a sequence of Y_1, Y_2, \dots holds, i.e.,

$$(3) \quad \frac{N_n - E(N_n)}{\sqrt{D(N_n)}} = \frac{N_n - n\alpha}{\tau\sqrt{n}} \xrightarrow{d} \mathcal{N}(0, 1) \quad \text{as } n \rightarrow \infty.$$

By an argument analogous, on account of above assumptions for sequence X, X_1, X_2, \dots , the central limit theorem confirms that

$$(4) \quad \frac{S_n - E(S_n)}{\sqrt{D(S_n)}} = \frac{S_n - n\mu}{\sigma\sqrt{n}} \xrightarrow{d} \mathcal{N}(0, 1) \quad \text{as } n \rightarrow \infty.$$

According to Corollary 4 from [21], combining the (3) with (4), we conclude that

$$\frac{S_{N_n} - E(S_{N_n})}{\sqrt{D(S_{N_n})}} = \frac{S_{N_n} - n\alpha\mu}{\sqrt{n[\alpha\sigma^2 + \mu^2\tau^2]}} \xrightarrow{d} \mathcal{N}(0, 1) \quad \text{as } n \rightarrow \infty.$$

(See also [5], Chapter VIII, Theorem 4, page 265 or [8], Theorem 3.2, page 346 or [20], Chapter VIII, Section 7, Theorem 2, page 473). The proof is completed. \square

Let X, X_1, X_2, \dots be a sequence of i.i.d. random variables with mean $E(X) = \mu$, and finite variance $D(X) = \sigma^2 < +\infty$. Suppose that Y, Y_1, Y_2, \dots, Y_n are independent, Bernoulli distributed random variables, $Y_j \sim \text{Bernoulli}(p)$, $p \in (0, 1)$. Moreover, assume that they are independent of all X_j , $j = 1, 2, \dots$. For $n \geq 1$, setting a partial sum $N_n = \sum_{j=1}^n Y_j$. It is easily seen that the N_n is a binomial random variable, $N_n \sim B_n(p)$, $n \geq 1$, $p \in (0, 1)$. Thus, the random

sum $S_{N_n} := X_1 + X_2 + \cdots + X_{N_n}$ has become a binomial random summation. The following corollary is a direct result of Theorem 3.2.

Corollary 3.1. *Assume that the hypotheses of Theorem 3.2 hold. Then*

$$\frac{S_{N_n} - np\mu}{\sqrt{n[p\sigma^2 + \mu^2p(1-p)]}} \xrightarrow{d} \mathcal{N}(0, 1) \quad \text{as } n \rightarrow \infty.$$

Proof. Under the above assumption that $Y \sim \text{Ber}(p)$, $p \in (0, 1)$, it follows that

$$E(Y) = p \quad \text{and} \quad D(Y) = p(1-p) \leq \frac{1}{4}$$

and

$$E(S_{N_n}) = n \times E(Y) \times E(X) = np\mu,$$

with

$$D(S_{N_n}) = n[E(Y) \times D(X) + D(Y) \times (E(X))^2] = np\sigma^2 + \mu^2np(1-p).$$

Then, on account of the above Theorem 3.2, we finish the proof. \square

Assume that Y, Y_1, Y_2, \dots, Y_n are independent, geometric distributed random variables with common success probabilities p , (in short, $Y_j \sim \text{Geo}(p)$, $p \in (0, 1)$). Setting $N_n = \sum_{j=1}^n Y_j$, $n \geq 1$ and $S_{N_n} = \sum_{j=1}^{N_n} X_j$. It is to be noticed that N_n is a negative-binomial random variable with two parameters n and p , in short, $N_n \sim \mathcal{NB}(n, p)$, $n \geq 1$, $p \in (0, 1)$. Thus, the random sum S_{N_n} has become a negative-binomial random summation. The forthcoming theorems will be related to these negative-binomial random summations (see Remark 2.2).

Theorem 3.3. *Let X, X_1, X_2, \dots be a sequence of i.i.d., positive random variables with a finite expectation $E(X) = \mu \in (0, +\infty)$. Suppose that Y, Y_1, Y_2, \dots, Y_n are independent, geometric random variables with common success probabilities p , $Y_j \sim \text{Geo}(p)$, $p \in (0, 1)$. Moreover, assume that random variables X, X_1, X_2, \dots and Y, Y_1, Y_2, \dots, Y_n are independent. Setting $N_n = \sum_{j=1}^n Y_j$, $n \geq 1$ and $S_{N_n} = \sum_{j=1}^{N_n} X_j$. Then,*

$$\frac{S_{N_n}}{E(N_n)} \xrightarrow{d} \mathcal{G} \quad \text{as } p \rightarrow 0^+,$$

where \mathcal{G} is a Gamma random variable with two parameters n and $\frac{n}{\mu}$ (in short, $\mathcal{G} \sim \text{Gamma}(n, \frac{n}{\mu})$).

Proof. Let us denote by $\varphi(t)$ and $\psi(t)$ the characteristic function and generating functions, respectively. It can be verified that the characteristic function of a Gamma distributed random variable is $\varphi_{\mathcal{G}}(t) = \left(\frac{1}{1-i\mu\frac{t}{n}}\right)^n$, and the generating function of $Y \sim \text{Geo}(p)$ is given by

$$\psi_Y(t) = \frac{pt}{1 - (1-p)t}.$$

On account of Proposition 2.2, the characteristic function of $\frac{S_{N_n}}{E(N_n)}$ will be given as follows

$$(5) \quad \varphi_{\frac{S_{N_n}}{E(N_n)}}(t) = \varphi_{S_{N_n}}\left(\frac{p}{n}t\right) = \left[\psi_Y \circ \varphi_X\left(\frac{p}{n}t\right)\right]^n = \left[\frac{p\varphi_X\left(\frac{p}{n}t\right)}{1 - (1-p)\varphi_X\left(\frac{p}{n}t\right)}\right]^n.$$

According to Taylor's expansion, there exists a real number c between 0 and $\frac{p}{n}t$, such that

$$\varphi_X\left(\frac{p}{n}t\right) = \varphi_X(0) + \frac{p}{n}t\varphi'_X(c) = 1 + \frac{p}{n}t\varphi'_X(c).$$

Therefore

$$\begin{aligned} \varphi_{\frac{S_{N_n}}{E(N_n)}}(t) &= \left[\frac{p\varphi_X\left(\frac{p}{n}t\right)}{1 - (1-p)\varphi_X\left(\frac{p}{n}t\right)}\right]^n = \left[\frac{p\left(1 + \frac{p}{n}t\varphi'_X(c)\right)}{1 - (1-p)\left(1 + \frac{p}{n}t\varphi'_X(c)\right)}\right]^n \\ &= \left[\frac{1 + \frac{p}{n}t\varphi'_X(c)}{1 - \frac{t}{n}\varphi'_X(c) + \frac{p}{n}t\varphi'_X(c)}\right]^n. \end{aligned}$$

Letting $p \rightarrow 0^+$, so $\frac{p}{n}t \rightarrow 0^+$, it follows that $c \rightarrow 0$. Then,

$$(6) \quad \begin{aligned} \lim_{p \rightarrow 0^+} \varphi_{\frac{S_{N_n}}{E(N_n)}}(t) &= \left[\frac{1}{1 - \frac{t}{n}\varphi'_X(0)}\right]^n \\ &= \left[\frac{1}{1 - \frac{t}{n}iE(X)}\right]^n = \left(\frac{1}{1 - i\mu\frac{t}{n}}\right)^n = \varphi_{\mathcal{G}}(t). \end{aligned}$$

This finished the proof. \square

It is worth pointing out that when $P(N_1 = Y_1) = 1$, we have a desired geometric sum S_{N_1} and the following corollary will be hold and it is analogous to Renyi's results in 1957 for geometric sum of i.i.d. positive-valued random variables (see [13, 17, 20], for more details).

Corollary 3.2. *Let X, X_1, X_2, \dots be a sequence of i.i.d., positive-valued random variables with finite expectation $E(X) = \mu \in (0, +\infty)$. Let Y_1 be a geometric random variable with success probability $p \in (0, 1)$. Moreover, assume that random variables X, X_1, X_2, \dots and Y_1 are independent. Write $S_{Y_1} = \sum_{j=1}^{Y_1} X_j$. Then,*

$$\frac{S_{Y_1}}{E(Y_1)} \xrightarrow{d} \mathcal{W}^{(\mu)} \quad \text{as } p \rightarrow 0^+.$$

where $\mathcal{W}^{(\mu)}$ is an exponential random variable with expectation $E(\mathcal{W}^{(\mu)}) = \mu$, (in short, $\mathcal{W}^{(\mu)} \sim \text{Exp}(\mu^{-1})$).

Proof. Applying (5) and (6) when $n = 1$, with $N_1 = Y_1$ and $S_{N_1} = \sum_{i=1}^{Y_1} X_i$, we conclude that

$$\lim_{p \rightarrow 0^+} \varphi_{\frac{S_{Y_1}}{E(Y_1)}} = \left(\frac{1}{1 - i\mu t} \right) = \varphi_{\mathcal{W}^\mu}(t).$$

It follows that

$$\frac{S_{Y_1}}{E(Y_1)} \xrightarrow{d} \mathcal{W}^{(\mu)} \quad \text{as } p \rightarrow 0^+.$$

The proof is complete finished \square

Suppose that Y, Y_1, Y_2, \dots, Y_n are independent, geometric distributed random variables with common success probability p , $Y_j \sim \mathcal{Geo}(p)$, $p \in (0, 1)$. For every $n \geq 1$, we will denote by $N_n = \sum_{j=1}^n Y_j$ the partial sum. The random sum N_n is a negative-binomial random variable with parameters n and p , $N_n \sim \mathcal{NB}(n, p)$, $n \geq 1$, $p \in (0, 1)$. Thus, the random summation $S_{N_n}^2 = \sum_{j=1}^{N_n} X_j^2$ is a negative-binomial random summation of squares of standard normal distributed random variables.

Theorem 3.4. *Let X, X_1, X_2, \dots be a sequence of independent, standard normal distributed random variables, $X_j \sim \mathcal{N}(0, 1)$, $j = 1, 2, \dots$. Suppose that Y, Y_1, Y_2, \dots, Y_n are independent, geometric distributed random variables with common success probability p , $Y_j \sim \mathcal{Geo}(p)$, $p \in (0, 1)$. Moreover, assume that random variables X, X_1, X_2, \dots and Y, Y_1, Y_2, \dots, Y_n are independent. For every $n \geq 1$, write $N_n = \sum_{j=1}^n Y_j$ and define $S_{N_n}^2 = \sum_{j=1}^{N_n} X_j^2$. Then,*

$$\frac{S_{N_n}^2}{E(N_n)} \xrightarrow{d} \mathcal{G} \quad \text{as } p \rightarrow 0^+,$$

where \mathcal{G} is a gamma distributed random variable, $\mathcal{G} \sim \text{Gamma}(n, n)$.

Proof. Let us denote by $\varphi_{X^2}(t)$ the characteristic function of random variable X^2 and by $\varphi_{\mathcal{G}}(t)$ the characteristic function of random variables \mathcal{G} . It is easily seen that the characteristic function of a gamma random variable $\varphi_{\mathcal{G}}(t)$ is given by

$$\varphi_{\mathcal{G}}(t) = \left(\frac{n}{n - it} \right)^n.$$

Since $X_j \sim \mathcal{N}(0, 1)$, we inference that X_j^2 will be a chi-squared random variable with degree of freedom 1. (in short, $X_j^2 \sim \chi_1^2(1)$.) Therefore, the characteristic function of X_j^2 is defined as follows

$$\varphi_{X^2}(t) = \frac{1}{\sqrt{1 - 2it}}.$$

Since $Y \sim \mathcal{Geo}(p)$, the generating function of Y is given by

$$\psi_Y(t) = \frac{pt}{1 - (1-p)t}.$$

Then, according to Proposition 2.2, the characteristic function of $S_{N_n}^2$ is defined by

$$\varphi_{S_{N_n}^2}(t) = [\psi_Y \circ \varphi_{X^2}(t)]^n = \left[\frac{p\varphi_{X^2}(t)}{1 - (1-p)\varphi_{X^2}(t)} \right]^n.$$

Thus, the characteristic function of $\frac{S_{N_n}^2}{E(N_n)}$ is given as follows

$$\begin{aligned} \varphi_{\frac{S_{N_n}^2}{E(N_n)}}(t) &= \varphi_{S_{N_n}^2}\left(\frac{p}{n}t\right) = \left[\frac{p\varphi_{X^2}\left(\frac{p}{n}t\right)}{1 - (1-p)\varphi_{X^2}\left(\frac{p}{n}t\right)} \right]^n \\ &= \left(\frac{\sqrt{1 - 2i\frac{p}{n}t + 1 - p}}{2 - 2i\frac{t}{n} - p} \right)^n. \end{aligned}$$

Letting $p \rightarrow 0^+$, we conclude that

$$(7) \quad \lim_{p \rightarrow 0^+} \varphi_{\frac{S_{N_n}^2}{E(N_n)}}(t) = \left(\frac{n}{n - it} \right)^n = \varphi_{\mathcal{G}}(t).$$

We have the complete proof. \square

It is worth noticing that in case of $P(N_1 = Y_1) = 1$, we have a result related to geometric sum of square standard normal distributed random variables, $S_{Y_1} = \sum_{j=1}^{Y_1} X_j^2$. This geometric sum should be considered as a $\chi_{Y_1}^2$ -squared random variable with geometric degrees of freedom Y_1 .

Corollary 3.3. *Let X, X_1, X_2, \dots be a sequence of independent, standard normal distributed random variables, $X \sim \mathcal{N}(0, 1)$. Let Y_1 be a geometric random variable with parameter p , $Y_1 \sim \text{Geo}(p)$, $p \in (0, 1)$. Moreover, assume that random variables X, X_1, X_2, \dots and Y_1 are independent. Set $S_{Y_1}^2 = \sum_{j=1}^{Y_1} X_j^2$. Then,*

$$\frac{S_{Y_1}^2}{E(Y_1)} \xrightarrow{d} \mathcal{W}^{(1)} \quad \text{as } p \rightarrow 0^+.$$

where $\mathcal{W}^{(1)} \sim \text{Exp}(1)$ is an exponential random variable with mean 1.

Proof. The proof is immediate from (7) with $n = 1$, i.e.,

$$\lim_{p \rightarrow 0^+} \varphi_{\frac{S_{Y_1}^2}{E(Y_1)}}(t) = \left(\frac{1}{1 - it} \right) = \varphi_{\mathcal{W}^{(1)}}(t). \quad \square$$

The following theorem is concerning with the negative-binomial random summation $S_{N_n} = \sum_{j=1}^{N_n} X_j$, where $N_n = Y_1 + Y_2 + \dots + Y_n$, with $Y_j \sim \text{Geo}(p)$, $p \in (0, 1)$.

Theorem 3.5. Let X, X_1, X_2, \dots be a sequence of i.i.d., non-negative random variables with mean zero $E(X) = 0$ and finite variance $D(X) = \sigma^2 < +\infty$. Suppose that Y, Y_1, Y_2, \dots, Y_n are independent, geometric distributed random variables with common success probability p , $Y_j \sim \mathcal{Geo}(p)$, $p \in (0, 1)$. Moreover, assume that random variables X, X_1, X_2, \dots and Y, Y_1, Y_2, \dots, Y_n are independent. Write $N_n = \sum_{j=1}^n Y_j$ and define $S_{N_n} = \sum_{j=1}^{N_n} X_j$. Then,

$$(8) \quad \frac{S_{N_n}}{\sqrt{E(N_n)}} \xrightarrow{d} \sum_{j=1}^n \mathcal{L}_j \quad \text{as } p \rightarrow 0^+,$$

where $\mathcal{L}_1, \mathcal{L}_2, \dots, \mathcal{L}_n$ are n independent, Laplace distributed random variables with parameters zero and $\frac{\sqrt{2n}}{\sigma}$, $\mathcal{L}_j \sim \text{Laplace}\left(0, \frac{\sqrt{2n}}{\sigma}\right)$, $j = 1, 2, \dots, n$.

Proof. An easily computation that, for a geometric random variable $Y \sim \mathcal{Geo}(p)$, the generating function is given by

$$\psi_Y(t) = \frac{pt}{1 - (1-p)t}.$$

According to Proposition 2.2, the characteristic function of $\frac{S_{N_n}}{\sqrt{E(N_n)}}$ will be calculated as follows

$$\begin{aligned} \varphi_{\frac{S_{N_n}}{\sqrt{E(N_n)}}}(t) &= \varphi_{S_{N_n}}\left(\sqrt{\frac{p}{n}}t\right) = \left[\psi_Y \circ \varphi_X\left(\sqrt{\frac{p}{n}}t\right)\right]^n \\ &= \left[\frac{p\varphi_X\left(\sqrt{\frac{p}{n}}t\right)}{1 - (1-p)\varphi_X\left(\sqrt{\frac{p}{n}}t\right)}\right]^n. \end{aligned}$$

On account of Taylor expansion, there exists a real number c between 0 and $\sqrt{\frac{p}{n}}t$, we obtain

$$\begin{aligned} \varphi_X\left(\sqrt{\frac{p}{n}}t\right) &= \varphi_X(0) + \sqrt{\frac{p}{n}}t\varphi_X'(0) + \frac{pt^2}{2n}\varphi_X''(c) \\ &= 1 + \sqrt{\frac{p}{n}}t\varphi_X'(0) + \frac{pt^2}{2n}\varphi_X''(c) \\ &= 1 + \frac{pt^2}{2n}\varphi_X''(c). \end{aligned}$$

Therefore

$$\begin{aligned} \varphi_{\frac{S_{N_n}}{\sqrt{E(N_n)}}}(t) &= \left[\frac{p\varphi_X\left(\sqrt{\frac{p}{n}}t\right)}{1 - (1-p)\varphi_X\left(\sqrt{\frac{p}{n}}t\right)}\right]^n = \left[\frac{p\left[1 + \frac{pt^2}{2n}\varphi_X''(c)\right]}{1 - (1-p)\left[1 + \frac{pt^2}{2n}\varphi_X''(c)\right]}\right]^n \\ &= \left[\frac{1 + \frac{pt^2}{2n}\varphi_X''(c)}{1 - \frac{t^2}{2n}\varphi_X''(c) + \frac{pt^2}{2n}\varphi_X''(c)}\right]^n. \end{aligned}$$

Letting $p \rightarrow 0^+$, it follows $\sqrt{\frac{E}{n}}t \rightarrow 0^+$. Thus $c \rightarrow 0$. Then,

$$\lim_{p \rightarrow 0^+} \varphi \frac{S_{N_n}}{\sqrt{E(N_n)}}(t) = \left[\frac{1}{1 - \frac{t^2}{2n} \varphi''(0)} \right]^n = \left[\frac{1}{1 + \frac{\sigma^2 t^2}{2n}} \right]^n = \varphi_{\sum_{j=1}^n \mathcal{L}_j}(t).$$

This finishes the proof. \square

Corollary 3.4. *Let X, X_1, X_2, \dots be a sequence of i.i.d., non-negative random variables with mean $E(X) = 0$ and finite variance $D(X) = \sigma^2 < +\infty$. Assume that $Y_1 \sim \text{Geo}(p)$ and independent of all $X_j, j \geq 1$. Then,*

$$\frac{S_{Y_1}}{\sqrt{E(Y_1)}} \xrightarrow{d} \mathcal{L} \quad \text{as } n \rightarrow \infty,$$

where $S_{Y_1} = \sum_{j=1}^{Y_1} X_j$ and $\mathcal{L} \sim \text{Laplace}(0, \frac{\sqrt{2}}{\sigma})$.

Proof. On account of (8) for $n=1$, the corollary is proven. \square

Acknowledgments. The research has been supported by the Vietnam's National Foundation For Science and Technology Development (NAFOSTED, Vietnam) under Grant N. 101.01-2010.02. The authors would like to thank the referees for several suggested comments to improve the manuscript.

References

- [1] J. Blanchet and P. Glynn, *Uniform renewal theory with applications to expansions of random geometric sums*, Adv. in Appl. Probab. **39** (2007), no. 4, 1070–1097.
- [2] L. H. Y. Chen, L. Goldstein, and Q.-M. Shao, *Normal approximation by Stein's method, Probability and its Applications* (New York), Springer, Heidelberg, 2011.
- [3] L. H. Y. Chen and Q.-M. Shao, *Normal approximation for nonlinear statistics using a concentration inequality approach*, Bernoulli **13** (2007), no. 2, 581–599.
- [4] R. T. Durrett and S. I. Resnick, *Weak convergence with random indices*, Stochastic Processes Appl. **5** (1977), no. 3, 213–220.
- [5] W. Feller, *An Introduction to Probability Theory and its Applications. Vol. I*, John Wiley and Sons, Inc., New York, 1957.
- [6] B. V. Gnedenko, *Limit theorems for sums of a random number of positive independent random variables*, in Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability (Univ. California, Berkeley, Calif., 1970/1971), Vol. II: Probability theory, 537–549, Univ. California Press, Berkeley, CA, 1972.
- [7] B. V. Gnedenko and V. Yu. Korolev, *Random Summation*, CRC Press, Boca Raton, FL, 1996.
- [8] A. Gut, *Probability: a graduate course*, Springer Texts in Statistics, Springer, New York, 2005.
- [9] C.-Y. Hu and T.-L. Cheng, *A characterization of distributions by random summation*, Taiwanese J. Math. **16** (2012), no. 4, 1245–1264.
- [10] T. L. Hung and T. T. Thanh, *Some results on asymptotic behaviors of random sums of independent identically distributed random variables*, Commun. Korean Math. Soc. **25** (2010), no. 1, 119–128.
- [11] T. L. Hung, T. T. Thanh, and B. Q. Vu, *Some results related to distribution functions of chi-square type random variables with random degrees of freedom*, Bull. Korean Math. Soc. **45** (2008), no. 3, 509–522.

- [12] U. Islak, *Asymptotic normality of random sums of m -dependent random variables*, arXiv:1303.2386 [Math.PR], 2013.
- [13] V. Kalashnikov, *Geometric sums: bounds for rare events with applications*, Mathematics and its Applications, **413**, Kluwer Academic Publishers Group, Dordrecht, 1997.
- [14] H. Klaver and N. Schmitz, *An inequality for the asymmetry of distributions and a Berry-Esseen theorem for random summation*, JIPAM. J. Inequal. Pure Appl. Math. **7** (2006), no. 1, Article 2, 12 pp.
- [15] V. Yu. Korolev and V. M. Kruglov, *Limit theorems for random sums of independent random variables*, in Stability problems for stochastic models (Suzdal, 1991), 100–120, Lecture Notes in Math., 1546, Springer, Berlin, 1991.
- [16] S. Kotz, T. J. Kozubowski, and K. Podgorski, *The Laplace Distribution and Generalizations*, Birkhäuser Boston, Inc., Boston, MA, 2001.
- [17] V. M. Kruglov and V. Yu. Korolev, *Limit theorems for random sums*, Moscow University Press, Moscow, (in Russian), 1990.
- [18] J. A. Melamed, *Limit theorems in the set-up of summation of a random number of independent identically distributed random variables*, in Stability problems for stochastic models (Sukhumi, 1987), 194–228, Lecture Notes in Math., 1412, Springer, Berlin, 1989.
- [19] E. Omev and R. Vesilo, *Random sums of random variables and vectors: including infinite means and unequal length sums*, Lith. Math. J. **55** (2015), no. 3, 433–450.
- [20] A. Renyi, *Probability theory*, translated by László Vekerdí, North-Holland Publishing Co., Amsterdam, 1970.
- [21] H. Robbins, *The asymptotic distribution of the sum of a random number of random variables*, Bull. Amer. Math. Soc. **54** (1948), 1151–1161.
- [22] J. K. Sunklodas, *On the rate of convergence of L_p norms in the CLT for Poisson and gamma random variables*, Lith. Math. J. **49** (2009), no. 2, 216–221.
- [23] ———, *On the normal approximation of a binomial random sum*, Lith. Math. J. **54** (2014), no. 3, 356–365.
- [24] ———, *On the normal approximation of a negative binomial random sum*, Lith. Math. J. **55** (2015), no. 1, 150–158.
- [25] P. Vellaisamy and B. Chaudhuri, *Poisson and compound Poisson approximations for random sums of random variables*, J. Appl. Probab. **33** (1996), no. 1, 127–137.

LE TRUONG GIANG
UNIVERSITY OF FINANCE AND MARKETING
2/4 TRAN XUAN SOAN STREET
DISTRICT N. 7, HO CHI MINH CITY, VIETNAM
Email address: ltgiang@ufm.edu.vn

TRAN LOC HUNG
UNIVERSITY OF FINANCE AND MARKETING
2/4 TRAN XUAN SOAN STREET
DISTRICT N. 7, HO CHI MINH CITY, VIETNAM
Email address: tlhung@ufm.edu.vn