

## THE FIRST EIGENVALUE OF SOME $(p, q)$ -LAPLACIAN AND GEOMETRIC ESTIMATES

SHAHROUD AZAMI

ABSTRACT. We study the nonlinear eigenvalue problem for some of the  $(p, q)$ -Laplacian on compact manifolds with zero boundary condition. In particular, we obtain some geometric estimates for the first eigenvalue.

### 1. Introduction

Let  $M$  be a compact domain in a complete, simply connected, Riemannian manifold  $\mathcal{M}$  of constant sectional curvature. Studying the eigenvalues of geometric operators is an important tool for Riemannian manifolds. There are many mathematicians who investigate properties of the spectrum of Laplacian and estimate the spectrum in terms of the other geometric quantities of  $M$  (see [4], [5], [6], [7], [9], [11], [12]). In [1], Abolarinwa obtained the classical estimates of Faber-Krahn inequality and Cheeger-type inequality for the first eigenvalue of  $p$ -Laplacian operator. In this paper we study the principal eigenvalue of some  $(p, q)$ -Laplacian on Riemannian manifold and prove some of classical results introduced in above for the  $(p, q)$ -Laplacian of any  $p > 1, q > 1$ .

#### 1.1. Eigenvalues of $(p, q)$ -Laplacian

Let  $(M, g)$  be a compact domain in a complete, simply connected, Riemannian manifold  $\mathcal{M}$  of constant sectional curvature and  $f : M \rightarrow \mathbb{R}$  be a smooth function on  $M$  or  $f \in W^{1,p}(M)$  where  $W^{1,p}(M)$  is the Sobolev space. The  $p$ -Laplacian of  $f$  for  $1 < p < \infty$  is defined as

$$(1) \quad \begin{aligned} \Delta_p f &= \operatorname{div}(|\nabla f|^{p-2} \nabla f) \\ &= |\nabla f|^{p-2} \Delta f + (p-2)|\nabla f|^{p-4} (\operatorname{Hess} f)(\nabla f, \nabla f), \end{aligned}$$

where  $f(x) \neq 0$  for  $x \in M$ ,  $f(x) = 0$  on  $\partial M$ ,

$$(\operatorname{Hess} f)(X, Y) = \nabla(\nabla f)(X, Y) = Y.(X.f) - (\nabla_Y X).f, \quad X, Y \in \mathcal{X}(M)$$

---

Received April 4, 2017; Revised August 17, 2017; Accepted August 17, 2017.

2010 *Mathematics Subject Classification.* 35P15, 47J10, 53C21.

*Key words and phrases.*  $(p, q)$ -Laplacian, eigenvalue problem, geometric constant.

and in local coordinate, we have

$$(Hessf)(\partial_i, \partial_j) = \partial_i \partial_j f - \Gamma_{ij}^k \partial_k f.$$

When  $p = 2$ ,  $\Delta = \Delta_2$  is the natural Laplace-Beltrami operator  $\Delta f = \text{div}(\text{grad } f)$ . In this paper, at the first, we consider the nonlinear system introduced in [8], that is

$$(2) \quad \begin{cases} \Delta_p u = -\lambda |u|^\alpha |v|^\beta v & \text{in } M \\ \Delta_q v = -\lambda |u|^\alpha |v|^\beta u & \text{in } M \\ (u, v) \in W^{1,p}(M) \times W^{1,q}(M), \end{cases}$$

where  $p > 1$ ,  $q > 1$  and  $\alpha, \beta$  are real numbers satisfying

$$(3) \quad \alpha > 0, \beta > 0, \quad \frac{\alpha + 1}{p} + \frac{\beta + 1}{q} = 1.$$

We say that  $\lambda$  is an eigenvalue of (2), whenever for some  $u \in W_0^{1,p}(M)$  and  $v \in W_0^{1,q}(M)$ ,

$$(4) \quad \int_M |\nabla u|^{p-2} \langle \nabla u, \nabla \phi \rangle d\mu = \lambda \int_M |u|^\alpha |v|^\beta v \phi d\mu,$$

$$(5) \quad \int_M |\nabla v|^{q-2} \langle \nabla v, \nabla \psi \rangle d\mu = \lambda \int_M |u|^\alpha |v|^\beta u \psi d\mu,$$

where  $\phi \in W^{1,p}(M)$ ,  $\psi \in W^{1,q}(M)$  and  $W_0^{1,p}(M)$  is the closure of  $C_0^\infty(M)$  in Sobolev space  $W^{1,p}(M)$ . The pair  $(u, v)$  is called eigenfunctions. A first positive eigenvalue  $\lambda_{1,p,q}$  of (2) obtained as

$$\inf\{A(u, v) : (u, v) \in W_0^{1,p}(M) \times W_0^{1,q}(M), B(u, v) = 1\},$$

where

$$A(u, v) = \frac{\alpha + 1}{p} \int_M |\nabla u|^p d\mu + \frac{\beta + 1}{q} \int_M |\nabla v|^q d\mu,$$

$$B(u, v) = \int_M |u|^\alpha |v|^\beta uv d\mu.$$

In what follows, we use some identities arising from the co-area formula ([3], [10]). Suppose that  $M(t) = \{x \in M : f(x) > t\}$  and  $N(t) = \{x \in M : f(x) = t\}$ , where  $t$  is a regular value of  $|f|$ . We denote by  $dN(t)$  the Riemannian measure of the set  $N(t)$ . For any  $0 < f \in C_0^\infty(M)$  and  $1 \leq p \leq \infty$ , we have

$$\int_M |\nabla f|^p d\mu = \int_0^\infty \left( \int_{N(t)} |\nabla f|^{p-1} dN(t) \right) dt$$

and

$$\text{Vol}(M(t)) = \int_{M(t)} d\mu = \int_0^\infty \left( \int_{N(t)} \frac{1}{|\nabla f|} dN(t) \right) dt,$$

$$\text{Vol}(\partial M(t)) = \int_{N(t)} |\nabla f| d\mu = \int_0^\infty dN(t) dt.$$

**Theorem 1.1.** *Let  $M$  be a domain in a complete, simply connected Riemannian manifold  $\mathcal{M}$  of constant sectional curvature. Let  $B_\eta(x)$  be a geodesic ball of radius  $\eta$ , centred  $x \in M$  in  $\mathcal{M}$ , such that  $\text{Vol}(M) = \text{Vol}(B_\eta(x))$ . Then the following estimate holds for (2),*

$$\lambda_{1,p,q}(M) \geq \lambda_{1,p,q}(B_\eta(x)).$$

The equality holds if and only if  $M = B_\eta(x)$ .

*Proof.* Let  $v > 0$ ,  $u > 0$  be the pair of eigenfunctions associated with  $\lambda_{1,p,q}(M)$ . Using a symmetrization procedure, we define a radially decreasing function  $u^* : B_\eta \rightarrow \mathbb{R}^+$ . Let  $M_t = \{x \in M : u(x) > t\}$ ,  $N_t = \{x \in M : u(x) = t\}$  and  $N_t^* = \{x \in B_\eta : u^*(x) = t\}$  such that  $\text{Vol}(M_t) = \text{Vol}(B_\eta)$  for each  $t$ . Let  $dN_t$  and  $dN_t^*$  are the  $(n-1)$ -dimensional volume elements of sets  $N_t$  and  $N_t^*$  respectively. By Hölder inequality we have

$$(6) \quad \int_{N_t} |\nabla u|^{p-1} dN_t \geq \frac{\left(\int_{N_t} dN_t\right)^p}{\left(\int_{N_t} |\nabla u|^{-1} dN_t\right)^{p-1}}.$$

Since  $\text{Vol}(M_t) = \text{Vol}(B_\eta)$  and by

$$\frac{d}{dt} \text{Vol}(M_t) = - \int_{N_t} \frac{1}{|\nabla u|} dN_t, \quad \frac{d}{dt} \text{Vol}(B_\eta) = - \int_{N_t^*} \frac{1}{|\nabla u^*|} dN_t^*,$$

we get

$$\left(\int_{N_t} \frac{1}{|\nabla u|} dN_t\right)^k = \left(\int_{N_t^*} \frac{1}{|\nabla u^*|} dN_t^*\right)^k$$

for any real number  $k$ . Now, by the co-area formula for gradient

$$\begin{aligned} \int_M |\nabla u|^p d\mu &= \int_0^\infty \left(\int_{N_t} |\nabla u|^{p-1} dN_t\right) dt \geq \int_0^\infty \frac{\left(\int_{N_t} dN_t\right)^p}{\left(\int_{N_t} |\nabla u|^{-1} dN_t\right)^{p-1}} dt \\ &\geq \int_0^\infty \frac{\left(\int_{N_t^*} dN_t^*\right)^p}{\left(\int_{N_t^*} |\nabla u^*|^{-1} dN_t^*\right)^{p-1}} dt = \int_0^\infty \left(\int_{N_t^*} |\nabla u^*|^{p-1} dN_t^*\right) dt \\ &= \int_{B_\eta} |\nabla u^*|^p d\mu. \end{aligned}$$

Similarly,  $\int_M |\nabla v|^q d\mu \geq \int_{B_\eta} |\nabla v^*|^q d\mu$ , where  $v^*$  defined as  $u^*$ . Therefore

$$\begin{aligned} \lambda_{1,p,q}(M) &= \frac{\alpha+1}{p} \int_M |\nabla u|^p d\mu + \frac{\beta+1}{q} \int_M |\nabla v|^q d\mu \\ &\geq \frac{\alpha+1}{p} \int_{B_\eta} |\nabla u^*|^p d\mu + \frac{\beta+1}{q} \int_{B_\eta} |\nabla v^*|^q d\mu \\ &\geq \lambda_{1,p,q}(B_\eta). \end{aligned}$$

If  $M = B_\eta(x)$  is the geodesic ball of radius  $\eta$  in  $\mathcal{M}$ , then equality is hold and conversely if the equality is hold, then  $\int_M |\nabla u|^p d\mu = \int_{B_\eta} |\nabla u^*|^p d\mu$  which implies that  $M = B_\eta(x)$ .  $\square$

We denote the Cheeger's isoperimetric constant by

$$\mathcal{C}(M) = \inf_{M_c} \frac{\text{Vol}(\partial M_c)}{\text{Vol}(M_c)}, \quad M_c \subset\subset M$$

where the infimum is taken all over manifold  $M_c$  with compact closure in  $M$  and  $\partial M_c$  is assumed to be smooth.

**Theorem 1.2.** *Let  $M$  be a compact manifold with smooth boundary in a complete Riemannian manifold. Then for the first eigenvalue of (2) we have*

$$\lambda_{1,p,q}(M) \geq \frac{\alpha+1}{p} \left(\frac{\mathcal{C}(M)}{p}\right)^p \int_M |u|^p d\mu + \frac{\beta+1}{q} \left(\frac{\mathcal{C}(M)}{q}\right)^q \int_M |v|^q d\mu,$$

where  $(u, v)$  be the pair of positive eigenfunctions corresponding to  $\lambda_{1,p,q}(M)$ .

*Proof.* Let  $(u, v)$  be the pair of eigenfunctions corresponding to  $\lambda_{1,p,q}(M)$  where  $u > 0$ ,  $v > 0$ . By applying co-area formula for  $\varphi \in C_0^\infty(M)$  we have

$$\begin{aligned} \int_M |\nabla \varphi| d\mu &= \int_{-\infty}^{\infty} \left( \int_{N_t} dN_t \right) dt \\ &= \int_{-\infty}^{\infty} \text{Vol}(\partial M_t) dt \\ &= \int_{-\infty}^{\infty} \frac{\text{Vol}(\partial M_t)}{\text{Vol}(M_t)} \text{Vol}(M_t) dt \\ &\geq \inf_t \left( \frac{\text{Vol}(\partial M_t)}{\text{Vol}(M_t)} \right) \int_{-\infty}^{\infty} \text{Vol}(M_t) dt \\ &= \inf_t \left( \frac{\text{Vol}(\partial M_t)}{\text{Vol}(M_t)} \right) \int_M \varphi d\mu = \mathcal{C}(M) \int_M \varphi d\mu. \end{aligned}$$

Suppose  $\varphi = u^p$ , by using the Hölder inequality

$$\begin{aligned} \mathcal{C}(M) \int_M |u|^p d\mu &\leq \int_M |\nabla u^p| d\mu = p \int_M |u^{p-1} \nabla u| d\mu \\ &\leq p \left( \int_M |u|^p d\mu \right)^{\frac{p-1}{p}} \left( \int_M |\nabla u|^p d\mu \right)^{\frac{1}{p}} \end{aligned}$$

by this we get

$$\int_M |\nabla u|^p d\mu \geq \left(\frac{\mathcal{C}(M)}{p}\right)^p \int_M |u|^p d\mu.$$

Hence

$$\lambda_{1,p,q}(M) \geq \frac{\alpha+1}{p} \left(\frac{\mathcal{C}(M)}{p}\right)^p \int_M |u|^p d\mu + \frac{\beta+1}{q} \left(\frac{\mathcal{C}(M)}{q}\right)^q \int_M |v|^q d\mu. \quad \square$$

Another class of  $(p, q)$ -Laplacian is defined as follows

$$(7) \quad \Delta_p u + \Delta_q u = \operatorname{div} ( (|\nabla u|^{p-2} + |\nabla u|^{q-2}) \nabla u ), \quad u \in W = W_0^{1,p}(M) \cap W_0^{1,q}(M),$$

where  $1 < q < p < \infty$  and introduced in [2]. We say that  $\lambda \in \mathbb{R}$  is an eigenvalue of (7) if there exists  $u \in W$ ,  $u \neq 0$  such that

$$(8) \quad -\Delta_p u - \Delta_q u = \lambda |u|^{p-2} u$$

or equivalently

$$(9) \quad \int_M |\nabla u|^{p-2} \nabla u \cdot \nabla v d\mu + \int_M |\nabla u|^{q-2} \nabla u \cdot \nabla v d\mu = \lambda \int_M |u|^{p-2} u v d\mu$$

for any  $v \in W^{1,p}(M) \cap W^{1,q}(M)$ . The first positive eigenvalue  $\lambda_{1,p,q}(M)$  of (7) is obtained as

$$(10) \quad \lambda_{1,p,q}(M) = \inf \left\{ \int_M |\nabla u|^p d\mu + \int_M |\nabla u|^q d\mu : \int_M |u|^p d\mu = 1 \right\}.$$

Similar to the proof of Theorem 1.1 we have:

**Corollary 1.3.** *Let  $M$  be a domain in a complete, simply connected Riemannian manifold  $\mathcal{M}$  of constant sectional curvature and  $B_\eta(x)$  be a geodesic ball of radius  $\eta$  in  $\mathcal{M}$  such that  $\operatorname{Vol}(M) = \operatorname{Vol}(B_\eta(x))$ . Then the following estimate holds for (10),*

$$\lambda_{1,p,q}(M) \geq \lambda_{1,p,q}(B_\eta(x)).$$

The equality holds if and only if  $M = B_\eta(x)$ .

**Theorem 1.4.** *Let  $M$  be a compact manifold with smooth boundary in a complete Riemannian manifold. Then for (10) we have*

$$(11) \quad \lambda_{1,p,q}(M) \geq \left( \frac{\mathcal{C}(M)}{p} \right)^p + \left( \frac{\mathcal{C}(M)}{q} \right)^q,$$

where  $(u, v)$  be the pair of positive eigenfunctions corresponding to  $\lambda_{1,p,q}(M)$ .

*Proof.* Let  $u > 0$  be the eigenfunctions corresponding to  $\lambda_{1,p,q}(M)$ . From the proof of Theorem 1.2 we have

$$\int_M |\nabla u|^p d\mu \geq \left( \frac{\mathcal{C}(M)}{p} \right)^p \int_M |u|^p d\mu = \left( \frac{\mathcal{C}(M)}{p} \right)^p,$$

and

$$\int_M |\nabla u|^q d\mu \geq \left( \frac{\mathcal{C}(M)}{q} \right)^q \int_M |u|^q d\mu \geq \left( \frac{\mathcal{C}(M)}{q} \right)^q.$$

These complete the proof.  $\square$

**Theorem 1.5.** *Let  $\mathcal{M}$  be an  $n$  dimensional complete simply connected Riemannian manifold of negative constant sectional curvature  $-k$ . Let  $M$  be a domain in  $\mathcal{M}$ . Then the follow inequality holds for (10)*

$$\lambda_{1,p,q}(M) \geq \left( \frac{(n-1)\sqrt{-k}}{p} \right)^p + \left( \frac{(n-1)\sqrt{-k}}{q} \right)^q.$$

*Proof.* Let  $B_\eta(x)$  be a geodesic ball of radius  $\eta$  in  $\mathcal{M}$  where  $x \in M$  and  $Vol(M) = Vol(B_\eta(x))$ . For  $y \notin B_\eta(x)$  we define  $\rho(y) = d(x, y)$ .  $\rho(y)$  is differentiable with  $|\nabla\rho| = 1$ , because of  $\mathcal{M}$  is simply connected with negative sectional curvature. By the Laplacian comparison theorem  $\Delta\rho \geq (n-1)\sqrt{-k}$ . If  $\nu$  is the outer unit normal along the boundary of  $B_\eta(x)$ , then  $\langle \nabla\rho, \nu \rangle \leq 1$ . Therefore

$$\begin{aligned} Vol(\partial B_\eta(x)) &= \int_{\partial B_\eta} dA \\ &\geq \int_{\partial B_\eta} \langle \nabla\rho, \nu \rangle dA = \int_{B_\eta} \Delta\rho d\mu \\ &\geq (n-1)\sqrt{-k} \int_{B_\eta} dA = (n-1)\sqrt{-k} Vol(B_\eta), \end{aligned}$$

which implies that

$$(n-1)\sqrt{-k} \leq \mathcal{C}(B_\eta(x)) = \frac{Vol(\partial B_\eta(x))}{Vol(B_\eta)}$$

thus (11) results that

$$\lambda_{1,p,q}(B_\eta(x)) \geq \left(\frac{(n-1)\sqrt{-k}}{p}\right)^p + \left(\frac{(n-1)\sqrt{-k}}{q}\right)^q.$$

Now, by Corollary 1.3, we get

$$\lambda_{1,p,q}(M) \geq \left(\frac{(n-1)\sqrt{-k}}{p}\right)^p + \left(\frac{(n-1)\sqrt{-k}}{q}\right)^q. \quad \square$$

## References

- [1] A. Abolarinwa, *The first eigenvalue of  $p$ -Laplacian and geometric estimates*, Nonl. Analysis and Differential Equations **2** (2014), no. 3, 105–115.
- [2] N. Benouhiba and Z. Belyacine, *A class of eigenvalue problems for the  $(p, q)$ -Laplacian in  $\mathbb{R}^N$* , Int. J. Pure Appl. Math. **80** (2012), no. 5, 727–737.
- [3] I. Chavel, *Eigenvalues in Riemannian geometry*, Pure and Applied Mathematics, 115, Academic Press, Inc., Orlando, FL, 1984.
- [4] S. Y. Cheng, *Eigenfunctions and eigenvalues of Laplacian*, in Differential geometry (Proc. Sympos. Pure Math., Vol. XXVII, Stanford Univ., Stanford, Calif., 1973), Part 2, 185–193, Amer. Math. Soc., Providence, RI.
- [5] Q.-M. Cheng and H. Yang, *Estimates on eigenvalues of Laplacian*, Math. Ann. **331** (2005), no. 2, 445–460.
- [6] E. M. Harrell, II and P. L. Michel, *Commutator bounds for eigenvalues, with applications to spectral geometry*, Comm. Partial Differential Equations **19** (1994), no. 11-12, 2037–2055.
- [7] S. Kawai and N. Nakauchi, *The first eigenvalue of the  $p$ -Laplacian on a compact Riemannian manifold*, Nonlinear Anal. **55** (2003), no. 1-2, 33–46.
- [8] A. El Khalil, S. El Manouni, and M. Ouanan, *Simplicity and stability of the first eigenvalue of a nonlinear elliptic system*, Int. J. Math. Math. Sci. **2005**, no. 10, 1555–1563.
- [9] P. F. Leung, *On the consecutive eigenvalues of the Laplacian of a compact minimal submanifold in a sphere*, J. Austral. Math. Soc. Ser. A **50** (1991), no. 3, 409–416.

- [10] P. Li, *Geometric analysis*, Cambridge Studies in Advanced Mathematics, 134, Cambridge University Press, Cambridge, 2012.
- [11] A.-M. Matei, *First eigenvalue for the  $p$ -Laplace operator*, *Nonlinear Anal.* **39** (2000), no. 8, Ser. A: Theory Methods, 1051–1068.
- [12] H. Takeuchi, *On the first eigenvalue of the  $p$ -Laplacian in a Riemannian manifold*, *Tokyo J. Math.* **21** (1998), no. 1, 135–140.

SHAHROUD AZAMI  
DEPARTMENT OF MATHEMATICS  
FACULTY OF SCIENCES ,  
IMAM KHOMEINI INTERNATIONAL UNIVERSITY  
QAZVIN, IRAN  
*E-mail address:* azami@sci.ikiu.ac.ir