

HARMONIC FINSLER MANIFOLDS WITH MINIMAL HOROSPHERES

CHANG-WAN KIM

ABSTRACT. In this paper we show that complete noncompact harmonic Finsler manifolds with minimal horospheres are flat.

Introduction

The Lichnerowicz conjecture asserts that all harmonic Riemannian manifolds are either flat or locally symmetric spaces of rank one. Szabò [12] proved the Lichnerowicz conjecture for *compact* simply connected Riemannian manifolds. Besson, Courtois, and Gallot [2] confirmed the conjecture for Riemannian manifolds of negative curvature admitting a compact quotient.

On the other hand, Damek and Ricci [5] provided examples showing that in the *noncompact* Riemannian case the conjecture is wrong. The classification of all noncompact harmonic Riemannian manifolds is still a very difficult open problem. Ranjan and Shah [10] showed that noncompact, simply connected harmonic manifolds with minimal horospheres are flat. In this paper, using the Borsuk-Ulam theorem and the existence of stable Jacobi tensors, we extend this result to Finsler manifolds.

Theorem. *Complete noncompact harmonic Finsler manifolds with minimal horospheres are flat.*

For compact Finsler manifolds M without conjugate points it is clear that the volume entropy $h_\mu \geq 0$, and it was conjectured by Burago and Ivanov that either $h_\mu > 0$ or the geodesic flow of M is conjugate to the flow of a flat Finsler metric ([4, 7, 13]). However, to the best of our knowledge, this conjecture is not yet solved. In this paper we provide convincing evidence for this conjecture.

Corollary. *If M is a compact harmonic Finsler manifold without conjugate points such that the volume entropy is zero, then M is flat.*

In particular, harmonic Finsler tori without conjugate points are flat (cf. [3, 9]).

Received April 5, 2017; Revised December 3, 2017; Accepted May 16, 2018.

2000 *Mathematics Subject Classification.* 53C23, 53C60.

Key words and phrases. Finsler geometry, harmonic manifolds, minimal horospheres.

The author is indebted to the referee for useful comments and remarks.

1. Preliminaries

In this section we recall some basics in Finsler geometry and prove some auxiliary facts. We follow the presentation in [8], where most concepts are developed from Riemannian point of view. We refer to [11] as more exhaustive references in Finsler geometry.

A function $F : TM \rightarrow [0, \infty)$ will be called a *Finsler metric* on the manifold M if it is smooth outside the zero section and its restriction to each tangent space $T_x M$ is a quadratically convex norm $F(x, \cdot)$. By a *geodesic* we always mean an affine geodesic, i.e., a constant-speed one. For a nonzero $v \in TM$, we denote by γ_v the unique geodesic with initial velocity $\dot{\gamma}_v(0) = v$. For every $p \in M$ and $v \in T_p M \setminus \{0\}$ there is a unique positive definite quadratic form g_v on $T_p M$ such that g_v and $F^2|_{T_p M}$ agree to second order at v . With the Chern connection, we can define the covariant derivative $D_{\dot{\gamma}_v(t)} J(t)$ of a vector field $J(t)$ along a geodesic $\gamma_v(t)$. A vector field $J(t)$ along $\gamma_v(t)$ is called a *Jacobi field* if it satisfies

$$D_{\dot{\gamma}_v(t)} D_{\dot{\gamma}_v(t)} J(t) + R_{\dot{\gamma}_v(t)}(J(t)) = 0,$$

where R is the curvature tensor.

The point $\gamma(t_0)$ is said to be *conjugate* to $\gamma(0)$ along geodesic γ , if there exists a Jacobi field J along γ , not identically zero, with $J(0) = 0 = J(t_0)$. We have the stable Jacobi tensors $C_v(t)$ with $C_v(0) = Id$ of Finsler manifolds without conjugate points as in the case of Riemannian manifolds (see [6, 8]). Making the change of variables

$$U_v(t) := (\ln C_v(t))' = C_v'(t) \cdot C_v^{-1}(t)$$

for t values for which $\det C_v(t) \neq 0$. In fact, if $\gamma_v(t)$ has no points conjugate to $\gamma(0)$ on $(0, \infty)$, then $U_v(t)$ is defined for all $t \in (0, \infty)$. Then the tensor U defined on the unit tangent bundle SM such that for every $v \in SM$, $U_v(t)$ is a self-adjoint linear operator on

$$\dot{\gamma}_v(t)^\perp := \{w \in T_{\gamma_v(t)} M \mid g_{\dot{\gamma}_v(t)}(\dot{\gamma}_v(t), w) = 0\}$$

and satisfies the *Riccati equation*

$$(1.1) \quad U_v'(t) + U_v^2(t) + R_v(t) = 0.$$

We will let $u(v) := \text{tr}(U_v(0))$ be the *mean curvature* (the trace of the second fundamental form) of the stable horosphere through v in the universal covering space. It is also known that $u(v)$ are measurable function, and the volume entropy

$$(1.2) \quad h_\mu = \int_{SM} u(v) d\mu(v).$$

2. Harmonic Finsler manifolds

A compact Finsler manifold is called a *Blaschke* manifold if every minimal geodesic of length less than the diameter is the unique shortest path between any of its points. Equivalently, for which all cut loci are round spheres of constant radius and dimension.

Remark 2.1. For a Blaschke manifold the exponential map restricted to the unit tangent sphere defines a great sphere foliation. Since every great sphere foliation of sphere is homeomorphic to a Hopf fibration, a simply connected Blaschke manifold is actually homeomorphic to compact rank one symmetric spaces.

A complete Finsler manifold is called *harmonic* if the mean curvature of all geodesic spheres is a function depending only on the radius. A historical break in the theory of harmonic Riemannian manifolds was made by Allamigeon when he proved the following: A simply connected harmonic Riemannian manifold is either diffeomorphic to Euclidean space or is a Blaschke manifold. The following theorem is to put them in a Finsler-geometric setting. For the sake of completeness we sketch the proof.

Theorem 2.2. *A simply connected harmonic Finsler manifold M is either diffeomorphic to Euclidean space or is a Blaschke manifold.*

Proof. Suppose there is no conjugate points. Then the exponential map is a covering map and since M is simply connected, a diffeomorphism. So take a $0 \neq v_0 \in T_x M$ and an $r_0 \in \mathbb{R}$ such that the first conjugate point along γ_{v_0} is $\gamma_{v_0}(r_0)$. Then the first conjugate point along γ_v is $\gamma_v(r_0)$ for all $v \in T_x M$, since the mean curvature is radial. Note that r_0 is the same for every point in M . This means that M is a Blaschke manifold by the Allamigeon-Warner theorem, cf. [1, Corollary 5.31]. \square

Therefore, noncompact harmonic Finsler manifolds have nonnegative constant mean curvature of the stable horospheres. Hence, we try to classify harmonic Finsler manifolds according to mean curvature stable horospheres. A harmonic Finsler manifold having stable horospheres of zero mean curvature is called as *manifold with minimal horospheres*.

The following proposition follows from standard techniques.

Proposition 2.3. *Let M be a complete noncompact harmonic Finsler manifold with minimal horospheres. Then for all $v \in SM$ we have*

$$U_v(0) = -U_{-v}(0).$$

Proof. Since M has no conjugate points, it is clear that the tensor $U_v(0) + U_{-v}(0)$ is nonpositive. Therefore we have

$$\operatorname{tr} \left(U_v(0) + U_{-v}(0) \right) \leq 0.$$

Furthermore M have minimal horospheres, we have $\text{tr } U_v(0) = 0$ for all $v \in SM$, and hence

$$\text{tr} \left(U_v(0) + U_{-v}(0) \right) = 0.$$

Since $U_v(0) + U_{-v}(0) \leq 0$ is symmetric negative semi-definite tensor, we can conclude that the tensor itself must be zero. \square

The above proposition shows that the stable and unstable horospheres of harmonic Finsler manifolds with minimal horospheres coincide like flat spaces.

The following lemma is the final ingredient needed for main theorem.

Lemma 2.4. *If M is as above, then for every point $x \in M$, there exists $v \in S_x M$ such that $U_v(0) = 0$.*

Proof. Since $U_v(0)$ is a symmetric matrix on v^\perp , there exists a basis of v^\perp such that $U_v(0)$ is represented by a diagonal matrix. Let $\lambda_1(v) \leq \lambda_2(v) \leq \dots \leq \lambda_{n-1}(v)$ be the eigenvalues of $U_v(0)$. Now consider the continuous function $f : S^{n-1} \rightarrow \mathbb{R}^{n-1}$ defined by

$$f(v) = (\lambda_1(v), \lambda_2(v), \dots, \lambda_{n-1}(v)).$$

Then by the Borsuk-Ulam theorem, there exists $v \in S^{n-1}$ such that $f(v) = f(-v)$. Therefore $\lambda_i(v) = \lambda_i(-v)$ for all $i = 1, 2, \dots, n-1$. Hence $U_v(0) = U_{-v}(0)$, and by Proposition 2.3, we have $U_v(0) = 0$. \square

Now we are ready to prove main theorem using Lemma 2.4.

Theorem 2.5. *Complete noncompact harmonic Finsler manifolds with minimal horospheres are flat.*

Proof. By Lemma 2.4, we have $U_v(0) = 0$. Inserting this back into the Riccati equation (1.1), we end up with $R_v(0) = 0$, which shows that given underline manifolds are flat, finishing the proof. \square

By using Theorem 2.5, we obtain the following Corollary.

Corollary. *If M is a compact harmonic Finsler manifold without conjugate points such that the volume entropy is zero, then M is flat.*

Proof. Since M have nonnegative constant mean curvature of the stable horospheres and the left-hand side of equation (1.2) has zero, we have that the integral term on the right-hand side of equation (1.2) has zero. This mean that the universal covering space \tilde{M} of M is a complete noncompact harmonic Finsler manifold with minimal horospheres. By Theorem 2.5, \tilde{M} is flat, we can conclude that M itself must be flat. \square

References

- [1] A. L. Besse, *Manifolds all of whose geodesics are closed*, Ergebnisse der Mathematik und ihrer Grenzgebiete, **93**, Springer-Verlag, Berlin, 1978.
- [2] G. Besson, G. Courtois, and S. Gallot, *Entropies et rigidités des espaces localement symétriques de courbure strictement négative*, Geom. Funct. Anal. **5** (1995), no. 5, 731–799.
- [3] D. Burago and S. Ivanov, *Riemannian tori without conjugate points are flat*, Geom. Funct. Anal. **4** (1994), no. 3, 259–269.
- [4] ———, *On asymptotic volume of Finsler tori, minimal surfaces in normed spaces, and symplectic filling volume*, Ann. of Math. (2) **156** (2002), no. 3, 891–914.
- [5] E. Damek and F. Ricci, *A class of nonsymmetric harmonic Riemannian spaces*, Bull. Amer. Math. Soc. (N.S.) **27** (1992), no. 1, 139–142.
- [6] P. Foulon, *Estimation de l'entropie des systèmes lagrangiens sans points conjugués*, Ann. Inst. H. Poincaré Phys. Théor. **57** (1992), no. 2, 117–146.
- [7] A. Freire and R. Mañé, *On the entropy of the geodesic flow in manifolds without conjugate points*, Invent. Math. **69** (1982), no. 3, 375–392.
- [8] C.-W. Kim, *Finsler manifolds without conjugate points and with integral Ricci curvature*, Israel J. Math. **189** (2012), 135–146.
- [9] ———, *Compact Finsler manifolds without focal points*, Arch. Math. (Basel) **99** (2012), no. 4, 387–392.
- [10] A. Ranjan and H. Shah, *Harmonic manifolds with minimal horospheres*, J. Geom. Anal. **12** (2002), no. 4, 683–694.
- [11] Z. Shen, *Lectures on Finsler Geometry*, World Scientific Publishing Co., Singapore, 2001.
- [12] Z. I. Szabò, *The Lichnerowicz conjecture on harmonic manifolds*, J. Differential Geom. **31** (1990), no. 1, 1–28.
- [13] N. Zinov'ev, *Examples of Finsler metrics without conjugate points: metrics of rotation*, St. Petersburg Math. J. **20** (2009), 361–379.

CHANG-WAN KIM
DIVISION OF LIBERAL ARTS AND SCIENCES
MOKPO NATIONAL MARITIME UNIVERSITY
MOKPO 58628, KOREA
Email address: cwkim@mmu.ac.kr