

TUBES OF FINITE CHEN-TYPE

HASSAN AL-ZOUBI, KHALID M. JABER, AND STYLIANOS STAMATAKIS

ABSTRACT. In this paper, we consider surfaces in the 3-dimensional Euclidean space \mathbb{E}^3 which are of finite *III*-type, that is, they are of finite type, in the sense of B.-Y. Chen, corresponding to the third fundamental form. We present an important family of surfaces, namely, tubes in \mathbb{E}^3 . We show that tubes are of infinite *III*-type.

1. Introduction

Let M^n be a (connected) submanifold in the m -dimensional Euclidean space \mathbb{E}^m . Let \mathbf{x}, \mathbf{H} be the position vector field and the mean curvature field of M^n respectively. Denote by Δ^I the second Beltrami-Laplace operator corresponding to the first fundamental form I of M^n . Then, it is well known that [7]

$$\Delta^I \mathbf{x} = -n\mathbf{H}.$$

From this formula one can see that M^n is a minimal submanifold if and only if all coordinate functions, restricted to M^n , are eigenfunctions of Δ^I with eigenvalue $\lambda = 0$. Moreover in [17] T. Takahashi showed that the submanifold M^n for which $\Delta^I \mathbf{x} = \lambda \mathbf{x}$, i.e., for which all coordinate functions are eigenfunctions of Δ^I with the same eigenvalue $\lambda \in \mathbb{R}$, are precisely either the minimal submanifold with eigenvalue $\lambda = 0$ or the minimal submanifold of hyperspheres S^{m-1} with eigenvalue $\lambda > 0$. Although the class of finite type submanifolds in an arbitrary dimensional Euclidean spaces is very large, very little is known about surfaces of finite type in the Euclidean 3-space \mathbb{E}^3 . Actually, so far, the only known surfaces of finite type corresponding to the first fundamental form in the Euclidean 3-space are the minimal surfaces, the circular cylinders and the spheres. So in [5] B.-Y. Chen mentions the following problem

Problem 1. Determine all surfaces of finite Chen *I*-type in \mathbb{E}^3 .

In order to provide an answer to the above problem, important families of surfaces were studied by different authors by proving that finite type ruled surfaces [9], finite type quadrics [8], finite type tubes [4], finite type cyclides

Received May 26, 2017; Revised July 12, 2017; Accepted July 26, 2017.

2010 *Mathematics Subject Classification.* 53A05.

Key words and phrases. surfaces in the Euclidean 3-space, surfaces of finite Chen-type, Beltrami operator.

frame. Further, we continue our study by proving finite type surfaces for an important class of surfaces, namely, tubes in \mathbb{E}^3 .

2. Preliminaries

Let S be a (connected) surface in the Euclidean 3-space \mathbb{E}^3 , whose Gaussian curvature K never vanishes. Let $\varphi = \{\varepsilon_1(u, v), \varepsilon_2(u, v), \varepsilon_3(u, v)\}$ be a moving frame of S , $\varepsilon_3 = \mathbf{n}$ be the Gauss map of S and $\det(\varepsilon_1, \varepsilon_2, \varepsilon_3) = 1$. Then it is well known that there are linear differential forms $\omega_1, \omega_2, \omega_{31}, \omega_{32}$ and ω_{12} , such that [12]

$$\begin{aligned} d\mathbf{x} &= \omega_1\varepsilon_1 + \omega_2\varepsilon_2, & d\mathbf{n} &= \omega_{31}\varepsilon_1 + \omega_{32}\varepsilon_2, \\ d\varepsilon_1 &= \omega_{12}\varepsilon_2 - \omega_{31}\varepsilon_3, & d\varepsilon_2 &= -\omega_{12}\varepsilon_1 - \omega_{32}\varepsilon_3, \end{aligned}$$

and functions a, b, c, q_1, q_2 of u, v such that

$$\omega_{31} = -a\omega_1 - b\omega_2, \quad \omega_{32} = -b\omega_1 - c\omega_2, \quad \omega_{12} = q_1\omega_1 + q_2\omega_2.$$

We can choose the moving frame of S , such that the vectors $\varepsilon_1, \varepsilon_2$ are the principle directions of S . Then a, c are the principle curvatures of S and $b = 0$, so the differential forms ω_1 and ω_2 become

$$\omega_1 = -\frac{1}{a}\omega_{31}, \quad \omega_2 = -\frac{1}{c}\omega_{32}.$$

The Gauss and mean curvature are respectively

$$K = ac, \quad H = \frac{a+c}{2}.$$

Let $\nabla_1 f, \nabla_2 f$ be the derivatives of Pfaff of a function $f(u, v) \in C^1$ along the curves $\omega_2 = 0, \omega_1 = 0$ respectively. Then we have the following well known relations [3]

$$\begin{aligned} \nabla_1 \mathbf{x} &= \varepsilon_1, & \nabla_2 \mathbf{x} &= \varepsilon_2, \\ \nabla_1 \varepsilon_1 &= q_1 \varepsilon_2 + a\mathbf{n}, & \nabla_2 \varepsilon_1 &= q_2 \varepsilon_2, \\ \nabla_1 \varepsilon_2 &= -q_1 \varepsilon_1, & \nabla_2 \varepsilon_2 &= -q_2 \varepsilon_1 + c\mathbf{n}, \\ \nabla_1 \mathbf{n} &= -a\varepsilon_1, & \nabla_2 \mathbf{n} &= -c\varepsilon_2, \end{aligned}$$

We denote by $\tilde{\nabla}_1 f$ and $\tilde{\nabla}_2 f$ the derivatives of Pfaff of f along the curves $\omega_{32} = 0, \omega_{31} = 0$ respectively. One finds

$$\tilde{\nabla}_1 f = -\frac{1}{a}\nabla_1 f, \quad \tilde{\nabla}_2 f = -\frac{1}{c}\nabla_2 f.$$

It follows that

$$(2.1) \quad \tilde{\nabla}_1 \mathbf{x} = -\frac{1}{a}\varepsilon_1, \quad \tilde{\nabla}_2 \mathbf{x} = -\frac{1}{c}\varepsilon_2,$$

$$(2.2) \quad \tilde{\nabla}_1 \varepsilon_1 = p_1 \varepsilon_2 - \mathbf{n}, \quad \tilde{\nabla}_2 \varepsilon_1 = p_2 \varepsilon_2,$$

$$(2.3) \quad \tilde{\nabla}_1 \varepsilon_2 = -p_1 \varepsilon_1, \quad \tilde{\nabla}_2 \varepsilon_2 = -p_2 \varepsilon_1 - \mathbf{n}, \\ \tilde{\nabla}_1 \mathbf{n} = \varepsilon_1, \quad \tilde{\nabla}_2 \mathbf{n} = \varepsilon_2,$$

where $p_1 = -\frac{1}{a}q_1$, $p_2 = -\frac{1}{c}q_2$ are the geodesic curvatures of the spherical curves $\omega_{32} = 0$ and $\omega_{31} = 0$ respectively. The Mainardi-Codazzi equations have the following forms

$$(2.4) \quad \tilde{\nabla}_1 \frac{1}{c} = p_2 \left(\frac{1}{a} - \frac{1}{c} \right), \quad \tilde{\nabla}_2 \frac{1}{a} = p_1 \left(\frac{1}{a} - \frac{1}{c} \right).$$

Let f be a sufficient differentiable function on S . Then the second differential parameter of Beltrami corresponding to the fundamental form III of S is defined by

$$(2.5) \quad \Delta^{III} f = -\tilde{\nabla}_1 \tilde{\nabla}_1 f - \tilde{\nabla}_2 \tilde{\nabla}_2 f - p_2 \tilde{\nabla}_1 f + p_1 \tilde{\nabla}_2 f.$$

Applying (2.5) to the position vector \mathbf{x} gives

$$\Delta^{III} \mathbf{x} = -\tilde{\nabla}_1 \tilde{\nabla}_1 \mathbf{x} - \tilde{\nabla}_2 \tilde{\nabla}_2 \mathbf{x} - p_2 \tilde{\nabla}_1 \mathbf{x} + p_1 \tilde{\nabla}_2 \mathbf{x}.$$

From (2.1) we obtain

$$(2.6) \quad \Delta^{III} \mathbf{x} = \tilde{\nabla}_1 \left(\frac{1}{a} \boldsymbol{\varepsilon}_1 \right) + \tilde{\nabla}_2 \left(\frac{1}{c} \boldsymbol{\varepsilon}_2 \right) + \frac{1}{a} p_2 \boldsymbol{\varepsilon}_1 - \frac{1}{c} p_1 \boldsymbol{\varepsilon}_2.$$

Using (2.2) and (2.3), equation (2.6) becomes

$$(2.7) \quad \Delta^{III} \mathbf{x} = \left(\tilde{\nabla}_1 \frac{1}{a} \right) \boldsymbol{\varepsilon}_1 + \left(\frac{1}{a} - \frac{1}{c} \right) p_2 \boldsymbol{\varepsilon}_1 + \left(\tilde{\nabla}_2 \frac{1}{c} \right) \boldsymbol{\varepsilon}_2 + \left(\frac{1}{a} - \frac{1}{c} \right) p_1 \boldsymbol{\varepsilon}_2 - \left(\frac{1}{a} + \frac{1}{c} \right) \mathbf{n}.$$

Taking into account the Mainardi-Codazzi equations (2.4), so equation (2.7) reduces to

$$\Delta^{III} \mathbf{x} = \left(\tilde{\nabla}_1 \left(\frac{1}{a} + \frac{1}{c} \right) \right) \boldsymbol{\varepsilon}_1 + \left(\tilde{\nabla}_2 \left(\frac{1}{a} + \frac{1}{c} \right) \right) \boldsymbol{\varepsilon}_2 - \left(\frac{1}{a} + \frac{1}{c} \right) \mathbf{n}$$

or equivalently, (see [15])

$$(2.8) \quad \Delta^{III} \mathbf{x} = \text{grad}^{III} \left(\frac{2H}{K} \right) - \left(\frac{2H}{K} \right) \mathbf{n}.$$

Remark 1. S. Stamatakis and H. Al-Zoubi proved in [15] relation (2.8) by using tensors calculus.

From (2.8) the following results were proved in [15].

Theorem 1. *A surface S in \mathbb{E}^3 is of 0-type 1 corresponding to the third fundamental form if and only if S is minimal.*

Theorem 2. *A surface S in \mathbb{E}^3 is of III-type 1 if and only if S is part of a sphere.*

Corollary 1. *The Gauss map of every surface S in \mathbb{E}^3 is of III-type 1. The corresponding eigenvalue is $\lambda = 2$.*

Up to now, the only known surfaces of finite III-type in \mathbb{E}^3 are parts of spheres, the minimal surfaces and the parallel of the minimal surfaces which are of null III-type 2. So the following question seems to be interesting:

Problem 2. Other than the surfaces mentioned above, which surfaces in \mathbb{E}^3 are of finite *III*-type?

Another generalization of the above problem is to study surfaces in \mathbb{E}^3 with the position vector \mathbf{x} satisfying

$$(2.9) \quad \Delta^{III} \mathbf{x} = A\mathbf{x},$$

where $A \in \mathbb{R}^{3 \times 3}$.

From this point of view, we also pose the following problem.

Problem 3. Classify all surfaces in \mathbb{E}^3 with the position vector \mathbf{x} satisfying relation (2.9).

Concerning this problem, in [16] S. Stamatakis and H. Al-Zoubi studied the class of surfaces of revolution and they proved that: A surface of revolution S satisfies (2.9) if and only if S is a catenoid or part of a sphere. Recently, the same authors in [1] studied the class of ruled surfaces and the class of quadric surfaces. In particular, they proved that helicoids and spheres are the only ruled and quadric surfaces satisfying (2.9) respectively.

This paper provides the first attempt at the study of finite type families of surfaces in \mathbb{E}^3 corresponding to the third fundamental form. Our main result is the following.

Theorem 3. *All tubes in \mathbb{E}^3 are of infinite III-type.*

Our discussion is local, which means that we show in fact that any open part of a tube is of infinite Chen type.

3. Tubes in \mathbb{E}^3

Let $C : \boldsymbol{\alpha} = \boldsymbol{\alpha}(t)$, $t \in (a, b)$ be a regular unit speed curve of finite length which is topologically imbedded in \mathbb{E}^3 . The total space $N_{\boldsymbol{\alpha}}$ of the normal bundle of $\boldsymbol{\alpha}((a, b))$ in \mathbb{E}^3 is naturally diffeomorphic to the direct product $(a, b) \times \mathbb{E}^2$ via the translation along $\boldsymbol{\alpha}$ with respect to the induced normal connection. For a sufficiently small $r > 0$ the tube of radius r about the curve $\boldsymbol{\alpha}$ is the set:

$$T_r(\boldsymbol{\alpha}) = \{\exp_{\boldsymbol{\alpha}(t)} \mathbf{u} \mid \mathbf{u} \in N_{\boldsymbol{\alpha}}, \quad \|\mathbf{u}\| = r, \quad t \in (a, b)\}.$$

Assume that $\mathbf{t}, \mathbf{h}, \mathbf{b}$ is the Frenet frame and κ the curvature of the unit speed curve $\boldsymbol{\alpha} = \boldsymbol{\alpha}(t)$. For a small real number r satisfies $0 < r < \min \frac{1}{|\kappa|}$, the tube $T_r(\boldsymbol{\alpha})$ is a smooth surface in \mathbb{E}^3 , [14]. Then a parametric representation of the tube $T_r(\boldsymbol{\alpha})$ is given by

$$(3.1) \quad F : \mathbf{x}(t, \varphi) = \boldsymbol{\alpha} + r \cos \varphi \mathbf{h} + r \sin \varphi \mathbf{b}.$$

It is easily verified that the first and the second fundamental forms of F are given by

$$\begin{aligned} I &= (\delta^2 + r^2 \tau^2) dt^2 + 2r^2 \tau dt d\varphi + r^2 d\varphi^2, \\ II &= (-\kappa \delta \cos \varphi + r\tau^2) dt^2 + 2r\tau dt d\varphi + rd\varphi^2, \end{aligned}$$

where $\delta := (1 - r\kappa \cos \varphi)$ and τ is the torsion of the curve α . The Gauss curvature of F is given by

$$(3.2) \quad K = -\frac{\kappa \cos \varphi}{r\delta}.$$

Notice that $\kappa \neq 0$ since the Gauss curvature vanishes. The Beltrami operator corresponding to the third fundamental form of F can be expressed as follows

$$(3.3) \quad \Delta^{III} = \frac{1}{(\kappa \cos \varphi)^2} \left[-\frac{\partial^2}{\partial t^2} + 2\tau \frac{\partial^2}{\partial t \partial \varphi} - (\tau^2 + \kappa^2 \cos^2 \varphi) \frac{\partial^2}{\partial \varphi^2} + \frac{\beta}{\kappa \cos \varphi} \frac{\partial}{\partial t} + \left(\tau' + \kappa^2 \cos \varphi \sin \varphi - \frac{\tau \beta}{\kappa \cos \varphi} \right) \frac{\partial}{\partial \varphi} \right],$$

where $\beta := \kappa' \cos \varphi + \kappa \tau \sin \varphi$ and $' := \frac{d}{dt}$.

Before we start of proving our main result, we mention and prove the following special case of tubular surfaces for later use.

3.1. Anchor rings

A tube in the Euclidean 3-space is called an anchor ring if the curve C is a plane circle (or is an open portion of a plane circle). In this case, the torsion τ of α vanishes identically and the curvature κ of α is a nonzero constant. Then the position vector \mathbf{x} of the anchor ring can be expressed as

$$(3.4) \quad F : \mathbf{x}(t, \varphi) = \{(a + r \cos t) \cos \varphi, (a + r \cos t) \sin \varphi, r \sin t\},$$

$$a > r, \quad a \in \mathbb{R}.$$

The first fundamental form is

$$I = r^2 dt^2 + (a + r \cos t)^2 d\varphi^2,$$

while the second is

$$II = r dt^2 + (a + r \cos t) \cos t d\varphi^2.$$

Hence, the Beltrami operator is given by

$$(3.5) \quad \Delta^{III} = -\frac{\partial^2}{\partial t^2} + \frac{\sin t}{\cos t} \frac{\partial}{\partial t} - \frac{1}{\cos^2 t} \frac{\partial^2}{\partial \varphi^2}.$$

Let x_1 be the first coordinate function of \mathbf{x} . By virtue of (3.5) one can find

$$(3.6) \quad \Delta^{III} x_1 = \frac{1}{\cos^2 t} a \cos \varphi + 2r \cos t \cos \varphi.$$

Moreover, by a direct computation, we obtain

$$(3.7) \quad (\Delta^{III})^2 x_1 = \left(\frac{2}{\cos^2 t} - \frac{3}{\cos^4 t} \right) a \cos \varphi + 4r \cos t \cos \varphi,$$

$$(3.8) \quad (\Delta^{III})^3 x_1 = \left(\frac{4}{\cos^2 t} - \frac{42}{\cos^4 t} + \frac{45}{\cos^6 t} \right) a \cos \varphi + 8r \cos t \cos \varphi.$$

It can be seen that $\Delta^{III}(\cos t \cos \varphi) = 2 \cos t \cos \varphi$, and for each integer $k > 0$, it is easy to see that

$$(3.9) \quad \Delta^{III} \frac{\cos \varphi}{\cos^k t} = \left(k^2 - k - \frac{k^2 - 1}{\cos^2 t} \right) \frac{\cos \varphi}{\cos^k t}.$$

Thus, by induction, one finds

$$(3.10) \quad (\Delta^{III})^m x_1 = \left(\frac{d_{0,m}}{\cos^2 t} - \frac{d_{1,m}}{\cos^4 t} + \cdots + \frac{d_{m-1,m}}{\cos^{2m} t} \right) a \cos \varphi + 2^m r \cos t \cos \varphi,$$

where $d_{j,m}$ are constants, $j = 1, 2, \dots, m-1$, and

$$d_{0,m} = 2^{m-1}, \quad d_{m-1,m} = (-1)^{m-1} (2m-1) \prod_{j=1}^m (2j-3)^2.$$

Notice that $d_{m-1,m} \neq 0$, for each integer $m \geq 1$. Now, if F is of finite type, then there exist real numbers, c_1, c_2, \dots, c_m such that

$$(3.11) \quad (\Delta^{III})^m \mathbf{x} + c_1 (\Delta^{III})^{m-1} \mathbf{x} + \cdots + c_{m-1} \Delta^{III} \mathbf{x} + c_m \mathbf{x} = \mathbf{0}.$$

Since $x_1 = (a + r \cos t) \cos \varphi$ is the first coordinate of \mathbf{x} , (3.11), one gets

$$(3.12) \quad (\Delta^{III})^m x_1 + c_1 (\Delta^{III})^{m-1} x_1 + \cdots + c_{m-1} \Delta^{III} x_1 + c_m x_1 = 0.$$

From (3.6-3.8), (3.10) and (3.12) we obtain that

$$\begin{aligned} & 2^m r \cos t \cos \varphi + a \cos \varphi \sum_{j=1}^m \frac{d_{j-1,m}}{\cos^{2j} t} + 2^{m-1} c_1 r \cos t \cos \varphi \\ & + c_1 a \cos \varphi \sum_{j=1}^{m-1} \frac{d_{j-1,m-1}}{\cos^{2j} t} + \cdots + 2c_{m-1} r \cos t \cos \varphi \\ & + c_{m-1} \frac{a \cos \varphi}{\cos^2 t} + c_m (a + r \cos t) \cos \varphi = 0 \end{aligned}$$

which can be rewritten as

$$(3.13) \quad \frac{d_{m-1,m}}{\cos^{2m} t} + \frac{1}{\cos^{2m-2} t} F(\cos t) = 0,$$

where $F(u)$ is a polynomial in $u = \cos t$ of degree $2m-2$.

This is impossible for any $m \geq 1$ since $d_{m-1,m} \neq 0$. Consequently, we have the following

Corollary 2. *Every anchor ring in the Euclidean 3-space is of infinite type.*

4. Proof of the main theorem

Applying relation (3.3) on the position vector \mathbf{x} of (3.1) gives

$$\Delta^{III} \mathbf{x} = \frac{\beta}{\kappa^3 \cos^3 \varphi} \mathbf{t} + \left(2r \cos \varphi - \frac{1}{\kappa \cos^2 \varphi} \right) \mathbf{h} + 2r \sin \varphi \mathbf{b},$$

which can be rewritten as

$$(4.1) \quad \Delta^{III} \mathbf{x} = \frac{\beta}{\kappa^3 \cos^3 \varphi} \mathbf{t} + \frac{1}{\kappa^2 \cos^2 \varphi} \mathbf{P}_1(\cos \varphi, \sin \varphi),$$

where $\mathbf{P}_1(u, v)$ is a vector valued polynomial in u, v of degree 3 with functions in t as coefficients. Moreover, by a long computation, we obtain

$$(4.2) \quad (\Delta^{III})^2 \mathbf{x} = \frac{\beta^3}{\kappa^7 \cos^7 \varphi} \mathbf{t} + \frac{1}{\kappa^6 \cos^6 \varphi} \mathbf{P}_2(\cos \varphi, \sin \varphi),$$

where $\mathbf{P}_2(u, v)$ is a vector valued polynomial in u, v of degree 7 with functions in t as coefficients.

We need the following lemma which can be proved directly by using (3.3).

Lemma 1. *For any natural numbers m and n we have*

$$\left(\Delta^{III} \frac{\beta^m}{(\kappa \cos \varphi)^n} \right) = -\frac{n(n+2)\beta^{m+2}}{(\kappa \cos \varphi)^{n+4}} + \frac{1}{(\kappa \cos \varphi)^{n+3}} P(\cos \varphi, \sin \varphi),$$

where P is a polynomial in u, v of degree $n+4$ with functions in t as coefficients.

Using Lemma 1 and relation (3.3) one finds

$$(4.3) \quad (\Delta^{III})^\lambda \mathbf{x} = d_\lambda \frac{\beta^{2\lambda-1}}{(\kappa \cos \varphi)^{4\lambda-1}} \mathbf{t} + \frac{1}{(\kappa \cos \varphi)^{4\lambda-2}} \mathbf{P}_\lambda(\cos \varphi, \sin \varphi),$$

where

$$d_\lambda = (-1)^{\lambda-1} \prod_{j=1}^{2\lambda-1} (2j-1).$$

It can be seen that $d_\lambda \neq 0$, for each natural number λ . Moreover, we have

$$(4.4) \quad (\Delta^{III})^{\lambda+1} \mathbf{x} = d_{\lambda+1} \frac{\beta^{2\lambda+1}}{(\kappa \cos \varphi)^{4\lambda+3}} \mathbf{t} + \frac{1}{(\kappa \cos \varphi)^{4\lambda+2}} \mathbf{P}_{\lambda+1}(\cos \varphi, \sin \varphi).$$

Let F be of finite type. Then there exist real numbers, $c_1, c_2, \dots, c_\lambda$ such that

$$(4.5) \quad (\Delta^{III})^{\lambda+1} \mathbf{x} + c_1 (\Delta^{III})^\lambda \mathbf{x} + \dots + c_\lambda \Delta^{III} \mathbf{x} = \mathbf{0}.$$

Using (4.1-4.4), one has

$$(4.6) \quad d_{\lambda+1} \frac{\beta^{2\lambda+1}}{\kappa \cos \varphi} \mathbf{t} = Q_1 \mathbf{t} + Q_2 \mathbf{h} + Q_3 \mathbf{b},$$

where $Q_i, i = 1, 2, 3$, are polynomials in u, v with functions in t as coefficients.

Now, if $\beta \neq 0$. From (4.6) we find

$$(4.7) \quad d_{\lambda+1} \frac{\beta^{2\lambda+1}}{\kappa \cos \varphi} = Q_1(\cos \varphi, \sin \varphi).$$

This is impossible, since Q_1 is polynomial in $\cos \varphi$ and $\sin \varphi$. Assume now $\beta = 0$. Then $\kappa' = 0$ and $\kappa\tau = 0$ so $\kappa = \text{const.} \neq 0$ and $\tau = 0$. Therefore

the curve C is a circle, and so F is anchor ring. Hence, F is of infinite type according to Corollary (2). This completes our proof.

Acknowledgement. The authors would like to express their thanks to the referee for his useful remarks.

References

- [1] H. Al-Zoubi and S. Stamatakis, *Ruled and Quadric surfaces satisfying $\Delta^{III} \mathbf{x} = A\mathbf{x}$* , J. Geom. Graph. **20** (2016), no. 2, 147–157.
- [2] C. Baikoussis and L. Verstraelen, *The Chen-type of the spiral surfaces*, Results Math. **28** (1995), no. 3-4, 214–223.
- [3] W. Blaschke and K. Leichtwiss, *Elementare Differentialgeometrie*, Springer, Berlin, 1973.
- [4] B.-Y. Chen, *Surfaces of finite type in Euclidean 3-space*, Bull. Soc. Math. Belg. Sér B **39** (1987), no. 2, 243–254.
- [5] ———, *Some open problems and conjectures on submanifolds of finite type*, Soochow J. Math. **17** (1991), no. 2, 169–188.
- [6] ———, *A report on submanifolds of finite type*, Soochow J. Math. **22** (1996), no. 2, 117–337.
- [7] ———, *Total Mean Curvature and Submanifolds of Finite Type*, 2nd edition. World Scientific Publisher, 2015.
- [8] B.-Y. Chen and F. Dillen, *Quadrics of finite type*, J. Geom. **38** (1990), no. 1-2, 16–22.
- [9] B.-Y. Chen, F. Dillen, L. Verstraelen, and L. Vrancken, *Ruled surfaces of finite type*, Bull. Austral. Math. Soc. **42** (1990), no. 3, 447–453.
- [10] F. Defever, R. Deszcz, and L. Verstraelen, *The compact cyclides of Dupin and a conjecture of B.-Y. Chen*, J. Geom. **46** (1993), no. 1-2, 33–38.
- [11] F. Dillen, J. Pas, and L. Verstraelen, *On surfaces of finite type in Euclidean 3-space*, Kodai Math. J. **13** (1990), no. 1, 10–21.
- [12] W. Haack, *Elementare Differentialgeometrie, Basel und Stuttgart*, Birkhäuser 1955.
- [13] O. Garay, *An extension of Takahashi's theorem*, Geom. Dedicata **34** (1990), no. 2, 105–112.
- [14] J. S. Ro and D. W. Yoon, *Tubes of Weingarten types in Euclidean 3-space*, J. Chungcheong Math. Soc. **22** (2009), 359–366.
- [15] S. Stamatakis and H. Al-Zoubi, *On surfaces of finite Chen-type*, Results. Math. **43** (2003), no. 1-2, 181–190.
- [16] ———, *Surfaces of revolution satisfying $\Delta^{III} \mathbf{x} = A\mathbf{x}$* , J. Geom. Graph. **14** (2010), no. 2, 181–186.
- [17] T. Takahashi, *Minimal immersions of Riemannian manifolds*, J. Math. Soc. Japan **18** (1966), 380–385.

HASSAN AL-ZOUBI
 DEPARTMENT OF MATHEMATICS
 AL-ZAYTOONAH UNIVERSITY OF JORDAN
 P.O. BOX 130, AMMAN, JORDAN 11733
E-mail address: dr.hassanz@zuj.edu.jo

KHALID M. JABER
 AL-ZAYTOONAH UNIVERSITY OF JORDAN
 DEPARTMENT OF COMPUTER SCIENCE
E-mail address: K.Jaber@zuj.edu.jo

STYLIANOS STAMATAKIS
DEPARTMENT OF MATHEMATICS
ARISTOTLE UNIVERSITY OF THESSALONIKI
E-mail address: `stamata@math.auth.gr`

Prehead of Print