

GRADED PRIMITIVE AND INC-EXTENSIONS

HALEH HAMDI AND PARVIZ SAHANDI

ABSTRACT. It is well-known that quasi-Prüfer domains are characterized as those domains D , such that every extension of D inside its quotient field is a primitive extension and that primitive extensions are characterized in terms of INC-extensions.

Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain graded by an arbitrary torsionless grading monoid Γ and \star be a semistar operation on R . The main purpose of this paper is to give new characterizations of $\text{gr}\text{-}\star$ -quasi-Prüfer domains in terms of graded primitive and INC-extensions. Applications include new characterizations of $\text{UM}t$ -domains.

1. Introduction

Let D be a (commutative) integral domain with quotient field $qf(D)$. Recall that D is called a *quasi-Prüfer domain* if D has Prüfer integral closure [8], and as a t -operation analogue, D is called a *UMt-domain* if every upper to zero in the polynomial ring $D[X]$ is a maximal t -ideal [12]. Gilmer and Hoffmann characterized quasi-Prüfer domains as those domains D , such that the embedding $D \subseteq qf(D)$ is a primitive-extension [10, Theorem 2], and Dobbs [6] characterized primitive-extensions in terms of INC-domains.

Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded (commutative) integral domain graded by an arbitrary grading torsionless monoid Γ . In [11] the authors studied quasi-Prüfer and $\text{UM}t$ -domain properties of graded integral domains. For this reason they introduced the graded analogue of \star -quasi-Prüfer domains [4] called $\text{gr}\text{-}\star$ -quasi-Prüfer domains. The graded integral domain R is called a *gr- \star -quasi-Prüfer domain* in case, if Q is a prime ideal in $R[X]$ and $Q \subseteq P[X]$, for some homogeneous quasi- \star -prime ideal P of R , then $Q = (Q \cap R)[X]$. When $\star = d$ the identity operation on R , then we call the $\text{gr}\text{-}d$ -quasi-Prüfer domain a *gr-quasi-Prüfer domain*. It is shown that R is a $\text{gr}\text{-}\star$ -quasi-Prüfer domain if and only if R_P is a quasi-Prüfer domain, for each homogeneous quasi- \star -prime ideal P of R [11, Proposition 2.2]. Also it is known that R is a $\text{UM}t$ -domain if and only if R is a $\text{gr}\text{-}t$ -quasi-Prüfer domain if and only if R_P is a quasi-Prüfer

Received May 31, 2017; Revised August 26, 2017; Accepted September 4, 2017.

2010 *Mathematics Subject Classification*. Primary 13A15, 13G05, 13A02, 13F05.

Key words and phrases. semistar operation, graded domain, $\text{UM}t$ -domain, primitive extension, INC-extension.

domain for each *homogeneous* prime t -ideal P of R [11, Theorem 3.2]. If \star is a (semi)star operation on R , then R is a $\text{gr-}\star_f$ -quasi-Prüfer domain if and only if R is a UMt-domain and $\tilde{\star}$ and w coincide on nonzero homogeneous ideals of R [11, Theorem 3.9]. In particular R is a gr- quasi-Prüfer domain if and only if R is a UMt-domain and d and w coincide on nonzero homogeneous ideals of R . (Relevant definitions are reviewed in the sequel.)

The main purpose of this paper is to give new characterizations of $\text{gr-}\star$ -quasi-Prüfer domains in terms of graded primitive extension and graded incomparable or INC-extension (see [3], [6] and [10]).

To facilitate the reading of the paper, we review some basic facts on semistar operations on (graded) integral domains. Let Γ be a nonzero torsionless grading monoid, that is, Γ is a torsionless commutative cancellative monoid (written additively), and $\langle \Gamma \rangle = \{a - b \mid a, b \in \Gamma\}$ be the quotient group of Γ ; so $\langle \Gamma \rangle$ is a torsionfree abelian group. It is known that a cancellative monoid is torsionless if and only if it can be given a total order compatible with the monoid operation [14, page 123]. Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a Γ -graded integral domain. That is, $\deg(x) = \alpha$ for each $0 \neq x \in R_\alpha$ and $\deg(0) = 0$, and thus each nonzero $f \in R$ can be written uniquely as $f = x_{\alpha_1} + \cdots + x_{\alpha_n}$ with $\deg(x_{\alpha_i}) = \alpha_i$ and $\alpha_1 < \cdots < \alpha_n$. A nonzero $x \in R_\alpha$ for all $\alpha \in \Gamma$ is said to be *homogeneous*, and so if $H = \bigcup_{\alpha \in \Gamma} (R_\alpha \setminus \{0\})$, then H is the saturated multiplicative set of nonzero homogeneous elements of R . Then $R_H = \bigoplus_{\alpha \in \langle \Gamma \rangle} (R_H)_\alpha$, called the *homogeneous quotient field of R* , is a $\langle \Gamma \rangle$ -graded integral domain whose nonzero homogeneous elements are units. An integral ideal I of R is said to be *homogeneous* if $I = \bigoplus_{\alpha \in \Gamma} (I \cap R_\alpha)$. A fractional ideal I of R is *homogeneous* if sI is an integral homogeneous ideal of R for some $s \in H$ (thus $I \subseteq R_H$). An overring T of R , with $R \subseteq T \subseteq R_H$ will be called a *homogeneous overring* if $T = \bigoplus_{\alpha \in \langle \Gamma \rangle} (T \cap (R_H)_\alpha)$. Thus T is a ($\langle \Gamma \rangle$)-graded integral domain with $T_\alpha = T \cap (R_H)_\alpha$ for all $\alpha \in \langle \Gamma \rangle$. For more on graded integral domains and their divisibility properties (see [1], [14]).

Let D be an integral domain with quotient field K . Let $\overline{\mathcal{F}}(D)$ denote the set of all nonzero D -submodules of K , $\mathcal{F}(D)$ be the set of all nonzero fractional ideals of D , and $f(D)$ be the set of all nonzero finitely generated fractional ideals of D . Obviously, $f(D) \subseteq \mathcal{F}(D) \subseteq \overline{\mathcal{F}}(D)$. As in [15], a *semistar operation on D* is a map $\star : \overline{\mathcal{F}}(D) \rightarrow \overline{\mathcal{F}}(D)$, $E \mapsto E^\star$, such that, for all $0 \neq x \in K$, and for all $E, F \in \overline{\mathcal{F}}(D)$, the following properties hold: (\star_1) $(xE)^\star = xE^\star$; (\star_2) $E \subseteq F$ implies that $E^\star \subseteq F^\star$; (\star_3) $E \subseteq E^\star$; and (\star_4) $E^{\star\star} := (E^\star)^\star = E^\star$.

A semistar operation \star is called a (*semi*)*star operation on D* , if $D^\star = D$. Let \star be a semistar operation on D . For every $E \in \overline{\mathcal{F}}(D)$, put $E^{\star_f} := \bigcup F^\star$, where the union is taken over all $F \in f(D)$ with $F \subseteq E$. It is easy to see that \star_f is a semistar operation on D . We say that a nonzero ideal I of D is a *quasi- \star -ideal* of D , if $I^\star \cap D = I$; a *quasi- \star -prime* (ideal of D), if I is a prime quasi- \star -ideal of D ; and a *quasi- \star -maximal* (ideal of D), if I is maximal in the set of all proper quasi- \star -ideals of D . Each quasi- \star -maximal ideal is a prime ideal. It is shown

in [7, Lemma 4.20] that if $D^* \neq K$, then each proper quasi- \star_f -ideal of D is contained in a quasi- \star_f -maximal ideal of D . We denote by $\text{QMax}^*(D)$ (resp., $\text{QSpec}^*(D)$) the set of all quasi- \star -maximal ideals (resp., quasi- \star -prime ideals) of D .

Given a semistar operation \star on D , it is possible to construct a semistar operation $\tilde{\star}$, which is defined as follows, for each $E \in \overline{\mathcal{F}}(D)$, $E^{\tilde{\star}} := \bigcap_{P \in \text{QMax}^{\star_f}(D)} ED_P$.

The most widely studied (semi)star operations on D have been the identity d_D , v_D , $t_D := (v_D)_f$, and $w_D := \tilde{v}_D$ operations, where $A^{v_D} := (A^{-1})^{-1}$, with $A^{-1} := (D : A) := \{x \in K \mid xA \subseteq D\}$. We usually use these operations without subscripts. If \star is a (semi)star operation on D , then $d \leq \star \leq v$.

Let \star be a semistar operation on a graded integral domain $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$. We say that \star is *homogeneous preserving* if \star sends homogeneous fractional ideals to homogeneous ones. It is known that d , t , and v are homogeneous preserving [1, Proposition 2.5], $\tilde{\star}$ is homogeneous preserving [16, Proposition 2.3], and that if \star is homogeneous preserving, then so is \star_f [16, Lemma 2.4]. Denote by $h\text{-QSpec}^*(R)$ the homogeneous ideals of $\text{QSpec}^*(R)$ and let $h\text{-QMax}^*(R)$ denote the set of ideals of R which are maximal in the set of all proper homogeneous quasi- \star -ideals of R (if \star is a (semi)star operation we denote these sets by $h\text{-Spec}^*(R)$ and $h\text{-Max}^*(R)$ respectively). It is shown that if $R^* \subsetneq R_H$ and $\star = \star_f$ homogeneous preserving, then $h\text{-QMax}^{\star_f}(R) (\subseteq h\text{-QSpec}^*(R))$ is nonempty, each proper homogeneous quasi- \star_f -ideal is contained in a homogeneous maximal quasi- \star_f -ideal [16, Lemma 2.1], and $h\text{-QMax}^{\star_f}(R) = h\text{-QMax}^{\tilde{\star}}(R)$ [16, Proposition 2.5].

2. Graded primitive extension

Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain with quotient field $qf(R)$, H be the set of nonzero homogeneous elements of R , and \star be a semistar operation on R such that $R^* \subsetneq R_H$. In this section we give a characterization of $gr\text{-}\star$ -quasi-Prüfer domains in terms of graded semistar primitive extensions.

For $a \in R$, denote by $C(a)$ the ideal of R generated by homogeneous components of a . The *homogeneous content ideal* for a polynomial $f = a_0 + a_1X + \cdots + a_nX^n \in R[X]$, is defined by $\mathcal{A}_f := \mathcal{A}_f^R := \sum_{i=0}^n C(a_i)$ [17]. It can be seen that if R has trivial grading, i.e., $\Gamma = \{0\}$, then \mathcal{A}_f coincides with the usual content ideal of f (that is the ideal generated by coefficients of f). Assume that L is a fractional ideal of $R[X]$ such that $L \subseteq R_H[X]$, and set $\mathcal{A}_L := \sum_{f \in L} \mathcal{A}_f$. Now assume that $R \subseteq T$ is an extension of graded integral domains such that each homogeneous element of R is a homogeneous element of T . We say that an element $u \in T$ is *gr- \star -primitive over R* if u is a root of a nonzero polynomial $g \in R[X]$ with $(\mathcal{A}_g^R)^\star = R^*$. The extension $R \subseteq T$ of graded integral domains is called a *gr- \star -primitive extension* if each homogeneous element of T is $gr\text{-}\star$ -primitive over R . We call the extension $R \subseteq T$ a *gr-primitive extension* if $R \subseteq T$ is a $gr\text{-}d_R$ -primitive extension. It is clear that if R and T have

trivial grading, then gr-primitive extension coincides with the usual primitive extension of Gilmer and Hoffmann [10].

Recall that R is said to be a *graded-Prüfer domain* if each nonzero finitely generated homogeneous ideal of R is invertible [2]. We say that R is a *graded valuation domain* (gr-valuation domain) if either $u \in R$ or $u^{-1} \in R$ for every nonzero homogeneous $u \in R_H$.

Proposition 2.1 ([16, Theorem 4.4]). *Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain. Then the following statements are equivalent.*

- (1) R is a graded-Prüfer domain.
- (2) R_P is a valuation domain for all $P \in h\text{-Spec}(R)$ (resp., $P \in h\text{-Max}(R)$).
- (3) $R_{H \setminus P}$ is a gr-valuation domain for all $P \in h\text{-Spec}(R)$ (resp., $P \in h\text{-Max}(R)$).

Proposition 2.2. *Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be an integrally closed graded domain. Then R is a graded-Prüfer domain if and only if R_H is a gr-primitive extension of R .*

Proof. Assume that R is a graded-Prüfer domain and let $u = a/b \in R_H$ be a nonzero homogeneous element, where $a, b \in H$. Then there exists an integer $n > 1$ such that $a^{n-1}b \in (a^n, b^n)$ by [16, Theorem 4.1]. It follows that $a^{n-1}b = r_1a^n + r_2b^n$ for some $r_1, r_2 \in R$; dividing both sides of this equation by b^n yields $f(X) = r_1X^n - X^{n-1} + r_2 \in R[X]$ with $f(u) = 0$ and $\mathcal{A}_f = R$, so u is gr-primitive over R .

Conversely, suppose that R_H is a gr-primitive extension of R . Let M be a homogeneous maximal ideal of R and u be a nonzero homogeneous element of R_H . Then there exists a polynomial f in $R[X]$ such that $f(u) = 0$ and $\mathcal{A}_f = R$. Since M is homogeneous, one has $f \notin M[X]$. It follows from [19, Lemma in Page 19], that u or u^{-1} is in R_M . Thus u or u^{-1} is in $R_{H \setminus M}$. Consequently $R_{H \setminus M}$ is a gr-valuation domain and hence R is a graded-Prüfer domain by Proposition 2.1. \square

Let $N := \{f \in R[X] \mid f \neq 0 \text{ and } \mathcal{A}_f = R\}$; then N is a multiplicatively closed subset of $R[X]$, and set $\text{NA}(R) := R[X]_N$. It is known that $N = R[X] \setminus \bigcup \{P[X] \mid P \in h\text{-Max}(R)\}$ and $\text{Max}(\text{NA}(R)) = \{P\text{NA}(R) \mid P \in h\text{-Max}(R)\}$ [17, Proposition 2.3].

The integral closure of R is denoted by \bar{R} . Then \bar{R} is a homogeneous overring of R (cf. [13, Theorem 2.10]).

Remark 2.3. *It is shown in the proof of part (2) \Rightarrow (7) of [11, Theorem 2.9] that $\bar{R}[X]_N = \text{NA}(\bar{R})$, where $N = R[X] \setminus \bigcup \{M[X] \mid M \in h\text{-Max}(R)\}$.*

Lemma 2.4. *Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain. If R_H is a gr-primitive extension of \bar{R} , then R_H is a gr-primitive extension of R .*

Proof. Let $u \in R_H$, $Q' = \{f \in \bar{R}[X] \mid f(u) = 0\}$, and set $Q = Q' \cap R[X]$. If $N = R[X] \setminus \bigcup \{M[X] \mid M \in h\text{-Max}(R)\}$ and $N' = \bar{R}[X] \setminus \bigcup \{M'[X] \mid M' \in h\text{-Max}(\bar{R})\}$, then $\text{NA}(R) = R[X]_N$ and $\text{NA}(\bar{R}) = \bar{R}[X]_{N'}$. The hypothesis that

R_H is a gr-primitive extension of \bar{R} implies that $Q' \cap N' \neq \emptyset$. It suffices to show that $Q \cap N \neq \emptyset$. We first observe that $Q \text{NA}(R) = Q' \bar{R}[X]_N \cap \text{NA}(R)$. That the right side contains the left side is clear, and if $f/n = d/m \in Q' \bar{R}[X]_N \cap \text{NA}(R)$, where $f \in Q'$, $d \in R[X]$, and $n, m \in N$, then $fm = dn \in Q' \cap R[X] = Q$, so that $f/n = fm/nm \in Q \text{NA}(R)$. Thus $Q' \bar{R}[X]_N \cap \text{NA}(R) \subseteq Q \text{NA}(R)$. It follows from Remark 2.3, that $\text{NA}(\bar{R}) = \bar{R}[X]_N$; hence

$$\begin{aligned} Q \text{NA}(R) &= Q' \bar{R}[X]_N \cap \text{NA}(R) = Q' \text{NA}(\bar{R}) \cap \text{NA}(R) \\ &= \text{NA}(\bar{R}) \cap \text{NA}(R) = \text{NA}(R), \end{aligned}$$

which means that $Q \cap N \neq \emptyset$. \square

Gilmer and Hoffmann characterized Prüfer domains as those integrally closed domains D , such that every extension of D inside its quotient field is a primitive extension [10, Theorem 2].

Theorem 2.5. *Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain. Then R is a gr-quasi-Prüfer domain if and only if R_H is a gr-primitive extension of R .*

Proof. Suppose that R_H is a gr-primitive extension of R . Then R_H is a gr-primitive extension of \bar{R} . Therefore by Proposition 2.2, \bar{R} is a graded-Prüfer domain and hence R is a gr-quasi-Prüfer domain by [11, Corollary 2.10]. Conversely, if R is a gr-quasi-Prüfer domain, then \bar{R} is a graded-Prüfer domain ([11, Corollary 2.10]). Then by Proposition 2.2, R_H is a gr-primitive extension of \bar{R} and hence, by Lemma 2.4, R_H is a gr-primitive extension of R . \square

The following is the main result of this section.

Theorem 2.6. *Let \star be a homogeneous preserving semistar operation on a graded integral domain $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ such that $R^\star \subsetneq R_H$. Then the following statements are equivalent:*

- (1) $R \subseteq R_H$ is a gr- \star_f -primitive extension.
- (2) $R_{H \setminus P} \subseteq R_H$ is a gr-primitive extension for each $P \in h\text{-QSpec}^{\star f}(R)$.
- (3) R_P is a quasi-Prüfer domain, for each $P \in h\text{-QSpec}^{\star f}(R)$.
- (4) $R_P \subseteq \text{qf}(R)$ is a primitive extension, for each $P \in h\text{-QSpec}^{\star f}(R)$.
- (5) R is a gr- \star_f -quasi-Prüfer domain.

Proof. (1) \Rightarrow (2) Let $P \in h\text{-QSpec}^{\star f}(R)$ and let u be a nonzero homogeneous element of R_H . Then by assumption there is a polynomial $0 \neq g \in R[X]$ such that $\mathcal{A}_g^{\star f} = R^\star$ and $g(u) = 0$. Clearly, $g \in R_{H \setminus P}[X]$ and $\mathcal{A}_g^R \not\subseteq P$. So

$$\mathcal{A}_g^{R_{H \setminus P}} = \mathcal{A}_g^R R_{H \setminus P} = R_{H \setminus P},$$

and u is primitive over $R_{H \setminus P}$.

(2) \Rightarrow (1) Let u be a nonzero homogeneous element of R_H and let I be the nonzero ideal of $R[X]$ generated by polynomials $f \in R[X]$ such that $f(u) = 0$. We show that $\mathcal{A}_I^{\star f} = R^\star$. Since $R_{H \setminus P} \subseteq R_H$ is a gr-primitive extension for each $P \in h\text{-QMax}^{\star f}(R)$, there is a nonzero polynomial $g \in R_{H \setminus P}[X]$, such

that $g(u) = 0$ and $\mathcal{A}_g^{R_{H \setminus P}} = R_{H \setminus P}$. Let $0 \neq s \in H \setminus P$ with $sg \in R[X]$. Then $\mathcal{A}_{sg}^R \not\subseteq P$ (otherwise, $R_{H \setminus P} = sR_{H \setminus P} = s\mathcal{A}_g^{R_{H \setminus P}} = \mathcal{A}_{sg}^{R_{H \setminus P}} = \mathcal{A}_{sg}^R R_{H \setminus P} \subseteq PR_{H \setminus P}$, a contradiction). Clearly, $sg \in I$ and so $\mathcal{A}_I \not\subseteq P$ for each $P \in h\text{-QMax}^{\star f}(R)$, therefore $\mathcal{A}_I^{\star f} = R^*$. Hence we can find $f \in I$ such that $\mathcal{A}_f^{\star f} = R^*$ and $f(u) = 0$. So u is $\text{gr-}\star f\text{-primitive}$ over R .

(2) \Leftrightarrow (3) For each $P \in h\text{-QSpec}^{\star f}(R)$, $R_{H \setminus P} \subseteq R_H$ is a gr-primitive extension if and only if $R_{H \setminus P}$ is a gr-quasi-Prüfer domain by Theorem 2.5, if and only if R_P is a quasi-Prüfer domain by [11, Theorem 2.9].

(5) \Leftrightarrow (3) Follows from [11, Theorem 2.9].

(3) \Leftrightarrow (4) Follows from [4, Theorem 1.1]. \square

Corollary 2.7. *Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain. Then the following statements are equivalent:*

- (1) $R \subseteq R_H$ is a gr-primitive extension.
- (2) $R_{H \setminus P} \subseteq R_H$ is a gr-primitive extension for each $P \in h\text{-Spec}(R)$.
- (3) $R_P \subseteq \text{qf}(R)$ is a primitive extension, for each $P \in h\text{-Spec}(R)$.
- (4) R is a gr-quasi-Prüfer domain.

Proof. Set $\star = d$ in Theorem 2.6. \square

Corollary 2.8. *Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain. Then the following statements are equivalent:*

- (1) $R \subseteq R_H$ is a $\text{gr-}t\text{-primitive}$ extension.
- (2) $R_{H \setminus P} \subseteq R_H$ is a gr-primitive extension for each $P \in h\text{-Spec}^t(R)$.
- (3) $R_P \subseteq \text{qf}(R)$ is a primitive extension, for each $P \in h\text{-Spec}^t(R)$.
- (4) R is a UMt-domain .

Proof. Follows from Theorem 2.6 by setting $\star = v$ and [11, Theorem 3.2]. \square

3. Graded INC-extension

Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain with quotient field $\text{qf}(R)$, H be the set of nonzero homogeneous elements of R , and \star be a semistar operation on R such that $R^* \subsetneq R_H$. In this section we give a characterization of $\text{gr-}\star\text{-quasi-Prüfer}$ domains in terms of graded semistar INC-extensions.

Assume that $R \subseteq T$ is an extension of graded integral domains. We say that T is a $\text{gr-}\star\text{-INC-extension}$ of R if whenever Q_1 and Q_2 are nonzero homogeneous prime ideals of T such that $Q_1 \cap R = Q_2 \cap R$ and $(Q_1 \cap R)^* \subsetneq R^*$ then Q_1 and Q_2 are incomparable. We also say that R is a $\text{gr-}\star\text{-INC-domain}$ if each homogeneous overring of R is a $\text{gr-}\star\text{-INC-extension}$ of R . We call R a gr-INC-domain if it is a $\text{gr-}d_R\text{-INC-domain}$. It is clear that if R has trivial grading, then gr-INC extension coincides with the usual INC extension of Dobbs [6].

For an ideal I of $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ let I_h denote the ideal of R generated by the set of homogeneous elements of R in I .

Lemma 3.1. *Let $R \subseteq T \subseteq S \subseteq R_H$ be such that T and S are homogeneous overrings of a graded integral domain $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$.*

- (1) *If $R \subseteq T$ and $T \subseteq S$ are gr-INC-extensions, then $R \subseteq S$ is a gr-INC-extension.*
- (2) *If $R \subseteq S$ is a gr-INC-extension and $T \subseteq S$ is an integral extension, then $R \subseteq T$ is a gr-INC-extension.*
- (3) *If $R \subseteq S$ is a gr-INC-extension, then $T \subseteq S$ is a gr-INC-extension.*

Proof. (1) Assume that Q_1 and Q_2 are homogeneous prime ideals of S such that $Q_1 \subsetneq Q_2$. Since $T \subseteq S$ is a gr-INC-extension, then $Q_1 \cap T \subsetneq Q_2 \cap T$. Note that $Q_1 \cap T$ and $Q_2 \cap T$ are homogeneous prime ideals of T . Since $R \subseteq T$ is a gr-INC-extension, then $Q_1 \cap T \cap R \subsetneq Q_2 \cap T \cap R$, i.e., $Q_1 \cap R \subsetneq Q_2 \cap R$.

(2) Assume that P_1 and P_2 are homogeneous prime ideals of T such that $P_1 \subsetneq P_2$. Since $T \subseteq S$ is an integral extension, there are prime ideals $Q_1 \subsetneq Q_2$ of S such that $P_i = Q_i \cap T$ for $i = 1, 2$ ([9, Corollary 11.6]). Since P_i is homogeneous,

$$P_i = (P_i)_h = (Q_i \cap T)_h = (Q_i)_h \cap T$$

for $i = 1, 2$ ([11, Lemma 2.7]). Hence we may assume that Q_i is homogeneous for $i = 1, 2$. Since $R \subseteq S$ is a gr-INC-extension, then $Q_1 \cap R \subsetneq Q_2 \cap R$ and so

$$P_1 \cap R = Q_1 \cap T \cap R = Q_1 \cap R \subsetneq Q_2 \cap R = Q_2 \cap T \cap R = P_2 \cap R.$$

(3) Assume that Q_1 and Q_2 are homogeneous prime ideals of S such that $Q_1 \subsetneq Q_2$. Since $R \subseteq S$ is a gr-INC-extension, we have $Q_1 \cap R \subsetneq Q_2 \cap R$. Now if $Q_1 \cap T = Q_2 \cap T$, then $Q_1 \cap R = Q_2 \cap R$ a contradiction. \square

Recall that Ayache and Jaballah introduced the residually algebraic extension of integral domains and characterized quasi-Prüfer domains as those domains D , such that every extension of D inside its quotient field is a residually algebraic extension [3, Corollary 2.8].

The extension $R \subseteq T$ of graded integral domains is called a *gr-residually algebraic extension*, if for each homogeneous prime ideal Q of T , T/Q is algebraic over $R/(Q \cap R)$, equivalently $qf(R/(Q \cap R)) \hookrightarrow qf(T/Q)$ is an algebraic extension. We say that R is a *gr-residually algebraic domain* if each homogeneous overring of R is a gr-residually algebraic extension of R . Recall that if $I = \bigoplus_{\alpha \in \Gamma} I_\alpha$ is a homogeneous ideal of a graded ring $T = \bigoplus_{\alpha \in \Gamma} T_\alpha$, then T/I is a Γ -graded ring such that for each $\alpha \in \Gamma$, $(T/I)_\alpha = (T_\alpha + I)/I \cong T_\alpha/(T_\alpha \cap I) = T_\alpha/I_\alpha$.

The following lemma is the graded version of [3, Theorem 2.3].

Lemma 3.2. *Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain. Then R is a gr-residually algebraic domain if and only if it is a gr-INC-domain.*

Proof. Assume that R is a gr-residually algebraic domain which is not a gr-INC-domain. So there exists a homogeneous overring T of R such that $R \subseteq T$ is not a gr-INC-extension. Let Q_1 and Q_2 be homogeneous prime ideals of T such that $P := Q_1 \cap R = Q_2 \cap R$ and $Q_1 \subsetneq Q_2$. Moreover, $R/P \subseteq T/Q_1$ is

an algebraic extension of domains, although Q_2/Q_1 is a nonzero ideal of T/Q_1 that intersects R/P in 0, the desired contradiction.

Conversely, assume that R is a gr-INC-domain which is not a gr-residually algebraic domain. So there exists a homogeneous overring T of R such that $R \subseteq T$ is not a residually algebraic extension. Let Q be a homogeneous prime ideal of T such that $R/(Q \cap R) \hookrightarrow T/Q$ is not an algebraic extension. Therefore there exists $u \in T$ such that $u + Q$ is a transcendental element of T/Q over R/P , where $P := Q \cap R$. Let $u + Q = \sum_{i=1}^n (u_i + Q)$ be the decomposition of $u + Q$ to homogeneous components. Hence there is a $u_j + Q$ which is a homogeneous transcendental element of T/Q (in fact if for each i , $u_i + Q$ is algebraic over R/P , then $u + Q$ is algebraic over R/P). Note that u_j is a homogeneous element of T and hence $R[u_j](= R_0[H \cup \{u_j\}])$ is a homogeneous overring of R . Set $x = u_j + Q$ and $T_1 = R[u_j] + Q$. Then T_1 is a homogeneous overring of R and $T_1/Q \cong (R/P)[x]$. Let \wp be the prime ideal of $(R/P)[x]$ generated by x . Then \wp contracts to the zero ideal in R/P , and if Q_1 is a homogeneous prime of T_1 minimal over $u_j T_1 + Q$, then $Q \subsetneq Q_1$ and $\wp = Q_1/Q$. It is easy to check that $Q_1 \cap R = Q \cap R = P$. This contradicts the graded incomparability of the extension $R \subseteq T_1$. \square

Lemma 3.3 (cf. [5, Lemma 10]). *Let $\{P_\lambda\}$ be a chain of homogeneous prime ideals of a graded domain $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$. Then there is a gr-valuation overring of R with a chain of homogeneous prime ideals that contract to $\{P_\lambda\}$.*

Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain and T be a homogeneous overring of R . We say that T is an *h-flat* overring of R if for each homogeneous prime ideal Q of T , one has $R_{Q \cap R} = T_Q$.

Lemma 3.4. *Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain. Then R is a graded-Prüfer domain if and only if each homogeneous overring of R is h-flat.*

Proof. Suppose that R is a graded-Prüfer domain and T is a homogeneous overring of R and let $Q \in h\text{-Spec}(T)$. Therefore $R_{Q \cap R} \subseteq T_Q$ is an inclusion of valuation domains. By [9, Theorem 17.6], $QT_Q \subseteq (Q \cap R)R_{Q \cap R} \subseteq QT_Q$, so $R_{Q \cap R} = T_Q$.

Conversely, assume that each homogeneous overring of R is h-flat. Let P be a homogeneous prime ideal of R . By Lemma 3.3, there exists a gr-valuation overring (V, M) of R such that $P = M \cap R$. Hence $V_M = R_{M \cap R} = R_P$ is a valuation domain. Thus R is a graded-Prüfer domain by Proposition 2.1. \square

Corollary 3.5. *If R is a graded-Prüfer domain, then it is a gr-INC-domain.*

Proof. Let T be a homogeneous overring of R and Q_1 and Q_2 be homogeneous prime ideals of T such that $Q_1 \subsetneq Q_2$. By Lemma 3.4, the map $\phi : h\text{-Spec}(T) \rightarrow h\text{-Spec}(R)$ which sends Q to $Q \cap R$ is injective. Therefore $Q_1 \cap R \subsetneq Q_2 \cap R$. \square

Lemma 3.6. *Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain. Then R is a gr-residually algebraic domain if and only if \bar{R} is a gr-residually algebraic domain.*

Proof. Since one implication is trivial, we will concentrate on the other. Let T be a homogeneous overring of R . Since $\bar{R} \subseteq \bar{T} \subseteq R_H$, then $\bar{R} \subseteq \bar{T}$ is a gr-residually algebraic extension by assumption. Moreover, $R \subseteq \bar{R}$ is a gr-residually algebraic extension, and thus $R \subseteq \bar{T}$ is also a gr-residually algebraic extension by Lemma 3.1(1), and Lemma 3.2. Hence $R \subseteq T$ is a gr-residually algebraic extension by Lemma 3.1(2), and Lemma 3.2. \square

Proposition 3.7. *Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be an integrally closed graded domain. If R is a gr-INC-domain, then $R \subseteq R_H$ is a gr-primitive extension.*

Proof. Assume that $R \subseteq R_H$ is not a gr-primitive extension. So R is not a graded-Prüfer domain by Proposition 2.2. Hence there exists $P \in \text{h-Max}(R)$ such that R_P is not a valuation domain by Proposition 2.1. Thus there is a homogeneous element $u \in R_H$ such that $u \notin R_P$ and $u^{-1} \notin R_P$ by [16, Lemma 4.3]. Consequently by [18, Theorem 7], $PR[u]$ is a (non-maximal) prime ideal of $R[u]$, $P = PR[u] \cap R$ and $R[u]/PR[u] \cong (R/P)[X]$. Since u is a homogeneous element of R_H , $R[u]$ is a homogeneous overring of R . Thus $R \subseteq R[u]$ is a gr-residually algebraic extension by Lemma 3.2. Hence

$$R/P \hookrightarrow R[u]/PR[u] \cong (R/P)[X]$$

is an algebraic extension, a contradiction. \square

The following theorem is the graded version of [6, Theorem].

Theorem 3.8. *Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain. Then R is a gr-INC-domain if and only if $R \subseteq R_H$ is a gr-primitive extension.*

Proof. Assume that R is a gr-INC-domain. Let T be a homogeneous overring of \bar{R} . Thus $R \subseteq T$ is a gr-INC-extension which implies that $\bar{R} \subseteq T$ is a gr-INC-extension by Lemma 3.1(3). Therefore \bar{R} is an integrally closed gr-INC-domain, hence $\bar{R} \subseteq R_H$ is a gr-primitive extension by Proposition 3.7. Now $R \subseteq R_H$ is a gr-primitive extension by Lemma 2.4.

Conversely, assume that $R \subseteq R_H$ is a gr-primitive extension. Then by the proof of Theorem 2.5, \bar{R} is a graded-Prüfer domain. So \bar{R} is a gr-INC-domain by Corollary 3.5. Thus \bar{R} is a gr-residually algebraic domain by Lemma 3.2. Therefore R is a gr-residually algebraic domain by Lemma 3.6, and hence R is a gr-INC-domain by Lemma 3.2. \square

Corollary 3.9. *Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain. Then the following statements are equivalent:*

- (1) R is a gr-quasi-Prüfer domain.
- (2) R is a gr-residually algebraic domain.

Proof. It follows by combining Corollary 2.7, Theorem 3.8 and Lemma 3.2. \square

The following is the main result of this section.

Theorem 3.10. *Let \star be a homogeneous preserving semistar operation on a graded integral domain $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ such that $R^\star \subsetneq R_H$. Then the following statements are equivalent:*

- (1) R is a $gr\text{-}\star_f\text{-INC}$ -domain.
- (2) $R_{H \setminus P}$ is a $gr\text{-INC}$ -domain, for each $P \in h\text{-QSpec}^{\star_f}(R)$.
- (3) R_P is a quasi-Prüfer domain, for each $P \in h\text{-QSpec}^{\star_f}(R)$.
- (4) R_P is an INC -domain, for each $P \in h\text{-QSpec}^{\star_f}(R)$.
- (5) R is a $gr\text{-}\star_f\text{-quasi-Prüfer}$ domain.

Proof. (1) \Rightarrow (2) Let $P \in h\text{-QSpec}^{\star_f}(R)$ and let T be a homogeneous overring of $R_{H \setminus P}$. Assume that Q_1 and Q_2 are homogeneous prime ideals of T such that $Q_1 \cap R_{H \setminus P} = Q_2 \cap R_{H \setminus P}$. We must show that Q_1 and Q_2 are incomparable. By assumption, T is a $gr\text{-}\star_f\text{-INC}$ -extension of R and $Q_1 \cap R = Q_2 \cap R \subseteq P$ with $P^{\star_f} \subsetneq R^\star$, hence Q_1 and Q_2 are incomparable.

(2) \Rightarrow (1) Let T be a homogeneous overring of R and $Q_1 \subsetneq Q_2$ be homogeneous prime ideals of T such that $P := Q_1 \cap R = Q_2 \cap R \subseteq M$ for some $M \in h\text{-QSpec}^{\star_f}(R)$. Note that $Q_1 T_{H \setminus M} \subsetneq Q_2 T_{H \setminus M}$ and they are prime ideals of $T_{H \setminus M}$ each of which intersects $R_{H \setminus M}$ in $PR_{H \setminus M}$, and so $R_{H \setminus M} \subseteq T_{H \setminus M}$ is not a $gr\text{-INC}$ -extension, contradicting (2).

(2) \Leftrightarrow (3) $R_{H \setminus P}$ is a $gr\text{-INC}$ -domain for each $P \in h\text{-QSpec}^{\star_f}(R)$ if and only if $R_{H \setminus P} \subseteq R_H$ is a $gr\text{-primitive}$ extension by Theorem 3.8, if and only if R_P is a quasi-Prüfer domain, for each $P \in h\text{-QSpec}^{\star_f}(R)$ by Theorem 2.6.

(3) \Leftrightarrow (4) and (3) \Leftrightarrow (5) Follow from [4, Theorem 1.1] and [11, Theorem 2.9], respectively. \square

Corollary 3.11. *Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain. Then the following statements are equivalent:*

- (1) R is a $gr\text{-INC}$ -domain.
- (2) $R_{H \setminus P}$ is a $gr\text{-INC}$ -domain, for each $P \in h\text{-Spec}(R)$.
- (3) R_P is an INC -domain, for each $P \in h\text{-Spec}(R)$.
- (4) R is a $gr\text{-quasi-Prüfer}$ domain.

Proof. Set $\star = d$ in Theorem 3.10. \square

Corollary 3.12. *Let $R = \bigoplus_{\alpha \in \Gamma} R_\alpha$ be a graded integral domain. Then the following statements are equivalent:*

- (1) R is a $gr\text{-}t\text{-INC}$ -domain.
- (2) $R_{H \setminus P}$ is a $gr\text{-INC}$ -domain, for each $P \in h\text{-Spec}^t(R)$.
- (3) R_P is an INC -domain, for each $P \in h\text{-Spec}^t(R)$.
- (4) R is a UMt -domain.

Proof. Follows from Theorem 3.10 by setting $\star = v$ and [11, Theorem 3.2]. \square

Acknowledgment. We wish to thank the referee for careful reading and an insightful report. We also would like to thank Dr. Nematollah Shirmohammadi for helpful discussion.

References

- [1] D. D. Anderson and D. F. Anderson, *Divisibility properties of graded domains*, Canad. J. Math. **34** (1982), no. 1, 196–215.
- [2] D. F. Anderson and G. W. Chang, *Graded integral domains and Nagata rings*, J. Algebra **387** (2013), 169–184.
- [3] A. Ayache and A. Jaballah, *Residually algebraic pairs of rings*, Math. Z. **225** (1997), no. 1, 49–65.
- [4] G. W. Chang and M. Fontana, *Uppers to zero in polynomial rings and Prüfer-like domains*, Comm. Algebra **37** (2009), no. 1, 164–192.
- [5] G. W. Chang and D. Y. Oh, *Discrete valuation overrings of a graded Noetherian domain*, J. Commut. Algebra to appear.
- [6] D. E. Dobbs, *On INC-extensions and polynomials with unit content*, Canad. Math. Bull. **23** (1980), no. 1, 37–42.
- [7] M. Fontana and J. A. Huckaba, *Localizing systems and semistar operations*, in Non-Noetherian commutative ring theory, 169–197, Math. Appl., 520, Kluwer Acad. Publ., Dordrecht.
- [8] M. Fontana, J. A. Huckaba, and I. J. Papick, *Prüfer domains*, Monographs and Textbooks in Pure and Applied Mathematics, 203, Marcel Dekker, Inc., New York, 1997.
- [9] R. Gilmer, *Multiplicative Ideal Theory*, Marcel Dekker, Inc., New York, 1972.
- [10] R. Gilmer and J. F. Hoffmann, *A characterization of Prüfer domains in terms of polynomials*, Pacific J. Math. **60** (1975), no. 1, 81–85.
- [11] H. Hamdi and P. Sahandi, *Uppers to zero in polynomial rings over graded domains and UMt-domains*, Bull. Korean Math. Soc. **55** (2018), no. 1, 187–204.
- [12] E. Houston and M. Zafrullah, *On t -invertibility. II*, Comm. Algebra **17** (1989), no. 8, 1955–1969.
- [13] J. L. Johnson, *Integral closure and generalized transforms in graded domains*, Pacific J. Math. **107** (1983), no. 1, 173–178.
- [14] D. G. Northcott, *Lessons on rings, modules and multiplicities*, Cambridge University Press, London, 1968.
- [15] A. Okabe and R. Matsuda, *Semistar-operations on integral domains*, Math. J. Toyama Univ. **17** (1994), 1–21.
- [16] P. Sahandi, *Characterizations of graded Prüfer \star -multiplication domains*, Korean J. Math. **22**, (2014), 181–206.
- [17] ———, *Characterizations of graded Prüfer \star -multiplication domains, II*, Bull. Iranian Math. Soc. to appear, DOI:10.1007/s1980-018-0005-1.
- [18] A. Seidenberg, *A note on the dimension theory of rings*, Pacific J. Math. **3** (1953), 505–512.
- [19] O. Zariski and P. Samuel, *Commutative algebra. Vol. II*, The University Series in Higher Mathematics, D. Van Nostrand Co., Inc., Princeton, NJ, 1960.

HALEH HAMDI
DEPARTMENT OF PURE MATHEMATICS
FACULTY OF MATHEMATICAL SCIENCES
UNIVERSITY OF TABRIZ
TABRIZ, IRAN
Email address: h.hamdimoghadam@tabrizu.ac.ir

PARVIZ SAHANDI
DEPARTMENT OF PURE MATHEMATICS
FACULTY OF MATHEMATICAL SCIENCES
UNIVERSITY OF TABRIZ
TABRIZ, IRAN
Email address: sahandi@ipm.ir