

## RATIONAL HOMOTOPY TYPE OF MAPPING SPACES BETWEEN COMPLEX PROJECTIVE SPACES AND THEIR EVALUATION SUBGROUPS

J.-B. GATSINZI

ABSTRACT. We use  $L_\infty$  models to compute the rational homotopy type of the mapping space of the component of the natural inclusion  $i_{n,k} : \mathbb{C}P^n \hookrightarrow \mathbb{C}P^{n+k}$  between complex projective spaces and show that it has the rational homotopy type of a product of odd dimensional spheres and a complex projective space. We also characterize the mapping  $\text{aut}_1 \mathbb{C}P^n \rightarrow \text{map}(\mathbb{C}P^n, \mathbb{C}P^{n+k}; i_{n,k})$  and the resulting  $G$ -sequence.

### 1. Introduction

Let  $f : X \rightarrow Y$  be a map between simply connected CW-complexes of finite type. We denote by  $\text{map}(X, Y; f)$  the path component of  $f$  in the space of continuous mappings from  $X$  to  $Y$ . The study of the rational homotopy type of  $\text{map}(X, Y; f)$  was initiated by Haefliger [10] who describes its Sullivan model. Afterwards there were attempts to find a Quillen model of  $\text{map}(X, Y; f)$  from either a Sullivan or a Quillen model of  $f$ . Chain complexes of which the homology coincides with rational homotopy groups of function spaces were investigated [8, 12, 13]. Those chain complexes were later developed into models of function spaces [2–5].

Following [5] we describe in this paper an  $L_\infty$  model of the inclusion  $i_{n,k} : \mathbb{C}P^n \hookrightarrow \mathbb{C}P^{n+k}$ . We shall use rational homotopy theory for which the standard reference is [6].

The notion of  $L_\infty$ -algebra was introduced by Lada [11] and we remind here the definition.

**Definition 1.** A permutation  $\sigma \in S_n$  is called an  $(i, n - i)$ -shuffle if  $\sigma(1) < \dots < \sigma(i)$  and  $\sigma(i + 1) < \dots < \sigma(n)$ , where  $i = 1, \dots, n$ . For graded objects  $x_1, \dots, x_n$ , the Koszul sign  $\epsilon(\sigma)$  is determined by

$$x_1 \wedge \dots \wedge x_n = \epsilon(\sigma) x_{\sigma(1)} \wedge \dots \wedge x_{\sigma(n)}.$$

It depends not only of the permutation  $\sigma$  but also of degrees of  $x_1, \dots, x_n$ .

---

Received November 16, 2020; Accepted January 20, 2021.

2010 *Mathematics Subject Classification.* Primary 55P62; Secondary 54C35.

*Key words and phrases.* Rational homotopy theory, mapping space,  $L_\infty$  algebra.

We assume that all vector spaces are over the field of rational numbers  $\mathbb{Q}$ .

**Definition 2.** An  $L_\infty$ -algebra or a strongly homotopy Lie algebra is a graded vector space  $L = \bigoplus_{i \geq 0} L_i$  with maps  $\ell_k : L^{\otimes k} \rightarrow L$  of degree  $k - 2$  such that

- (1)  $\ell_k$  is graded skew symmetric, that is, for a  $k$ -permutation  $\sigma$

$$\ell_k(x_{\sigma(1)}, \dots, x_{\sigma(k)}) = \text{sgn}(\sigma) \epsilon(\sigma) \ell_k(x_1, \dots, x_k),$$

where  $\text{sgn}(\sigma)$  is the sign of  $\sigma$ .

- (2) There are some generalized Jacobi identities

$$\sum_{i+j=n+1} \sum_{\sigma} \epsilon(\sigma) (-1)^{i(j-1)} \ell_j(\ell_i(x_{\sigma(1)}, \dots, x_{\sigma(i)}), x_{\sigma(i+1)}, \dots, x_{\sigma(n)}) = 0,$$

where the summation extends to all  $(i, n - i)$  shuffles of the symmetric group  $S_n$ .

If  $\ell_k = 0$  for  $k \geq 3$ , one retrieves the definition of a graded differential Lie algebra  $(L, d)$  where  $d = \ell_1$  and  $\ell_2$  is the Lie bracket.

Let  $(L, \ell_k)$  be an  $L_\infty$  algebra and  $sL$  the suspension of  $L$ , and  $C_\infty(L) = (\wedge sL, d)$  the generalized Cartan-Chevalley-Eilenberg functor (see [6, §22]). One gets linear mappings  $d_k : \wedge^k(sL) \rightarrow sL$  defined by

$$d_k(sx_1 \wedge \dots \wedge sx_k) = (-1)^{\frac{k(k-1)}{2}} \ell_k(x_1, \dots, x_k),$$

each of which extends into a codifferential on the coalgebra  $\wedge sL$ . This gives an equivalence between  $L_\infty$  structures on  $L$  and codifferentials on  $\wedge sL$  [11]. Moreover if  $L$  is of finite type, then  $C^\infty(L) = (\wedge(sL)^\#, d)$  is a commutative differential graded algebra (cdga for short). The differential  $d = d_1 + \dots + d_k + \dots$  is defined by

$$\langle d_k v, sx_1 \wedge \dots \wedge sx_k \rangle = (-1)^\epsilon \langle v, \ell_k(x_1, \dots, x_k) \rangle,$$

where  $v \in (sL)^\#$  and  $\epsilon = \sum_{i=1}^{k-1} (k-i)|x_i|$ .

**Definition 3.** Two cdga's  $(A, d)$  and  $(B, d)$  have the same homotopy type if they are linked by a sequence of quasi-isomorphisms

$$(A, d) = A_0 \rightarrow A_1 \leftarrow A_2 \cdots \rightarrow A_{n-1} \leftarrow A_n = (B, d).$$

Let  $V$  be a graded vector space. A Sullivan algebra  $(\wedge V, d)$  is the free graded commutative algebra generated by  $V$  together with a filtration  $V(0) \subset V(1) \subset \dots \subset V$  such that  $dV(i) \subset \wedge V(i-1)$ . It is called minimal if  $dV \subset \wedge^{\geq 2} V$ . A Sullivan model of a simply connected space  $X$  is a Sullivan algebra  $(\wedge V, d)$  such that there exists a quasi-isomorphism  $\varphi : (\wedge V, d) \rightarrow A_{PL}(X)$ , where  $A_{PL}(X)$  denotes the cdga of piecewise linear forms of  $X$  [16]. A cdga model of  $X$  is a cdga  $(A, d)$  which has the same homotopy type as  $A_{PL}(X)$ .

**Definition 4.** If  $f : X \rightarrow Y$  is a map between simply connected spaces of finite type, then there is a cdga map  $\phi : (\wedge V, d) \rightarrow (B, d)$ , called a model of  $f$ , where  $(B, d)$  and  $(\wedge V, d)$  are respective cdga models of  $Y$  and  $X$ , respectively.

**Definition 5.** Let  $L$  be an  $L_\infty$ -algebra of finite type. Then  $L$  is called an  $L_\infty$  model of a topological space  $X$  if  $C^\infty(L)$  is a Sullivan model of  $X$ . It is minimal if  $\ell_1 = 0$ . In this case  $\pi_*(\Omega X) \otimes \mathbb{Q} \cong L$ .

In this note, we give another proof of the following result using  $L_\infty$  models of function spaces (see [15], Example 3.4).

**Theorem 6.** *The function space  $\text{map}(\mathbb{C}P^n, \mathbb{C}P^{n+k}; i_{n,k})$  has the rational homotopy type of  $\mathbb{C}P^k \times S^{2k+3} \times \dots \times S^{2(n+k)+1}$ .*

Moreover we study evaluation subgroups of the mapping  $\text{aut}_1 \mathbb{C}P^n \rightarrow \mathbb{C}P^{n+k}$  and prove the following result.

**Theorem 7.** *The  $G$ -sequence associated with the inclusion*

$$\text{aut}_1 \mathbb{C}P^n \rightarrow \text{map}(\mathbb{C}P^n, \mathbb{C}P^{n+k}; i_{n,k})$$

*is not exact.*

## 2. $L_\infty$ -models of function spaces

**Definition 8.** Let  $\phi : (\wedge V, d) \rightarrow (B, d)$  be a morphism of cdga's. A  $\phi$ -derivation of degree  $k$  is a linear mapping  $\theta : (\wedge V)^n \rightarrow B^{n-k}$  such that  $\theta(ab) = \theta(a)\phi(b) + (-1)^{k|a|}\phi(a)\theta(b)$ . We denote by  $\text{Der}(\wedge V, B; \phi)$  the  $\mathbb{Z}$ -graded vector space of all  $\phi$ -derivations. The differential on  $\text{Der}(\wedge V, B; \phi)$  is defined by  $\delta\theta = d\theta - (-1)^k\theta d$ .

Define  $\widetilde{\text{Der}}(\wedge V, B; \phi)$  as

$$\widetilde{\text{Der}}(\wedge V, B, \phi)_i = \begin{cases} \text{Der}(\wedge V, B; \phi)_i, & i > 1, \\ \{\theta \in \text{Der}(\wedge V, B; \phi)_1 : \delta\theta = 0\}, & i = 1. \end{cases}$$

If  $\varphi_1, \dots, \varphi_k \in \widetilde{\text{Der}}(\wedge V, B; \phi)$  are  $\phi$ -derivations of respective degrees  $n_1, \dots, n_k$ , define

$$\begin{aligned} & [\varphi_1, \dots, \varphi_k](v) \\ &= (-1)^{n_1 + \dots + n_k - 1} \sum_{i_1, \dots, i_k} \epsilon \phi(v_1 \cdots \hat{v}_{i_1} \cdots \hat{v}_{i_k} \cdots v_m) \varphi_1(v_{i_1}) \cdots \varphi_k(v_{i_k}), \end{aligned}$$

where  $dv = \sum v_1 \cdots v_m$  and  $\epsilon$  is the corresponding Koszul sign of the permutation

$$(\varphi_1, \dots, \varphi_k, v_1, \dots, v_m) \rightarrow (v_1, \dots, \hat{v}_{i_1}, \dots, \hat{v}_{i_k}, \dots, v_m, \varphi_1, v_{i_1}, \dots, \varphi_k, v_{i_k}).$$

We note that  $[\varphi_1, \dots, \varphi_k]$  is of degree  $n_1 + \dots + n_k - 1$ . Now define linear maps  $\ell_k$  of degree  $k - 2$  on  $s^{-1}\widetilde{\text{Der}}(\wedge V, B, \phi)$  by

$$\ell_1(s^{-1}\varphi) = -s^{-1}\delta\varphi, \quad \ell_k(s^{-1}\varphi_1, \dots, s^{-1}\varphi_k) = (-1)^{\epsilon_k} s^{-1}[\varphi_1, \dots, \varphi_k],$$

where  $\epsilon_k = \sum_{i=1}^{k-1} (k-i)|\varphi_i|$ .

**Proposition 9** (Lemma 3.3, [5]). *If  $\phi : (\wedge V, d) \rightarrow (B, d)$  is a Sullivan model of a mapping  $f : X \rightarrow Y$  between simply connected spaces and  $V$  is finite dimensional, then  $(s^{-1}\widetilde{\text{Der}}(\wedge V, B; \phi), \ell_k)$  is an  $L_\infty$  model of  $\text{map}(X, Y; f)$ .*

### 3. Component of the inclusion $\mathbb{C}P^n \rightarrow \mathbb{C}P^{n+k}$

Recall that the minimal Sullivan model of  $\mathbb{C}P^n$  is given by  $(\wedge(x_2, x_{2n+1}), d)$  where  $dx_2 = 0$ ,  $dx_{2n+1} = x_2^{n+1}$ . Our objective is to compute an  $L_\infty$  model of the component of the inclusion  $\mathbb{C}P^n \rightarrow \mathbb{C}P^{n+k}$ . For  $k = 0$ , one gets a model of  $\text{aut}_1 \mathbb{C}P^n = \text{map}(\mathbb{C}P^n, \mathbb{C}P^n; \text{Id})$  from the differential Lie algebra  $(L, \delta)$  of derivations of  $(\wedge(x_2, x_{2n+1}), d)$ , of which  $H_*(L, \delta)$  is spanned by  $\{z_3, z_5, \dots, z_{2n+1}\}$  [7, §3]. Therefore  $\text{aut}_1 \mathbb{C}P^n$  has the rational homotopy type of the product  $S^3 \times S^5 \times \dots \times S^{2n+1}$ . This result was also proved by Møller and Raussen using another method [15, Example 3.4].

Let  $f : (\wedge V, d) \rightarrow (B, d)$  be a morphism of differential graded algebras. For  $v \in V$  and  $b \in B$  we denote by  $(v, b)$  the unique  $f$ -derivation  $\theta$  such that  $\theta(v) = b$  and zero on the remaining generators of  $\wedge V$ .

From now on we assume that  $k \geq 1$ . A model of the inclusion

$$i_{n,k} : \mathbb{C}P^n \rightarrow \mathbb{C}P^{n+k}$$

is given by

$$\psi : (A, d) = (\wedge(x_2, x_{2n+2k+1}) \rightarrow (\wedge(y_2, y_{2n+1}), d) = (B, d),$$

where  $\psi(x_2) = y_2$ ,  $\psi(x_{2n+2k+1}) = y_2^k y_{2n+1}$ . We consider the composition

$$\phi : A = (\wedge(x_2, x_{2n+2k+1}) \xrightarrow{\psi} (\wedge(y_2, y_{2n+1}), d) = B \simeq (\wedge(y_2)/(y_2^{n+1}), 0).$$

Hence  $\phi(x_2) = y_2$  and  $\phi(x_{2n+2k+1}) = 0$ . The induced map

$$(\text{Der}(A, B; \psi), \delta) \rightarrow (\text{Der}(A, H^*(B); \phi), \delta)$$

is a quasi-isomorphism [1]. In the sequel we compute

$$\widetilde{\text{Der}}(\wedge(x_2, x_{2n+2k+1}), \wedge(y_2)/(y_2^{n+1}); \phi)$$

and determine its brackets. As a vector space

$$\widetilde{\text{Der}}(\wedge(x_2, x_{2n+2k+1}), \wedge(y_2)/(y_2^{n+1}); \phi)$$

is spanned by

$$\{\beta_2, \alpha_{2k+2i-1}, i = 1, \dots, n+1\},$$

where  $\alpha_{2k+2i-1} = (x_{2n+2k+1}, y_2^{n-i+1})$  and  $\beta_2 = (x_2, 1)$ . Note that  $|\beta_2| = 2$  and  $|\alpha_{2k+2i-1}| = 2k + 2i - 1$ . Computations show that the only non zero brackets are given by  $\underbrace{[\beta_2, \dots, \beta_2]}_{k+i} = \alpha_{2k+2i-1}$  for  $i = 1, \dots, n+1$ .

We deduce the following result (see [15] for a different proof).

**Proposition 10.** *The function space  $\text{map}(\mathbb{C}P^n, \mathbb{C}P^{n+k}; i_{n,k})$  has a Sullivan model of the form*

$$(\wedge(z_2, z_{2k+1}, \dots, z_{2k+2n+1}), d),$$

where  $dz_2 = 0$ ,  $dz_{2k+1} = z_2^{k+1}$ ,  $\dots$ ,  $dz_{2k+2n+1} = z_2^{k+n+1}$ .

*Proof.* An  $L_\infty$  model  $(L, \ell_k)$  of  $\text{map}(\mathbb{C}P(n), \mathbb{C}P(n+k); i_{n,k})$  is spanned by

$$\langle s^{-1}\beta_2, s^{-1}\alpha_{2k+2i-1}, i = 1, \dots, n+1 \rangle.$$

Moreover  $\ell_j = 0$  for  $j = 1, \dots, k$  and  $\ell_{k+i}(s^{-1}\beta_2, \dots, s^{-1}\beta_2) = s^{-1}\alpha_{2k+2i-1}$ , for  $i = 1, \dots, n+1$ . Therefore

$$C^\infty(L) = \wedge(z_2, z_{2k+1}, z_{2k+3}, \dots, z_{2k+2n+1}), d, \quad dz_2 = 0, dz_{2k+2i+1} = z_2^{k+i+1},$$

where  $0 \leq i \leq n$ .  $\square$

**Theorem 11.** *The function space  $\text{map}(\mathbb{C}P^n, \mathbb{C}P^{n+k}; i_{n,k})$  has the rational homotopy type of  $\mathbb{C}P^k \times S^{2k+3} \times \dots \times S^{2(n+k)+1}$ .*

*Proof.* By the above result, a Sullivan model of  $\text{map}(\mathbb{C}P^n, \mathbb{C}P^{n+k}; i_{n,k})$  is given by

$$(\wedge(x_2, x_{2k+1}, x_{2k+3}, \dots, x_{2n+2k+1}),$$

where  $dx_2 = 0, dx_{2i+1} = x_2^{i+1}, i = k, k+1, \dots, k+n$ . We consider the relative Sullivan model

$$(\wedge(x_2, x_{2k+1}), d) \rightarrow (\wedge(x_2, x_{2k+1}) \otimes \wedge x_{2k+3}, D),$$

where

$$dx_2 = 0, dx_{2k+1} = x_2^{k+1}, Dx_2 = dx_2, Dx_{2k+1} = dx_{2k+1}, Dx_{2k+3} = x_2^{k+2}.$$

It is a Sullivan model of the fibration  $S^{2k+3} \rightarrow E \xrightarrow{p} \mathbb{C}P^k$ , where  $p$  is classified by a map  $f : \mathbb{C}P^k \rightarrow B \text{aut}_1 S^{2k+3}$ . Using the algebra of derivations on the minimal Sullivan model of  $S^{2k+3}$  [16], it is easily seen that  $B \text{aut}_1 S^{2k+3}$  has the rational homotopy type of  $K(\mathbb{Q}, 2k+4)$  [7, Proposition 2.1].

Moreover equivalence classes

$$[\mathbb{C}P^k, K(\mathbb{Q}, 2k+4)]$$

are in a bijective correspondence with  $H^{2k+4}(\mathbb{C}P^k, \mathbb{Q}) = \{0\}$ . Therefore the classifying map  $f$  is rationally trivial. So we deduce that the fibration is trivial. Hence the cdga

$$(A, d) = (\wedge(x_2, x_{2k+1}, x_{2k+3}), d), dx_2 = 0, dx_{2k+1} = x_2^{k+1}, dx_{2k+3} = x_2^{k+2}$$

and

$$(\wedge(x_2, x_{2k+1}) \otimes \wedge z_{2k+3}, d), dx_2 = 0, dx_{2k+1} = x_2^{k+1}, dz_{2k+3} = 0$$

are isomorphic. We deduce that the cdga  $(A, d)$  is a Sullivan model of  $\mathbb{C}P^k \times S^{2k+3}$ . It follows from an induction argument that  $\text{map}(\mathbb{C}P^k, \mathbb{C}P^{n+k}; i_{n,k})$  has the rational homotopy type of  $\mathbb{C}P^k \times S^{2k+3} \times \dots \times S^{2(n+k)+1}$ .  $\square$

Recall that a Sullivan algebra  $(\wedge V, d)$  is called formal if there is a quasi-isomorphism  $(\wedge V, d) \rightarrow H^*(\wedge V, d)$ . Spheres and complex projective spaces are formal. Moreover a product of formal spaces is also formal. We deduce that:

**Corollary 12.** *The function space  $\text{map}(\mathbb{C}P^n, \mathbb{C}P^{n+k}; i_{n,k})$  is formal.*

#### 4. Evaluation subgroups of the inclusion $i_{n,k} : \mathbb{C}P^n \rightarrow \mathbb{C}P^{n+k}$

We consider the inclusion  $i_{n,k} : \mathbb{C}P^n \rightarrow \mathbb{C}P^{n+k}$  and the corresponding Sullivan model  $\phi$  of the previous section given by the composition

$$\phi : A = (\wedge(x_2, x_{2n+2k+1}), d) \xrightarrow{\psi} \wedge(y_2, y_{2n+1}), d = B \xrightarrow[\simeq]{\gamma} H^*(B).$$

Forgetting the desuspension, a model of the inclusion  $(i_{n,k})_* : \text{aut}_1 \mathbb{C}P^n \rightarrow \text{map}(\mathbb{C}P^n, \mathbb{C}P^{n+k}; i_{n,k})$  is given by

$$\phi^* : (\text{Der}(B, H^*(B); \gamma), \delta) \rightarrow (\text{Der}(A, H^*(B); \phi), \delta).$$

We know characterize the map  $\phi^*$  when  $k > n$ .

**Theorem 13.** *If  $k > n$ , then the induced map*

$$\phi^* : (\text{Der}(B, H^*(B); \gamma), \delta) \longrightarrow (\text{Der}(A, H^*(B), \phi), \delta)$$

*is homotopy trivial.*

*Proof.* We note that  $L = \text{Der}(B, H^*(B); \gamma)$  is spanned by

$$\{\delta_2, \theta_1, \theta_3, \dots, \theta_{2n+1}\},$$

where  $\delta_2 = (y_2, 1)$ ,  $\theta_{2i+1} = (y_{2n+1}, y_2^{n-i})$ ,  $i = 0, \dots, n$ . The differential is given by  $\delta\delta_2 = (n+1)\theta_1$  and zero otherwise. Therefore

$$\pi_*(\text{aut}_1 \mathbb{C}P^n) \otimes \mathbb{Q} = H_*(L, \delta) = \langle [\theta_3], \dots, [\theta_{2n+1}] \rangle.$$

Hence  $\text{aut}_1 \mathbb{C}P^n$  has the rational homotopy type of  $S^3 \times S^5 \times \dots \times S^{2n+1}$ . Let

$$L' = (\text{Der}(A, H^*(B), \phi), \delta) = (\langle \beta_2, \alpha_{2k+1}, \dots, \alpha_{2n+2k+1} \rangle, \delta).$$

The mapping  $\phi^* : L \rightarrow L'$  is defined by  $\phi^*(\delta_2) = \beta_2$ ,  $\phi^*(\theta_{2i+1}) = 0$  for  $i < k$ , and  $\phi^*(\theta_{2i+1}) = \alpha_{2i+1}$  for  $i \geq k$ . If  $k > n$ , then  $\phi^*(\delta_2) = \beta_2$  and zero otherwise. Moreover

$$C^\infty(s^{-1}L) = (\wedge(x_2, y_1, \dots, y_{2i-1}, \dots, y_{2n+1}), d),$$

where  $dx_2 = 0$  and  $dy_{2i-1} = x_2^i$ . In particular  $dy_1 = x_2$ . In the same way

$$C^\infty(s^{-1}L') = (\wedge(u_2, v_{2k+1}, \dots, v_{2n+2k+1}), d),$$

where  $du_2 = 0$ ,  $dv_{2i+1} = u_2^{i+1}$ . Hence

$$\Phi = C^\infty(\phi^*) : C^\infty(s^{-1}L') \rightarrow C^\infty(s^{-1}L)$$

is defined by  $\Phi(u_2) = x_2$  and vanishes on other generators. As  $C^\infty(s^{-1}L')$  is quasi-isomorphic to

$$(\wedge(w_2, w_{2k+1}), d) \otimes (\wedge(w_{2k+3}, \dots, w_{2n+2k+1}), 0),$$

where  $dw_2 = 0$ ,  $dw_{2k+1} = w_2^{k+1}$  and,  $C^\infty(s^{-1}L)$  is quasi-isomorphic to

$$(\wedge(z_3, \dots, z_{2n+1}), 0),$$

then induced map

$$\tilde{\Phi} : (\wedge(w_2, w_{2k+1}, w_{2k+3}, \dots, w_{2n+2k+1}), d) \rightarrow (\wedge(z_3, \dots, z_{2n+1}), 0)$$

between minimal Sullivan models is zero.  $\square$

**Definition 14.** Let  $X$  be a topological space. We say  $\alpha \in \pi_n(X)$  is a Gottlieb element if the map:  $f \vee 1_X : S^n \vee X \rightarrow X$  extends to  $S^n \times X$ , where  $f$  represents the homotopy class  $\alpha$  [9].

Gottlieb elements form a subgroup of  $\pi_*(X)$  which will be denoted by  $G_*(X)$ . It comes from the definition that  $G_*(X)$  is the image of  $\pi_*(\text{ev}) : \pi_*(\text{aut}_1 X, 1_X) \rightarrow \pi_*(X, x_0)$ , where  $\text{ev}$  is the evaluation map at  $x_0$ . If  $f : X \rightarrow Y$ , then  $G_*(Y, X; f)$  is the image of  $\pi_*(\text{ev})$  where  $\text{ev} : \text{map}(X, Y; f) \rightarrow Y$  is the evaluation map at the base point.

Let  $(\wedge V, d)$  be the minimal Sullivan model of a simply connected space  $X$ . Define the Gottlieb group of  $(\wedge V, d)$

$$G_n(\wedge V, d) = \{[\theta] \in H_n(\text{Der } \wedge V, \delta) : \theta(v) = 1, v \in V^n\}.$$

Hence  $G_*(\wedge V, d) \cong \text{im } H_*(\epsilon_*)$ , where  $\epsilon_* : \text{Der } \wedge V \rightarrow \text{Der}(\wedge V, \mathbb{Q}; \epsilon)$  is the post composition with the augmentation map  $\epsilon : \wedge V \rightarrow \mathbb{Q}$ . Then  $G_n(\wedge V) \cong G_n(X_{\mathbb{Q}})$ , where  $h : X \rightarrow X_{\mathbb{Q}}$  is the rationalization [6, Propostion 29.8]. There are also relative Gottlieb groups  $G_*^{rel}(Y, X; f)$  and a  $G$ -sequence

$$\cdots \rightarrow G_{n+1}^{rel}(Y, X; f) \rightarrow G_n(X) \rightarrow G_n(Y, X; f) \rightarrow \cdots$$

which was introduced by Lee and Woo. The sequence is exact in some cases, for instance if  $f$  has a left homotopy inverse [17]. We follow the description of rational evaluation homotopy groups as given by Lupton and Smith [12].

Using augmentation maps we obtain the commutative diagram.

$$\begin{array}{ccc} \text{Der}(B, H^*(B); \gamma) & \xrightarrow{\phi^*} & \text{Der}(A, H^*(B); \phi) \\ \epsilon_* \downarrow & & \downarrow \epsilon_* \\ \text{Der}(B, \mathbb{Q}; \epsilon) & \xrightarrow{\widehat{\phi}^*} & \text{Der}(A, \mathbb{Q}; \epsilon) \end{array}$$

In the same way we define  $G_*(A, H^*(B); \phi)$  as the image of  $H_*(\epsilon_*)$  in  $H_*(\text{Der}(A, \mathbb{Q}, \epsilon))$ .

In order to define relative rational Gottlieb groups, we recall that if  $\phi : (C, d_C) \rightarrow (C', d_{C'})$  is a map of chain complexes, the mapping cone of  $\phi$ , denoted by  $\text{Rel}(\phi)$ , is the complex of which the underlying graded vector space is  $sC \oplus C'$  and the differential is given by  $D(sx, y) = (-sd_C(x), \phi(x) + d_{C'}y)$  [12] or [14, p. 46]. Define chain maps  $J : C'_n \rightarrow \text{Rel}_n(\phi)$  and  $P : \text{Rel}_n(\phi) \rightarrow C_{n-1}$  by  $J(y) = (0, y)$  and  $P(sx, y) = x$ . This yields an exact sequence of chain complexes

$$0 \rightarrow C'_* \xrightarrow{J} \text{Rel}_*(\phi) \xrightarrow{P} C_{*-1} \rightarrow 0,$$

which induces a long exact sequence in homology [14, Proposition 4.3]. We consider the mapping cone  $\text{Rel}(\phi^*)$  of

$$\phi^* : (\text{Der}(B, H^*(B), \gamma), \delta) \rightarrow (\text{Der}(A, H^*(B), \phi), \delta),$$

$\text{Rel}(\widehat{\phi}^*)$  the mapping cone of  $\widehat{\phi}^* : \text{Der}(B, \mathbb{Q}; \epsilon) \rightarrow \text{Der}(A, \mathbb{Q}; \epsilon)$  and the induced map  $(\epsilon_*, \epsilon_*) : \text{Rel}(\phi^*) \rightarrow \text{Rel}(\widehat{\phi}^*)$ . The relative Gottlieb group  $G_*^{\text{rel}}(A, B; \phi)$  is the image of  $H_*(\epsilon_*, \epsilon_*)$ . From the tower

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Der}(A, H^*(B); \phi) & \xrightarrow{J} & \text{Rel}(\phi^*) & \xrightarrow{P} & \text{Der}(B, H^*(B); \gamma) \longrightarrow 0 \\ & & \downarrow \epsilon_* & & \downarrow (\epsilon_*, \epsilon_*) & & \downarrow \epsilon_* \\ 0 & \longrightarrow & \text{Der}(A, \mathbb{Q}; \epsilon) & \xrightarrow{\widehat{J}} & \text{Rel}(\widehat{\phi}^*) & \xrightarrow{\widehat{P}} & \text{Der}(B, \mathbb{Q}; \epsilon) \longrightarrow 0 \end{array}$$

one gets a sequence

$$\cdots \rightarrow G_{k+1}(B, H^*(B), \gamma) \rightarrow G_k(A, H^*(B), \phi^*) \rightarrow G_k^{\text{rel}}(A, H^*(B), \phi^*) \rightarrow \cdots$$

called  $G$ -sequence of  $\phi$ .

**Proposition 15.** *The  $G$ -sequence associated to the inclusion  $\text{aut}_1 \mathbb{C}P^n \rightarrow \text{map}(\mathbb{C}P^n, \mathbb{C}P^{n+k}; i_{n,k})$  is not exact.*

*Proof.* Clearly  $G_*(B, H^*(B); \gamma) = \langle [(y_{2n+1}, 1)] \rangle$  and similarly

$$G_*(A, H^*(B), \phi) = \langle [(x_2, 1)], [(x_{2n+2k+1}, 1)] \rangle.$$

We consider first the case where  $k > n$ . Then the only non zero differential on  $\text{Rel}(\phi^*) = (sL \oplus L', d)$  is given by

$$d(s\delta_2, 0) = (-s\theta_1, 0) + (0, \phi^*(\delta_2)) = (-s\theta_1, 0) + (0, \beta_2).$$

Similarly the only non zero differential on

$$\text{Rel}(\widehat{\phi}^*) = \langle (sy_2^*, 0), (sy_{2n+1}^*, 0), (0, x_2^*), (0, x_{2n+2k+1}^*) \rangle$$

is  $d(sy_2^*, 0) = (0, x_2^*)$ . We conclude that

$$\begin{aligned} G_*^{\text{rel}}(A, H^*(B), \phi) &= \langle [(sy_{2n+1}^*, 0)], (0, x_{2n+2k+1}^*) \rangle \\ &\cong sG_*(\mathbb{C}P^n) \oplus G_*(\mathbb{C}P^{n+k}). \end{aligned}$$

Hence in the  $G$ -sequence reduces to fragments

$$0 \rightarrow G_{2n+2}^{\text{rel}}(A, H^*(B); \phi^*) \xrightarrow{\cong} G_{2n+1}(B, H^*(B); \gamma) \rightarrow 0,$$

$$0 \rightarrow G_{2n+2k+1}(A, H^*(B); \phi^*) \xrightarrow{\cong} G_{2n+2k+1}^{\text{rel}}(A, H^*(B); \phi^*) \rightarrow 0$$

and terminates with

$$0 \rightarrow G_2(A, H^*(B); \phi^*) \rightarrow 0.$$

As  $G_2(A, H^*(B); \phi^*) \cong \mathbb{Q}$ , we conclude that the last fragment of the  $G$ -sequence is not exact.

If  $k \leq n$ , then  $\phi^*(\theta_{2n+1}) = \alpha_{2n+1}$ , hence  $d(s\theta_{2n+1}, 0) = (0, \alpha_{2n+1})$ , therefore  $[(sy_{2n+1}^*, 0)] \in H_*(\text{Rel}(\widehat{\phi}^*))$  is not in the image of  $H_*(\epsilon_*, \epsilon_*)$ . The only change in the  $G$ -sequence is the fragment

$$0 \rightarrow G_{2n+2}^{\text{rel}}(A, H^*(B); \phi^*) \rightarrow 0,$$

which is not exact as well, as  $G_{2n+2}^{\text{rel}}(A, H^*(B)) \cong \mathbb{Q}$ .  $\square$



## References

- [1] J. Block and A. Lazarev, *André-Quillen cohomology and rational homotopy of function spaces*, Adv. Math. **193** (2005), no. 1, 18–39. <https://doi.org/10.1016/j.aim.2004.04.014>
- [2] E. H. Brown, Jr., and R. H. Szczarba, *On the rational homotopy type of function spaces*, Trans. Amer. Math. Soc. **349** (1997), no. 12, 4931–4951. <https://doi.org/10.1090/S0002-9947-97-01871-0>
- [3] U. Buijs, Y. Félix, and A. Murillo, *Lie models for the components of sections of a nilpotent fibration*, Trans. Amer. Math. Soc. **361** (2009), no. 10, 5601–5614. <https://doi.org/10.1090/S0002-9947-09-04870-3>
- [4] ———,  *$L_\infty$  models of based mapping spaces*, J. Math. Soc. Japan **63** (2011), no. 2, 503–524. <http://projecteuclid.org/euclid.jmsj/1303737796>
- [5] ———,  *$L_\infty$  rational homotopy of mapping spaces*, Rev. Mat. Complut. **26** (2013), no. 2, 573–588. <https://doi.org/10.1007/s13163-012-0105-z>
- [6] Y. Félix, S. Halperin, and J.-C. Thomas, *Rational homotopy theory*, Graduate Texts in Mathematics, 205, Springer-Verlag, New York, 2001. <https://doi.org/10.1007/978-1-4613-0105-9>
- [7] J.-B. Gatsinzi, *On the genus of elliptic fibrations*, Proc. Amer. Math. Soc. **132** (2004), no. 2, 597–606. <https://doi.org/10.1090/S0002-9939-03-07203-4>
- [8] J.-B. Gatsinzi and R. Kwashira, *Rational homotopy groups of function spaces*, in Homotopy theory of function spaces and related topics, 105–114, Contemp. Math., 519, Amer. Math. Soc., Providence, RI, 2010. <https://doi.org/10.1090/comm/519/10235>
- [9] D. H. Gottlieb, *Evaluation subgroups of homotopy groups*, Amer. J. Math. **91** (1969), 729–756. <https://doi.org/10.2307/2373349>
- [10] A. Haefliger, *Rational homotopy of the space of sections of a nilpotent bundle*, Trans. Amer. Math. Soc. **273** (1982), no. 2, 609–620. <https://doi.org/10.2307/1999931>
- [11] T. Lada and M. Markl, *Strongly homotopy Lie algebras*, Comm. Algebra **23** (1995), no. 6, 2147–2161. <https://doi.org/10.1080/00927879508825335>
- [12] G. Lupton and S. B. Smith, *Rationalized evaluation subgroups of a map. I. Sullivan models, derivations and G-sequences*, J. Pure Appl. Algebra **209** (2007), no. 1, 159–171. <https://doi.org/10.1016/j.jpaa.2006.05.018>
- [13] ———, *Rationalized evaluation subgroups of a map. II. Quillen models and adjoint maps*, J. Pure Appl. Algebra **209** (2007), no. 1, 173–188. <https://doi.org/10.1016/j.jpaa.2006.05.019>
- [14] S. Mac Lane, *Homology*, Die Grundlehren der mathematischen Wissenschaften, Bd. 114, Academic Press, Inc., Publishers, New York, 1963.
- [15] J. M. Møller and M. Raussen, *Rational homotopy of spaces of maps into spheres and complex projective spaces*, Trans. Amer. Math. Soc. **292** (1985), no. 2, 721–732. <https://doi.org/10.2307/2000242>
- [16] D. Sullivan, *Infinitesimal computations in topology*, Inst. Hautes Études Sci. Publ. Math. No. 47 (1977), 269–331 (1978).
- [17] M. H. Woo and K. Y. Lee, *On the relative evaluation subgroups of a CW-pair*, J. Korean Math. Soc. **25** (1988), no. 1, 149–160.

J.-B. GATSINZI

DEPARTMENT OF MATHEMATICS AND STATISTICAL SCIENCES  
BOTSWANA INTERNATIONAL UNIVERSITY OF SCIENCE AND TECHNOLOGY  
PRIVATE BAG 16, PALAPYE, BOTSWANA  
Email address: [gatsinzi@biust.ac.bw](mailto:gatsinzi@biust.ac.bw)