

## WHEN IS $C(X)$ AN EM-RING?

EMAD ABUOSBA AND ISAAF ATASSI

ABSTRACT. A commutative ring with unity  $R$  is called an EM-ring if for any finitely generated ideal  $I$  there exist  $a$  in  $R$  and a finitely generated ideal  $J$  with  $\text{Ann}(J) = 0$  and  $I = aJ$ . In this article it is proved that  $C(X)$  is an EM-ring if and only if for each  $U \in \text{Coz}(X)$ , and each  $g \in C^*(U)$  there is  $V \in \text{Coz}(X)$  such that  $U \subseteq V$ ,  $\bar{V} = X$ , and  $g$  is continuously extendable on  $V$ . Such a space is called an EM-space. It is shown that EM-spaces include a large class of spaces as F-spaces and cozero complemented spaces. It is proved among other results that  $X$  is an EM-space if and only if the Stone-Čech compactification of  $X$  is.

### 1. Introduction

Let  $X$  be a topological space,  $C(X)$  be the ring of all continuous real valued functions defined on  $X$  and  $C^*(X)$  be its subring of bounded functions. For each  $f \in C(X)$ , let  $Z(f) = f^{-1}(0)$ ,  $\text{coz}(f) = X - Z(f)$  and  $\text{supp}(f) = \overline{\text{coz}(f)}$ . Let  $Z(X)$  be the set of all zero sets in  $X$  and  $\text{Coz}(X)$  be the set of all cozero sets in  $X$ . For any undefined terms, the reader may refer to [9], and for a new survey and results on  $C(X)$ , see [4].

If  $X$  is any topological space, then there is a Tychonoff space  $Y$  such that  $C(X)$  is isomorphic to  $C(Y)$ . Thus we will assume that all spaces  $X$  are Tychonoff spaces, and so we are able to extend  $X$  into the Stone-Čech compactification  $\beta X$ .

A lot of work is done in the literature to characterize algebraic properties of  $C(X)$  using the topological properties of  $X$  and viceversa. In the following some of these characterizations that will be used in this article.

A space  $X$  is called basically disconnected if for each  $f \in C(X)$ ,  $\text{supp}(f)$  is open. It is known that  $X$  is basically disconnected if and only if  $C(X)$  is a PP-ring (every principal ideal is projective), see [5].

---

Received December 4, 2020; Revised April 7, 2021; Accepted April 22, 2021.

2010 *Mathematics Subject Classification*. 13A15, 54C30, 54C45.

*Key words and phrases*.  $C(X)$ , EM-ring, generalized morphic ring, PP-ring, PF-ring, basically disconnected space, F-space, cozero complemented space.

This article is a part of the Ph.D. thesis prepared by the second author under supervision of the first in The University of Jordan.

A space  $X$  is called an F-space if for each  $f \in C(X)$ ,  $\text{coz}(f)$  is  $C^*$ -embedded in  $X$ . It is known that  $X$  is an F-space if and only if  $C(X)$  is a Bezout ring (every finitely generated ideal is principal) if and only if  $C(X)$  is a PF-ring (every principal ideal is flat), see [8, 9].

A space  $X$  is called cozero complemented if and only if for each  $U \in \text{Coz}(X)$  there exists  $V \in \text{Coz}(X)$  such that  $U \cap V = \emptyset$  and  $\overline{U \cup V} = X$ . It is known that  $X$  is cozero complemented if and only if  $\text{Min}(C(X))$  (the space of all minimal prime ideals in  $C(X)$  with the Zariski topology) is compact, see [12].

Let  $R$  be a commutative ring. In [1], the authors introduced the notion of an annihilating content for a polynomial: if  $f(x) \in R[x]$  and there exist  $a \in R$  and a regular (non-zero divisor) polynomial  $g(x) \in R[x]$  such that  $f(x) = ag(x)$ , then  $a$  is called an annihilating content for  $f(x)$ . Annihilating content simplifies computing the annihilator of a polynomial, which is not always an easy task, also it is used to find the annihilator of a finitely generated ideal. In [2], the authors defined EM-rings as those rings for which any polynomial in  $R[x]$  has an annihilating content. Among other things it was shown that a Bezout ring is an EM-ring, the class of EM-rings is closed under localization, direct products, polynomial adjunction and that a Noetherian ring is an EM-ring if and only if each minimal prime ideal is principal. More investigation was done in [3], and it was proved that  $R$  is an EM-ring if and only if for each finitely generated ideal  $I$  of  $R$ , there exist  $a \in R$  and a finitely generated ideal  $J$  of  $R$  such that  $I = aJ$  and  $\text{Ann}(J) = 0$ .

In [7], the authors proved that a ring  $R$  is a PP-ring if and only if  $R(+)$  is an EM-ring if and only if  $R(+)$  is a generalized morphic ring (for each  $f \in R$ ,  $\text{Ann}(f) = \{g \in R : fg = 0\}$  is principal). They also showed in [2] that a Noetherian ring is an EM-ring if and only if it is a generalized morphic ring, however they showed that if  $X = \beta\mathbb{N} - \mathbb{N}$ , then  $C(X)$  is an EM-ring that is not generalized morphic. This motivated us to study EM-rings and generalized morphic rings in  $C(X)$ .

The purpose of this article is to characterize those topological spaces  $X$  for which  $C(X)$  is an EM-ring or a generalized morphic ring. Although we don't know yet the exact relation between EM-rings and generalized morphic rings, but we manage to show that if  $C(X)$  is generalized morphic, then indeed it is an EM-ring.

In Section 2, we study the annihilating content of a finitely generated ideal in  $C(X)$  and simplify some of its computations. It is shown (Theorem 2.3) that the finitely generated ideal  $I = (f_1, f_2, \dots, f_n)$  of  $C(X)$  has an annihilating content  $h \in C(X)$  if and only if there exist  $g_1, \dots, g_{n+1} \in C(X)$  such that  $f_i = hg_i$  for  $1 \leq i \leq n$  and  $0 = hg_{n+1}$  with  $\bigcup_{i=1}^{n+1} \text{supp}(g_i) = X$ . Also it is shown (Theorem 2.4) that we can pick the annihilating content  $h$  (which is not unique) to be bounded.

In Section 3, we define EM-spaces as follows:  $X$  is an EM-space if and only if  $C(X)$  is an EM-ring. It is shown that the set of all real numbers with the

Euclidean topology is an EM-space, while the one-point compactification of an uncountable discrete space is not. It is shown (Corollary 3.9) that  $X$  is an EM-space if and only if  $\beta X$  is. Theorem 3.11 characterizes Tychonoff EM-spaces: a Tychonoff space  $X$  is an EM-space if and only if for each  $U \in \text{Coz}(X)$ , and each  $g \in C^*(U)$  there is  $V \in \text{Coz}(X)$  such that  $U \subseteq V$ ,  $\overline{V} = X$ , and  $g$  is continuously extendable on  $V$ .

In Section 4, we relate EM-spaces with other spaces. We observe that the ring  $C(X)$  is generalized morphic if and only if  $X$  is basically disconnected, and so if  $C(X)$  is generalized morphic, then it is an EM-ring. It is shown that EM-spaces include wide range of spaces such as F-spaces and cozero complemented spaces (Theorem 4.1). It is also deduced that a quazi F-space EM-space is an F-space. Finally it is shown that a locally connected EM-space is cozero complemented.

## 2. Annihilating content

The idea of an annihilating content was first defined in [1] to factor a polynomial into an element in the ring multiplied by a regular polynomial to simplify calculating the zero divisor graph of the polynomial ring  $R[x]$ . In this section we use topological characterization for the annihilating content in  $C(X)$ .

**Definition 2.1.** Let  $R$  be a commutative ring and let  $f(x) \in R[x]$ . If there exist  $a \in R$  and a regular polynomial  $g(x) \in R[x]$  such that  $f(x) = ag(x)$ , then  $a$  is called an *annihilating content* for  $f(x)$ .

The idea of annihilating content was more developed in [2] and [3] to be used for finitely generated ideals of  $R$ .

**Definition 2.2.** Let  $R$  be a commutative ring with unity, and let  $I$  be a finitely generated ideal of  $R$  such that there exist  $a \in R$  and a finitely generated ideal  $J$  with  $\text{Ann}(J) = 0$  and  $I = aJ$ . In this case  $a$  is called an *annihilating content* for  $I$ .

It was shown in [3] that the annihilating content is not unique, and if  $a$  is an annihilating content for  $f(x)$  (the finitely generated ideal  $I$ ), then  $\text{Ann}(f(x)) = \text{Ann}(a)$  ( $\text{Ann}(I) = \text{Ann}(a)$ ), also if  $aR = bR$ , then  $b$  is an annihilating content for  $f(x)$  (for  $I$ ), but not conversely.

We now give the analogue definition of an annihilating content in the ring  $C(X)$ . But we recall first that for  $f, h \in C(X)$ ,  $\text{supp}(f) = \text{supp}(h)$  if and only if  $\text{Ann}(f) = \text{Ann}(h)$  and that  $\text{Ann}(f) = 0$  if and only if  $\text{coz}(f)$  is dense in  $X$ .

**Theorem 2.3.** In a Tychonoff space  $X$ , the finitely generated ideal  $I = (f_1, f_2, \dots, f_n)$  has an annihilating content  $h \in C(X)$  if and only if there exist  $g_1, g_2, \dots, g_{n+1} \in C(X)$  such that

$$\begin{aligned} f_i &= hg_i, & i &= 1, 2, \dots, n, \\ 0 &= hg_{n+1}, \end{aligned}$$

$$\bigcup_{i=1}^{n+1} \text{supp}(g_i) = X.$$

In this case,  $\text{supp}(h) = \bigcup_{i=1}^n \text{supp}(f_i)$ .

*Proof.* Assume  $I = (f_1, f_2, \dots, f_n)$  has an annihilating content. Then there exist  $h, g_1, g_2, \dots, g_n, g'_{n+1}, \dots, g'_m \in C(X)$  such that

$$\begin{aligned} f_i &= hg_i, \quad i = 1, 2, \dots, n, \\ 0 &= hg'_i, \quad i = n+1, \dots, m, \\ \text{Ann}(g_1, g_2, \dots, g_n, g'_{n+1}, \dots, g'_m) &= 0. \end{aligned}$$

Let  $g_{n+1} = \sum_{i=n+1}^m |g'_i|$ . Then it is clear that  $hg_{n+1} = 0$ , and that  $\text{Ann}(g_1, g_2, \dots, g_n, g_{n+1}) = 0$ , and so  $\bigcup_{i=1}^{n+1} \text{supp}(g_i) = X$ . The converse is straightforward.

Now to show that  $\text{supp}(h) = \bigcup_{i=1}^n \text{supp}(f_i)$ , it is sufficient to show that  $\text{Ann}(h) = \text{Ann}(f_1, f_2, \dots, f_n)$ . If  $\alpha f_i = 0$  for each  $i$ , then  $(\alpha h)g_i = 0$  for each  $i$ , and so  $\alpha h \in \text{Ann}(g_1, g_2, \dots, g_m) = 0$ . Hence the result.  $\square$

In the following, we show that the annihilating content for an ideal in  $C(X)$ , if exists, can be chosen to be bounded.

**Theorem 2.4.** *In a Tychonoff space  $X$ , if the finitely generated ideal  $I = (f_1, f_2, \dots, f_n)$  in  $C(X)$  has an annihilating content, then  $I$  has a bounded annihilating content.*

*Proof.* Let  $h_0 \in C(X)$  be an annihilating content for  $I = (f_1, f_2, \dots, f_n)$ . Then there exist  $g_1, g_2, \dots, g_{n+1} \in C(X)$  such that

$$\begin{aligned} f_i &= h_0 g_i, \quad i = 1, 2, \dots, n, \\ 0 &= h_0 g_{n+1}, \\ \bigcup_{i=1}^{n+1} \text{supp}(g_i) &= X. \end{aligned}$$

Let

$$q(x) = \begin{cases} -h_0(x), & x \in h_0^{-1}((-\infty, -1]), \\ 1, & x \in h_0^{-1}(-1, 1), \\ h_0(x), & x \in h_0^{-1}([1, \infty)). \end{cases}$$

Then  $q \in C(X)$ . Since  $Z(q) = \emptyset$  and  $\left| \frac{h_0}{q} \right| \leq 1$ , we have  $h = \frac{h_0}{q} \in C^*(X)$ . If  $g_i^* = g_i q$ , then

$$hg_i^* = hg_i q = h_0 g_i = \begin{cases} f_i, & i = 1, 2, \dots, n, \\ 0, & i = n+1. \end{cases}$$

Actually, for every  $i \in \{1, 2, \dots, n+1\}$ ,  $\text{supp}(g_i^*) = \text{supp}(g_i)$ .

Therefore,  $\bigcup_{i=1}^{n+1} \text{supp}(g_i^*) = X$ , and  $h$  is a bounded annihilating content for  $I$ .  $\square$

### 3. EM-spaces

In this section, we will give a topological characterization for EM-spaces, give examples and counter examples, and show that  $X$  is an EM-space if and only if  $\beta X$  is.

The idea of EM-rings was firstly defined in [2], then further investigations were carried out in [3].

**Definition 3.1.** A commutative ring  $R$  is called an *EM-ring* if every polynomial in  $R[x]$  has an annihilating content.

The following Theorem can be found in [3], it gives more equivalent conditions to EM-rings and simplifies computations.

**Theorem 3.2.** For a commutative ring  $R$ , the following are equivalent.

- (1)  $R$  is an EM-ring.
- (2) For  $a + bx \in R[x]$ , there exist an element  $c \in R$  and a regular polynomial  $f_1 \in R[x]$  with  $a + bx = cf_1$ .
- (3) For  $a, b \in R$ , there are an element  $c \in R$  and a finitely generated ideal  $J$  of  $R$  with  $(a, b) = cJ$  and  $\text{Ann}(J) = 0$ .

Now we define EM-spaces.

**Definition 3.3.** A Tychonoff spaces  $X$  is called an *EM-space* if  $C(X)$  is an EM-ring.

In view of Definition 3.3, Theorem 2.3, and Theorem 3.2, we give a simple formula for being an EM-space.

**Corollary 3.4.** A Tychonoff spaces  $X$  is an EM-space if and only if whenever  $f_1, f_2 \in C(X)$ , there exist  $h, g_1, g_2, g_3 \in C(X)$  such that  $f_i = hg_i$  for  $i = 1, 2$ ,  $hg_3 = 0$ , and  $\bigcup_{i=1}^3 \text{supp}(g_i) = X$ .

**Example 3.5.** It is proved in [2] that any Bezout ring is an EM-ring, and it is proved in [8] that  $C(X)$  is Bezout if and only if  $X$  is an F-space. Now we give a direct proof that if  $X$  is an F-space, then  $X$  is an EM-space. For an ideal  $I = (f_1, f_2) \subseteq C(X)$ , take  $h = |f_1| + |f_2|$ , and let  $g_i \in C(X)$  such that  $g_i|_{\bigcup_{j=1,2} \text{coz}(f_j)} = \frac{f_i}{h}$  for  $i = 1, 2$ , and  $g_3 = 1 - |g_1| - |g_2|$ . Then  $f_i = hg_i$  for  $i = 1, 2$ ,  $hg_3 = 0$  and  $\bigcup_{i=1}^3 \text{supp}(g_i) = X$ . This implies that  $h$  is an annihilating content for  $I$ .

**Example 3.6.** Let  $\mathbb{R}$  be the set of all real numbers with the Euclidean topology, and let  $f_1, f_2 \in C(\mathbb{R})$ . Define  $h = (|f_1| + |f_2|)^{\frac{1}{2}}$ . It is clear that  $\text{coz}(h) = \text{coz}(f_1) \cup \text{coz}(f_2)$ . For  $i = 1, 2$ , define

$$\alpha_i(x) = \begin{cases} \frac{f_i}{h}(x), & x \in \text{coz}(h), \\ 0, & \text{otherwise.} \end{cases}$$

Then  $\alpha_i \in C(X)$ , and  $f_i = h\alpha_i$  for  $i = 1, 2$ . Now, define  $\gamma(x) = \inf\{|x - a| : a \in \text{supp}(h)\}$ . It is clear that  $h\gamma = 0$ ,  $Z(\gamma) = \text{supp}(h)$ , and  $\text{coz}(\alpha_1) \cup \text{coz}(\alpha_2) \cup$

$\text{coz}(\gamma) = \text{coz}(f_1) \cup \text{coz}(f_2) \cup \text{coz}(\gamma) = \text{coz}(h) \cup \text{coz}(\gamma)$  is dense in  $\mathbb{R}$ . Thus,  $C(\mathbb{R})$  is an EM-ring.

In general if  $X$  is a metric space, then  $C(X)$  is an EM-ring.

One can wonder if for any Tychonoff space  $X$ , the ring  $C(X)$  is an EM-ring, since if  $f_1, f_2 \in C(X)$ , then using the same technique in the last example one shows that  $(f_1, f_2) = h(\alpha_1, \alpha_2)$ , but it is not always the case that we can find  $\gamma$  satisfying  $\text{supp}(\alpha_1) \cup \text{supp}(\alpha_2) \cup \text{supp}(\gamma) = X$  with  $h\gamma = 0$ . This will be shown in the following example.

**Example 3.7.** Let  $X = Y \cup \{\infty\}$  be the one-point compactification of an uncountable discrete space. For  $i = 1, 2$ , take  $f_i \in C(X)$  such that  $\text{coz}(f_i)$  are infinite disjoint countable sets. Suppose that  $h, g_i \in C(X)$  for  $i = 1, 2$ , and  $f_i = hg_i$ . Then  $\text{coz}(g_1) \cap \text{coz}(f_2) = \text{coz}(g_1) \cap \text{coz}(h) \cap \text{coz}(f_2) = \text{coz}(f_1) \cap \text{coz}(f_2) = \emptyset$ . But  $\text{coz}(f_2)$  is infinite, so  $\infty \notin \text{coz}(g_1)$ . Hence,  $\text{coz}(g_1)$  is countable being an  $F_\sigma$ -set in  $X$ . Similar argument shows that  $\text{coz}(g_2)$  is also countable.

Let  $g$  be any function in  $C(X)$ . If  $\infty \in \text{coz}(g)$ , then  $\text{coz}(g) \cap \text{coz}(h) \supseteq \text{coz}(g) \cap \text{coz}(f_1) \neq \emptyset$ , because  $\text{coz}(f_1)$  is infinite, and thus  $gh \neq 0$ .

If  $\infty \notin \text{coz}(g)$ , then  $\text{coz}(g)$  is countable, and  $\bigcup_{i=1}^2 \text{supp}(g_i) \cup \text{supp}(g) = \bigcup_{i=1}^2 \text{coz}(g_i) \cup \text{coz}(g) \cup \{\infty\} \neq X$  since the left hand side is countable, but the right hand side is uncountable. Therefore, the ideal  $(f_1, f_2)$  does not have an annihilating content, and so  $C(X)$  is not an EM-ring.

**Theorem 3.8.** *Let  $X$  be a Tychonoff space. Then  $C^*(X)$  is an EM-ring if and only if  $C(X)$  is an EM-ring.*

*Proof.* Let  $X$  be a space such that  $C^*(X)$  is an EM-ring. Let  $f, g \in C(X)$ , and let  $f^* = (f \wedge 1) \vee (-1)$ , and  $g^* = (g \wedge 1) \vee (-1)$ . By assumption, there exist  $h, \alpha^*, \beta^*, \gamma \in C^*(X)$ , such that  $f^* = h\alpha^*$ ,  $g^* = h\beta^*$ ,  $0 = h\gamma$ , and  $\text{supp}(\alpha^*) \cup \text{supp}(\beta^*) \cup \text{supp}(\gamma) = X$ . Consider

$$q_1(x) = \begin{cases} \frac{f}{f^*}(x), & x \in f^{-1}((-\infty, -1] \cup [1, \infty)), \\ 1, & \text{elsewhere,} \end{cases}$$

$$q_2(x) = \begin{cases} \frac{g}{g^*}(x), & x \in g^{-1}((-\infty, -1] \cup [1, \infty)), \\ 1, & \text{elsewhere,} \end{cases}$$

and let  $\alpha = \alpha^*q_1$ , and  $\beta = \beta^*q_2$ . Then  $\alpha h = \alpha^*q_1 h = f^*q_1 = f$ , and  $\beta h = \beta^*q_2 h = g^*q_2 = g$ . Furthermore,  $0 = h\gamma$ , and

$$\text{supp}(\alpha) \cup \text{supp}(\beta) \cup \text{supp}(\gamma) = \text{supp}(\alpha^*) \cup \text{supp}(\beta^*) \cup \text{supp}(\gamma) = X.$$

This implies that  $h$  is an annihilating content for  $(f, g)$ , and  $C(X)$  is an EM-ring.

Conversely, let  $C(X)$  be an EM-ring. Let  $f, g \in C^*(X) \subseteq C(X)$ . Then, by assumption, and by Theorem 2.4, we can find  $h_0 \in C^*(X)$ , and  $\alpha_0, \beta_0, \gamma_0 \in C(X)$  such that  $f = \alpha_0 h_0$ ,  $g = \beta_0 h_0$ ,  $0 = \gamma_0 h_0$ , and  $\text{supp}(\alpha_0) \cup \text{supp}(\beta_0) \cup$

$\text{supp}(\gamma_0) = X$ . Evidently, there is a bound  $M > 0$ , with  $|f|, |g|, |h_0| < M$ .  
Take

$$\gamma = (\gamma_0 \wedge M) \vee (-M)$$

and let  $\alpha_0^* = (\alpha_0 \wedge M) \vee (-M)$ . For  $A = \alpha_0^{-1}((-\infty, -M] \cup [M, \infty))$ , take

$$q_1(x) = \begin{cases} \frac{\alpha_0}{\alpha_0^*}(x), & x \in A, \\ 1, & \text{elsewhere,} \end{cases}$$

and then let  $\beta_1 = \frac{\beta_0}{q_1} \in C(X)$  since  $Z(q_1) = \emptyset$ , and  $\beta = (\beta_1 \wedge M) \vee (-M)$ .  
For  $B = \beta_1^{-1}((-\infty, -M] \cup [M, \infty))$ , take

$$q_2(x) = \begin{cases} \frac{\beta_1}{\beta}(x), & x \in B, \\ 1, & \text{elsewhere,} \end{cases}$$

and then let  $\alpha = \frac{\alpha_0^*}{q_2} \in C(X)$ ,  $Z(q_2) = \emptyset$ , and  $h = q_1 q_2 h_0$ . In fact,

$$|\alpha| = \left| \frac{\alpha_0^*}{q_2} \right| = \begin{cases} \left| \frac{\alpha_0^* \beta}{\beta_1} \right| \leq |\alpha_0^*| \leq M & \text{on } B, \\ |\alpha_0^*| \leq M & \text{on } X - B, \end{cases}$$

$$|h| = |q_1 q_2 h_0| = \begin{cases} |h_0| \leq M & \text{on } X - (A \cup B), \\ \left| \frac{\alpha_0}{\alpha_0^*} h_0 \right| = \left| \frac{f}{\alpha_0^*} \right| = \frac{|f|}{M} \leq 1 & \text{on } A - B, \\ \left| \frac{\beta_1 h_0}{\beta} \right| = \left| \frac{\beta_0 h_0}{\beta} \right| = \left| \frac{g}{\beta} \right| = \frac{|g|}{M} \leq 1 & \text{on } B - A, \\ \left| \frac{\alpha_0 \beta_1 h_0}{\alpha_0^* \beta} \right| = \left| \frac{\beta_0 h_0}{\beta} \right| = \frac{|g|}{M} \leq 1 & \text{on } A \cap B. \end{cases}$$

By this we have  $|h| \leq \max\{M, 1\}$ ,  $\alpha, \beta, \gamma, h \in C^*(X)$ , and having same support as  $\alpha_0, \beta_0, \gamma_0, h_0$ , respectively, since  $q_1, q_2$  are units.

Additionally,

$$\begin{aligned} \alpha h &= \alpha_0^* q_1 h_0 = \alpha_0 h_0 = f, \\ \beta h &= \beta q_1 q_2 h_0 = \beta_1 q_1 h_0 = \beta_0 h_0 = g, \\ \gamma h &= 0, \\ \text{supp}(\alpha) \cup \text{supp}(\beta) \cup \text{supp}(\gamma) &= X. \end{aligned}$$

Consequently,  $C^*(X)$  is an EM-ring.  $\square$

Since  $C^*(X)$  is isomorphic to  $C(\beta X)$ , see [9], we get the following result.

**Corollary 3.9.** *Let  $X$  be a Tychonoff space. Then  $C(X)$  is an EM-ring if and only if  $C(\beta X)$  is an EM-ring.*

We now go forward to give a topological characterization of an EM-space, but first we will need the following lemma.

**Lemma 3.10.** *Let  $X$  be a Tychonoff EM-space. Then for every  $U \in \text{Coz}(X)$  and every  $g \in C^*(U)$ , there exists a zero set  $Z \in Z(X)$  such that  $U \subseteq X - Z$ ,  $\text{Int } Z = \emptyset$ , and  $g$  is continuously extendable on  $U \cup (\partial U - Z)$ .*

*Proof.* Let  $U = \text{coz}(f)$  be a cozero set where  $f \in C^*(X)$ , and let  $g \in C^*(U)$ . Then

$$k = \begin{cases} fg & \text{on } U, \\ 0 & \text{on } X - U, \end{cases}$$

is continuous. As  $C(X)$  is an EM-ring, there exist  $\alpha, \beta, \gamma, h \in C(X)$  such that  $f = \alpha h$ ,  $k = \beta h$ ,  $0 = \gamma h$ , and  $\text{supp}(\alpha) \cup \text{supp}(\beta) \cup \text{supp}(\gamma) = X$ . Let  $Z = Z(|\alpha| + |\beta| + |\gamma|)$ . Then  $U \subseteq X - Z$ , and  $\text{Int } Z = \emptyset$ .

Notice that  $U \subseteq \text{coz}(\alpha) \cap \text{coz}(f) \cap \text{coz}(h)$ . Thus, on  $U$ ,

$$\left| \frac{\beta}{\alpha} \right| = \left| \frac{k}{f} \right| = |g| < M$$

for some  $M > 0$ . Thus for all  $x \in U$ ,

$$(*) \quad \frac{|\beta(x)| + |\alpha(x)|}{M + 1} < |\alpha(x)|.$$

Let  $x_0 \in \partial U - Z$ . Then  $\gamma(x_0) = 0$ , since  $\bar{U} = \text{supp}(f) \subseteq \text{supp}(h) \subseteq Z(\gamma)$ . But  $x_0 \notin Z$ , thus  $x_0 \in \text{coz}(|\alpha| + |\beta| + |\gamma|)$ , and so  $(|\alpha| + |\beta|)(x_0) > 0$ . By continuity of  $|\alpha| + |\beta|$ , there is a neighborhood  $U_1$  of  $x_0$  such that for each  $x \in U_1$ ,

$$(**) \quad \frac{3}{4} (|\alpha| + |\beta|)(x_0) < (|\alpha| + |\beta|)(x) < \frac{5}{4} (|\alpha| + |\beta|)(x_0).$$

If  $\alpha(x_0) = 0$ , then by continuity of  $\alpha$ , there is a neighborhood  $U_2$  of  $x_0$  such that for each  $x \in U_2$ ,

$$|\alpha(x)| < \frac{1}{4} \left( \frac{(|\alpha| + |\beta|)(x_0)}{1 + M} \right)$$

and by (\*), for each  $x \in U \cap U_2$ ,

$$(|\alpha| + |\beta|)(x) < \frac{1}{4} (|\alpha| + |\beta|)(x_0).$$

Clearly, this contradicts (\*\*) as  $U \cap U_1 \cap U_2 \neq \emptyset$ . Therefore,  $\alpha(x_0) \neq 0$ . In other words,  $\partial U - Z \subseteq \text{coz}(\alpha)$ . Evidently,

$$g^* = \begin{cases} g & \text{on } U \\ \frac{\beta}{\alpha} & \text{on } \partial U - Z \end{cases} \in C^*(U \cup (\partial U - Z))$$

is the desired extension.  $\square$

We now give a topological characterization for EM-spaces.

**Theorem 3.11.** *Let  $X$  be a Tychonoff space. Then  $X$  is an EM-space if and only if for each  $U \in \text{Coz}(X)$ , and each  $g \in C^*(U)$  there is  $V \in \text{Coz}(X)$  such that  $U \subseteq V$ ,  $\bar{V} = X$ , and  $g$  is continuously extendable on  $V$ .*

*Proof.* ( $\Rightarrow$ ) Assume  $X$  is an EM-space. Let  $U \in \text{Coz}(X)$  and  $f \in C^*(X)$  such that  $U = \text{coz}(f)$ . Let  $g \in C^*(U)$ , and then

$$k = \begin{cases} fg & \text{on } U \\ 0 & \text{on } X - U \end{cases} \in C^*(X).$$



Let  $f^\beta, k^\beta \in C(\beta X)$  be the extensions of  $f$  and  $k$  in  $\beta X$ , respectively. If  $\tilde{U} = \text{coz}(f^\beta)$ , then  $\tilde{g} = \frac{k^\beta}{f^\beta} \in C^*(\tilde{U})$ , and  $\tilde{g}|_U = g$ . On the other hand,  $C(\beta X)$  is an EM-ring by Corollary 3.9. Hence, by Lemma 3.10, there are  $\tilde{\sigma} \in C(\beta X)$  with  $\tilde{U} \subseteq \beta X - Z(\tilde{\sigma})$ ,  $\text{Int}_{\beta X} Z(\tilde{\sigma}) = \emptyset$ , and  $\tilde{g}_1 \in C^*(\tilde{U} \cup (\partial\tilde{U} - Z(\tilde{\sigma})))$  such that  $\tilde{g}_1$  is an extension of  $\tilde{g}$ . Now,

$$\tilde{g}_\sigma = \begin{cases} \tilde{g}_1 \tilde{\sigma} & \text{on } \tilde{U} \cup (\partial\tilde{U} - Z(\tilde{\sigma})) \\ 0 & \text{on } \partial\tilde{U} \cap Z(\tilde{\sigma}) \end{cases} \in C^*(\tilde{U}).$$

By Tietze-Urysohn Theorem,  $\tilde{g}_\sigma$  has an extension on  $\beta X$ , say  $\tilde{g}_\sigma$ . Moreover, on  $U$ ,  $\tilde{g}_\sigma = g\sigma$  for  $\sigma = \tilde{\sigma}|_X$ . Take  $V = \text{coz}(\sigma)$ . Then  $\bar{V} = X$  (since  $\text{Int}_{\beta X} Z(\tilde{\sigma}) = \emptyset$ ), and

$$g^* = \frac{\tilde{g}_\sigma|_V}{\sigma} \in C^*(V)$$

is the desired extension of  $g$ .

( $\Leftarrow$ ) Let  $X$  be the prescribed space. Let  $f_1, f_2 \in C(X)$ , and  $U = \text{coz}(f_1) \cup \text{coz}(f_2)$ . Then  $\frac{f_1}{|f_1|+|f_2|}, \frac{f_2}{|f_1|+|f_2|} \in C^*(U)$ . So by assumption, there exist dense  $V_1, V_2 \in \text{Coz}(X)$ ,  $f_1^* \in C^*(V_1)$ , and  $f_2^* \in C^*(V_2)$ , such that  $f_1^*, f_2^*$  are extensions of  $\frac{f_1}{|f_1|+|f_2|}, \frac{f_2}{|f_1|+|f_2|}$ , respectively. Consider  $V = V_1 \cap V_2$ , and let  $\sigma_1 \in C^*(X)$  be such that  $V = \text{coz}(\sigma_1)$ . Let  $\sigma = |\sigma_1| + (|f_1| + |f_2|)^{\frac{1}{2}}$ . Then  $\text{coz}(\sigma) = V$ . Define

$$h = \begin{cases} \frac{|f_1|+|f_2|}{\sigma} & \text{on } V, \\ 0 & \text{on } X - V. \end{cases}$$

For continuity of  $h$ , it is sufficient to show that it is continuous on  $\partial V$ . Let  $x_0 \in \partial V$ , and  $\epsilon > 0$ . Then  $h(x_0) = 0$ , and there is a neighborhood  $U_0$  of  $x_0$ , such that for each  $x \in U_0$ ,  $(|f_1| + |f_2|)^{\frac{1}{2}}(x) < (|f_1| + |f_2|)^{\frac{1}{2}}(x_0) + \epsilon = \epsilon$ . In fact, for each  $x \in U_0 \cap V \cap U$ ,

$$|h(x)| = \frac{|f_1| + |f_2|}{\sigma}(x) < \frac{|f_1| + |f_2|}{(|f_1| + |f_2|)^{\frac{1}{2}}}(x) = (|f_1| + |f_2|)^{\frac{1}{2}}(x) < \epsilon.$$

So,  $h \in C(X)$ .

Now let

$$\alpha = \begin{cases} f_1^* \sigma & \text{on } V, \\ 0 & \text{on } X - V, \end{cases} \quad \beta = \begin{cases} f_2^* \sigma & \text{on } V, \\ 0 & \text{on } X - V. \end{cases}$$

Then  $\alpha, \beta \in C(X)$ . By letting  $\gamma = \sigma - |\alpha| - |\beta|$ , we get  $X = \text{supp}(\sigma) = \text{supp}(\gamma + |\alpha| + |\beta|) \subseteq \text{supp}(|\gamma| + |\alpha| + |\beta|) = \text{supp}(\alpha) \cup \text{supp}(\beta) \cup \text{supp}(\gamma) \subseteq X$ .

Since  $f_1 = \alpha h$ ,  $f_2 = \beta h$ , and  $0 = \gamma h$ , we conclude that  $C(X)$  is an EM-ring.  $\square$

**Corollary 3.12.** *For a Tychonoff space  $X$ , the following statements are equivalent:*

- (1)  $C(X)$  is an EM-ring;
- (2)  $C^*(X)$  is an EM-ring;
- (3)  $X$  is an EM-space;
- (4)  $\beta X$  is an EM-space;
- (5) For each  $f, g \in C(X)$ , there exist  $h, \alpha, \beta, \gamma \in C(X)$  such that  $f = h\alpha, g = h\beta, 0 = h\gamma$  and  $\text{Ann}(\alpha, \beta, \gamma) = 0$ ;
- (6) For each  $U \in \text{Coz}(X)$ , and each  $g \in C^*(U)$  there is  $V \in \text{Coz}(X)$  such that  $U \subseteq V, \bar{V} = X$ , and  $g$  is continuously extendable on  $V$ .

Someone may wonder whether being an EM-space is preserved by continuous functions. Actually this is not true. The following is an example of a continuous function that maps an EM-space to a space that is not an EM-space.

**Example 3.13.** Let  $W^*$  be the space of ordinals that are less than or equal to  $\omega_1$ , the first uncountable ordinal. It was proved in [12] that every ordinal space is cozero complemented, and it is shown in Theorem 4.1 in this paper that a cozero complemented space is an EM-space.

Let  $S$  be the subspace of  $W^*$  that results after removing all limit ordinals, and let  $Y = S \cup \{\infty\}$  be the one point compactification of  $S$ . It is shown in Example 3.7 that  $Y$  is not an EM-space.

Consider the function  $f : W^* \rightarrow Y$  defined by

$$f(x) = \begin{cases} x & x \text{ is a non-limit ordinal,} \\ \infty & x \text{ is a limit ordinal.} \end{cases}$$

Let  $V$  be open in  $Y$ . If  $\infty \notin V$ , then  $f^{-1}(V) = V$ , and  $V$  is a union of isolated points thus open. If  $\infty \in V$ , then  $V$  is cofinite, and again  $f^{-1}(V)$  is open. So,  $f$  is continuous.

#### 4. Relation with other spaces

In this section, we relate EM-spaces to some other well known spaces, and show that EM-spaces include a wide class of spaces.

Concerning the relation between EM-rings and generalized morphic rings, which was the motivation to start this article, using the result in [6] that any countably generated  $z$ -ideal is generated by an idempotent, we get that for a Tychonoff space  $X$ , the ring  $C(X)$  is generalized morphic if and only if  $X$  is basically disconnected if and only if  $C(X)$  is a PP-ring. For more equivalent conditions, see [14]. Note that this result is not necessarily true outside  $C(X)$ , since there are commutative rings that are generalized morphic but not PP-rings. Thus, if  $C(X)$  is a generalized morphic ring, then it is an EM-ring, since any PP-ring is an EM-ring, see [2]. It is still an open question to characterize the relation between EM-rings and generalized morphic rings outside Noetherian rings and  $C(X)$ .

It was shown in Example 3.5 that any F-space is an EM-space, and now using Corollary 3.12, we get an even clearer proof. Using the technique in Example 3.6, one deduce that any metric space is an EM-space, or more generally, if

for each  $f \in C(X)$ , there exists  $g \in C(X)$  with  $\text{supp}(f) = Z(g)$ , then  $X$  is an EM-space.

**Theorem 4.1.** *A cozero complemented space is an EM-space.*

*Proof.* Let  $X$  be a cozero complemented space. Let  $U \in \text{Coz}(X)$ . Then there exists  $U' \in \text{Coz}(X)$  such that  $U \cap U' = \emptyset$  and  $\overline{U} \cup \overline{U'} = X$ . Let  $g \in C^*(U)$ .

Then  $g^* = \begin{cases} g & \text{on } U \\ 1 & \text{on } U' \end{cases} \in C^*(U \cup U')$  is the desired extension.  $\square$

It is well known that a Tychonoff space  $X$  is basically disconnected if and only if  $X$  is a cozero complemented F-space. So, the set  $\mathbb{R}$  of real numbers with Euclidean topology is an EM-space which is not an F-space, while  $\beta\mathbb{N} - \mathbb{N}$  is an EM-space that is not cozero complemented, since both spaces are not basically disconnected. One may ask if there is an EM-space that is not an F-space nor cozero complemented! The answer is yes; let  $X_1$  be a connected F-space, and  $X_2$  be a connected cozero complemented space. Then the free union space  $X = X_1 + X_2$  is an EM-space that is neither an F-space nor a cozero complemented space.

Recalling that a space  $X$  is a quasi F-space if every dense cozero set of  $X$  is  $C^*$ -embedded, it is directly deduced that a space  $X$  is an F-space if and only if it is an EM-space and a quasi F-space. Quasi F-spaces have been studied in number of articles including [11].

Recall that a space  $X$  is called an almost P-space if every  $G_\delta$ -set has dense interior, see [13]. An almost P-space which is an EM-space is an F-space, since an almost P-space is a quasi F-space.

Now we give an extra condition on an EM-space to get a cozero complemented space.

**Theorem 4.2.** *A locally connected EM-space is cozero complemented.*

*Proof.* Let  $X$  be an EM-space that is locally connected,  $U \in \text{Coz}(X)$ ,  $f \in C^*(X)$  such that  $f \geq 0$ , and  $U = \text{coz}(f)$ . Consider  $g = \cos\left(\frac{1}{f}\right) + 2 \in C^*(U)$ . Then, by Theorem 3.11, there exists a dense cozero set  $V$  that contains  $U$ , and  $g$  is continuously extendable on  $V$ . We need to show that  $\partial U \cap V = \emptyset$ .

Let  $b \in \partial U$ , and let  $W$  be any neighborhood of  $b$ . Since  $X$  is locally connected,  $W$  can be considered to be connected. Then for some  $a > 0$ ,  $f(W \cap U) = (0, a)$ . Moreover, for some  $n \in \mathbb{N}$ ,  $\frac{1}{2n\pi} < a$ . Therefore,

$$\left[ \frac{1}{(2n+1)\pi}, \frac{1}{2n\pi} \right] \subseteq f(W \cap U).$$

So,

$$(***) \quad [1, 3] \subseteq g(W \cap U).$$

If  $b \in V$ , then  $g$  has a continuous extension at  $b$ , and this contradicts (\*\*\*). This implies that  $\partial U \cap V = \emptyset$ .


Assume that  $l \in C^*(X)$ , with  $V = \text{coz}(l)$ . Then  $l(\partial U) = 0$ . Furthermore,  
 $l^* = \begin{cases} 0 & \text{on } \overline{U} \\ l & \text{elsewhere} \end{cases} \in C^*(X)$ .

Eventually,  $U \cap \text{coz}(l^*) = \emptyset$ , and  $\overline{U \cup \text{coz}(l^*)} = \overline{V} = X$ .  $\square$

**Corollary 4.3.** *A locally connected space is:*

- (1) *an EM-space if and only if it is cozero complemented.*
- (2) *an F-space if and only if it is basically disconnected.*

The following diagram illustrates the relations just obtained, where all the implications are strict, see also [10].



**Acknowledgment.** The authors are very grateful for the referee's efforts and valuable comments, which improved the article and brought it this way.

### References

- [1] E. Abuosba and O. Alkam, *When zero-divisor graphs are divisor graphs*, Turkish J. Math. **41** (2017), no. 4, 797–807. <https://doi.org/10.3906/mat-1601-102>
- [2] E. Abuosba and M. Ghanem, *Annihilating content in polynomial and power series rings*, J. Korean Math. Soc. **56** (2019), no. 5, 1403–1418. <https://doi.org/10.4134/JKMS.j180698>
- [3] D. D. Anderson, E. Abuosba, and M. Ghanem, *Annihilating content polynomials and EM-rings*, To appear in J. Algebra and its Application; <https://doi.org/10.1142/S021949882250092X>.
- [4] F. Azarpanah, E. Ghashghaei, and M. Ghoulipour, *C(X): something old and something new*, Comm. Algebra **49** (2021), no. 1, 185–206. <https://doi.org/10.1080/00927872.2020.1797070>
- [5] J. G. Brookshear, *Projective ideals in rings of continuous functions*, Pacific J. Math. **71** (1977), no. 2, 313–333. <http://projecteuclid.org/euclid.pjm/1102811430>
- [6] G. De Marco, *On the countably generated z-ideals of C(X)*, Proc. Amer. Math. Soc. **31** (1972), 574–576. <https://doi.org/10.2307/2037574>
- [7] M. Ghanem and E. Abu Osba, *Some extensions of generalized morhic rings and EM-rings*, An. Ştiinţ. Univ. “Ovidius” Constanţa Ser. Mat. **26** (2018), no. 1, 111–123. <https://doi.org/10.2478/auom-2018-0007>
- [8] L. Gillman and M. Henriksen, *Rings of continuous functions in which every finitely generated ideal is principal*, Trans. Amer. Math. Soc. **82** (1956), 366–391. <https://doi.org/10.2307/1993054>
- [9] L. Gillman and M. Jerison, *Rings of Continuous Functions*, reprint of the 1960 edition, Springer-Verlag, New York, 1976.
- [10] M. Henriksen and M. Jerison, *The space of minimal prime ideals of a commutative ring*, Trans. Amer. Math. Soc. **115** (1965), 110–130. <https://doi.org/10.2307/1994260>
- [11] M. Henriksen, J. Vermeer, and R. G. Woods, *Quasi F-covers of Tychonoff spaces*, Trans. Amer. Math. Soc. **303** (1987), no. 2, 779–803. <https://doi.org/10.2307/2000697>

- [12] M. Henriksen and R. G. Woods, *Cozero complemented spaces; when the space of minimal prime ideals of a  $C(X)$  is compact*, Top. Appl. **141** (2004), 147–170.
- [13] R. Levy, *Almost- $P$ -spaces*, Canadian J. Math. **29** (1977), no. 2, 284–288. <https://doi.org/10.4153/CJM-1977-030-7>
- [14] C. W. Neville, *When is  $C(X)$  a coherent ring?*, Proc. Amer. Math. Soc. **110** (1990), no. 2, 505–508. <https://doi.org/10.2307/2048096>

EMAD ABUOSBA  
DEPARTMENT OF MATHEMATICS  
THE UNIVERSITY OF JORDAN  
AMMAN, JORDAN  
*Email address:* eabuosba@ju.edu.jo

ISAAF ATASSI  
DEPARTMENT OF MATHEMATICS  
THE UNIVERSITY OF JORDAN  
AMMAN, JORDAN  
*Email address:* isaaf.atassi@gmail.com