

CERTAIN REDUCTION AND TRANSFORMATION FORMULAS FOR THE KAMPÉ DE FÉRIET FUNCTION

MEDHAT A. RAKHA AND ARJUN K. RATHIE

ABSTRACT. In 2014, Liu and Wang established a large number of interesting reduction, transformation and summation formulas for the Kampé de Fériet function. Inspired by the work, we aim to find further several transformation and reduction formulas for the Kampé de Fériet function. These formulas are mainly based on the formulas given by Liu and Wang [33].

1. Introduction

The familiar and classical Pochhammer symbol $(a)_\nu$ is defined as follows [35]:

$$(1.1) \quad (a)_\nu = \begin{cases} a(a+1)\cdots(a+n-1), & \nu = n \in \mathbb{N}, \\ 1, & \nu = 0, \end{cases} \\ = \frac{\Gamma(a+\nu)}{\Gamma(a)},$$

and $\Gamma(a)$ is the well-known Gamma function whose Euler's integral is defined by [44]:

$$(1.2) \quad \Gamma(z) = \int_0^\infty e^{-t} t^{z-1} dt, \quad (\operatorname{Re}(z) > 0).$$

In terms of Pochhammer's symbol, we define the generalized hypergeometric function ${}_pF_q$ for complex parameters and argument as follows [1, 2, 5, 43, 44, 46] by

$$(1.3) \quad {}_pF_q \left[\begin{matrix} a_1, & \dots, & a_p \\ b_1, & \dots, & b_q \end{matrix} ; x \right] = \sum_{k=0}^{\infty} \frac{(a_1)_k \cdots (a_p)_k}{(b_1)_k \cdots (b_q)_k} \frac{x^k}{k!}$$

Received March 11, 2021; Revised April 17, 2021; Accepted May 6, 2021.

2010 *Mathematics Subject Classification*. Primary 33B15, 33C05, 33C15; Secondary 33C65, 33C20.

Key words and phrases. Generalized hypergeometric function, Kampé de Fériet function, reduction, summation and transformation formulas.

when $p = q$ then this series converges for $|x| < \infty$, but when $q = p - 1$ convergence occurs when $|x| < 1$. However, when only one of the numerator parameters a_j is a negative integer or zero, then the series always converges since it is simply a polynomial in x of degree $-a_j$. For detailed account of ${}_pF_q$ including its properties, we refer standard text [1, 2, 5, 43, 44, 46].

It is interesting to mention here that for the hypergeometric function ${}_2F_1$, we have, four classical summation theorems namely Gauss, Gauss second, Kummer and Bailey and for generalized hypergeometric function ${}_3F_2$, we have four classical summation theorems namely Watson, Dixon, Whipple and Saalschütz. These theorems have wide applications. During 1992-2011, the above mentioned classical summation theorems have been generalized in the most general form by several researchers. For this, we refer some of the interesting research papers by Rakha and Rathie [37], Kim, et al. [27] and Lavoie, et al. [30–32].

It is not out of place to mention here that the vast popularity and immense usefulness of the hypergeometric function ${}_2F_1$, generalized hypergeometric function ${}_pF_q$ and confluent hypergeometric function ${}_1F_1$ in one variable have inspired and stimulated a number of researchers to discover and study hypergeometric functions in two and more variables.

A serious and very significant study of the function in two variables was initiated by Appell [4] who discovered the so-called four functions namely F_1, F_2, F_3 and F_4 which, in literature, popularly known as the Appell's functions which are natural generalization of the Gaussian hypergeometric function. The confluent forms of the Appell's functions were studied by Humbert [22]. A complete literature of the above mentioned Appell functions and their confluent forms is available in the standard text of Erdélyi et al. [20] and also in [3, 4]. Later on, the four Appell's functions and their confluent forms were generalized by Kampé de Fériet [24], who introduced a more general function in two variables. The notation defined by Kampé de Fériet for his double hypergeometric functions was subsequently abbreviated by Burchanall and Chaundy [6, 7]. In this paper, however, we shall recall the definition of a more general double hypergeometric function (than the one defined by Kampé de Fériet) in a slightly modified notation suggested by Srivastava and Panda [47].

So, for this, let (h_H) denote the sequence of H parameters (h_1, h_2, \dots, h_H) and for $n \in \mathbb{N}_0$, define the Pochhammer symbol

$$((h_H))_n := (h_1)_n \cdots (h_H)_n,$$

with the convention that when $n = 0$, the product is interpreted as unity. With this, we define the generalized Kampé de Fériet function in the following manner:

$$(1.4) \quad F_{G:C:D}^{H:A:B} \left[\begin{array}{l} (h_H) : (a_A); (b_B) \\ (g_G) : (c_C); (d_D) \end{array} ; x, y \right]$$

$$= \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{((h_H))_{m+n} ((a_A))_m ((b_B))_n x^m y^n}{((g_G))_{m+n} ((c_C))_m ((d_D))_n m! n!}.$$

For more details about this function (including its convergence conditions), we refer research paper [45].

For the Kampé de Fériet function, there are lots of transformation formulas available in the literature. Early works can be found in the following research papers [8, 9, 19, 21, 23, 25, 29, 34, 38–40, 42, 49, 50]

Moreover, researchers [10–18, 26, 33, 36, 41, 48] made further research and obtained many interesting and useful results in recent years.

In addition, the well known Beta function $B(a, b)$ is defined by the first integral and known to be evaluated as the second one in the following manner:

$$(1.5) \quad B(a, b) = \begin{cases} \int_0^1 t^{a-1} (1-t)^{b-1} dt, & ; (\operatorname{Re}(a) > 0, \operatorname{Re}(b) > 0) \\ \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}, & ; (a, b \in \mathbb{C} \setminus \mathbb{Z}_0^-). \end{cases}$$

Krattenthaler and Rao [28] made a systematic use of so-called Beta integral method, a method of establishing new hypergeometric identities in one and two variables from the old ones by mainly using the Beta integral in (1.5) based on the MATHEMATICA package HYP, to illustrate several interesting identities for the hypergeometric function and the Kampé de Fériet functions of two variables.

The paper is organized as follows. In Section 3, we establish twenty one reduction formulas for the Kampé de Fériet function. In Section 4, we establish twenty one transformation formulas and finally in Section 5, we further obtain twenty reduction formulas with the help of the results given in Sections 3 and 4.

The results established in this paper are simple, interesting, easily established and may be potentially useful. It hoped that these results will be a definite contribution in the theory of hypergeometric function in two variables.

2. Results required

In our present investigation, we shall require the following results due to Liu and Wang [33].

$$(2.1) \quad F_{1:0;0}^{1:1;1} \left[\begin{matrix} \alpha : \epsilon; & \beta - \epsilon \\ & & ; x, x \end{matrix} \right] = (1-x)^{-\alpha}.$$

$$(2.2) \quad F_{1:0;1}^{1:1;2} \left[\begin{matrix} \alpha : \epsilon; & \beta - \epsilon, \gamma \\ & & ; x, x \\ \beta : -; & \gamma + \beta \end{matrix} \right] \\ = (1-x)^{\beta-\alpha-\epsilon} {}_2F_1 \left[\begin{matrix} \beta - \epsilon, & \gamma + \beta - \alpha \\ & \gamma + \beta \end{matrix} ; x \right].$$

$$(2.3) \quad F_{1:0;1}^{1:1;2} \left[\begin{array}{c} \alpha : \epsilon; \quad \beta - \epsilon, 1 + \frac{1}{2}\alpha \\ \beta : -; \quad \frac{1}{2}\alpha \end{array} ; x, x \right] \\ = (1-x)^{\beta-\alpha-\epsilon} {}_2F_1 \left[\begin{array}{c} \beta - \epsilon, \quad 1 + \frac{1}{2}\beta \\ \frac{1}{2}\beta \end{array} ; x \right].$$

$$(2.4) \quad F_{1:0;1}^{1:1;2} \left[\begin{array}{c} \alpha : \epsilon; \quad \beta - \epsilon, \frac{1}{2}(\alpha - \beta) \\ \beta : -; \quad 1 + \frac{1}{2}(\alpha + \beta) \end{array} ; x, x \right] \\ = (1-x)^{\beta-\alpha-\epsilon} {}_3F_2 \left[\begin{array}{c} \beta - \epsilon, \quad 1 + \frac{1}{2}\beta, \quad \frac{1}{2}(\beta - \alpha) \\ \frac{1}{2}\beta, \quad 1 + \frac{1}{2}(\alpha + \beta) \end{array} ; x \right].$$

$$(2.5) \quad F_{1:0;2}^{1:1;3} \left[\begin{array}{c} \alpha : \epsilon; \quad \beta - \epsilon, 1 + \frac{1}{2}\alpha, \frac{1}{2}(\alpha - \beta) \\ \beta : -; \quad \frac{1}{2}\alpha, 1 + \frac{1}{2}(\alpha + \beta) \end{array} ; x, x \right] \\ = (1-x)^{\beta-\alpha-\epsilon} {}_2F_1 \left[\begin{array}{c} \beta - \epsilon, \quad \frac{1}{2}(\beta - \alpha) \\ 1 + \frac{1}{2}(\alpha + \beta) \end{array} ; x \right].$$

$$(2.6) \quad F_{1:0;2}^{1:1;3} \left[\begin{array}{c} \alpha : \epsilon; \quad \beta - \epsilon, 1 + \frac{1}{2}\alpha, \frac{1}{2}(\alpha - \beta - 1) \\ \beta : -; \quad \frac{1}{2}\alpha, \frac{1}{2}(\alpha + \beta + 3) \end{array} ; x, x \right] \\ = (1-x)^{\beta-\alpha-\epsilon} {}_3F_2 \left[\begin{array}{c} \beta - \epsilon, \quad 1 + \frac{1}{2}\beta, \quad \frac{1}{2}(\beta - \alpha - 1) \\ \frac{1}{2}\beta, \quad \frac{1}{2}(\alpha + \beta + 3) \end{array} ; x \right].$$

$$(2.7) \quad F_{1:0;1}^{0:2;3} \left[\begin{array}{c} - : \alpha, \epsilon; \quad \beta - \alpha, \beta - \epsilon, \gamma \\ \beta : -; \quad \delta \end{array} ; x, -\frac{x}{1-x} \right] \\ = (1-x)^{\beta-\alpha-\epsilon} {}_3F_2 \left[\begin{array}{c} \beta - \alpha, \quad \beta - \epsilon, \quad \delta - \gamma \\ \beta, \quad \delta \end{array} ; x \right].$$

$$(2.8) \quad F_{1:0;2}^{0:2;4} \left[\begin{array}{c} - : \alpha, \epsilon; \quad \beta - \alpha, \beta - \epsilon, \gamma, 1 + \frac{1}{2}\gamma \\ \beta : -; \quad \delta, \frac{1}{2}\gamma \end{array} ; x, -\frac{x}{1-x} \right] \\ = (1-x)^{\beta-\alpha-\epsilon} {}_4F_3 \left[\begin{array}{c} \beta - \alpha, \quad \beta - \epsilon, \quad \delta - \gamma - 1, \quad \gamma - \delta + 2 \\ \beta, \quad \delta, \quad \gamma - \delta + 1 \end{array} ; x \right].$$

$$(2.9) \quad F_{1:0;0}^{0:2;2} \left[\begin{array}{l} - : \alpha, \epsilon; \beta - \epsilon, \gamma \\ \beta : -; - \end{array} ; x, -\frac{x}{1-x} \right] \\ = (1-x)^{-\alpha} {}_2F_1 \left[\begin{array}{l} \beta - \epsilon, \alpha + \gamma \\ \beta \end{array} ; -\frac{x}{1-x} \right].$$

$$(2.10) \quad F_{1:0;1}^{0:2;3} \left[\begin{array}{l} - : \alpha, \epsilon; \beta - \epsilon, \gamma, \delta \\ \beta : -; \alpha + \gamma + \delta \end{array} ; x, -\frac{x}{1-x} \right] \\ = (1-x)^{-\alpha} {}_3F_2 \left[\begin{array}{l} \beta - \epsilon, \alpha + \gamma, \alpha + \delta \\ \beta, \alpha + \gamma + \delta \end{array} ; -\frac{x}{1-x} \right].$$

$$(2.11) \quad F_{1:0;1}^{0:2;3} \left[\begin{array}{l} - : \alpha, \epsilon; \beta - \epsilon, \gamma, 1 + \frac{1}{2}\gamma \\ \beta : -; \frac{1}{2}\gamma \end{array} ; x, -\frac{x}{1-x} \right] \\ = (1-x)^{-\alpha} {}_3F_2 \left[\begin{array}{l} \beta - \epsilon, \alpha + \gamma, 1 + \frac{1}{2}(\alpha + \gamma) \\ \beta, \frac{1}{2}(\alpha + \gamma) \end{array} ; -\frac{x}{1-x} \right].$$

$$(2.12) \quad F_{1:0;1}^{0:2;3} \left[\begin{array}{l} - : \alpha, \epsilon; \beta - \epsilon, \gamma, -\frac{1}{2}\alpha \\ \beta : -; 1 + \frac{1}{2}\alpha + \gamma \end{array} ; x, -\frac{x}{1-x} \right] \\ = (1-x)^{-\alpha} {}_4F_3 \left[\begin{array}{l} \alpha + \gamma, 1 + \frac{1}{2}(\alpha + \gamma), \frac{1}{2}\gamma, \beta - \epsilon \\ \frac{1}{2}(\alpha + \gamma), 1 + \frac{1}{2}\alpha + \gamma, \beta \end{array} ; -\frac{x}{1-x} \right].$$

$$(2.13) \quad F_{1:0;2}^{0:2;4} \left[\begin{array}{l} - : \alpha, \epsilon; \beta - \epsilon, \gamma, 1 + \frac{1}{2}\gamma, -\frac{1}{2}\alpha \\ \beta : -; \frac{1}{2}\gamma, 1 + \frac{1}{2}\alpha + \gamma \end{array} ; x, -\frac{x}{1-x} \right] \\ = (1-x)^{-\alpha} {}_3F_2 \left[\begin{array}{l} \beta - \epsilon, \alpha + \gamma, \frac{1}{2}\alpha \\ \beta, 1 + \frac{1}{2}\alpha + \gamma \end{array} ; -\frac{x}{1-x} \right].$$

$$(2.14) \quad F_{1:0;2}^{0:2;4} \left[\begin{array}{l} - : \alpha, \epsilon; \beta - \epsilon, \gamma, 1 + \frac{1}{2}\gamma, -\frac{\alpha+1}{2} \\ \beta : -; \frac{1}{2}\gamma, \frac{3}{2} + \frac{1}{2}\alpha + \gamma \end{array} ; x, -\frac{x}{1-x} \right] \\ = (1-x)^{-\alpha} {}_4F_3 \left[\begin{array}{l} \alpha + \gamma, 1 + \frac{1}{2}(\alpha + \gamma), \frac{1}{2}(\alpha - 1), \beta - \epsilon \\ \frac{1}{2}(\alpha + \gamma), \frac{3}{2} + \frac{1}{2}\alpha + \gamma, \beta \end{array} ; -\frac{x}{1-x} \right].$$

$$(2.15) \quad F_{1:0;1}^{2:0;1} \left[\begin{array}{c} \alpha, \gamma : -; \quad \epsilon \\ \beta : -; \quad \beta + \epsilon \end{array} ; x, -x \right] \\ = (1-x)^{-\alpha} {}_2F_1 \left[\begin{array}{c} \alpha, \quad \beta + \epsilon - \gamma \\ \beta + \epsilon \end{array} ; -\frac{x}{1-x} \right].$$

$$(2.16) \quad F_{1:0;1}^{2:0;1} \left[\begin{array}{c} \alpha, \gamma : -; \quad 1 + \frac{1}{2}\gamma \\ - : -; \quad \frac{1}{2}\gamma \end{array} ; x, -x \right] \\ = (1-x)^{-\alpha} {}_2F_1 \left[\begin{array}{c} \alpha, \quad 1 + \frac{1}{2}\beta \\ \frac{1}{2}\beta \end{array} ; -\frac{x}{1-x} \right].$$

$$(2.17) \quad F_{1:0;1}^{2:0;1} \left[\begin{array}{c} \alpha, \gamma : -; \quad \frac{1}{2}(\gamma - \beta) \\ \beta : -; \quad 1 + \frac{1}{2}(\gamma + \beta) \end{array} ; x, -x \right] \\ = (1-x)^{-\alpha} {}_3F_2 \left[\begin{array}{c} \alpha, \quad 1 + \frac{1}{2}\beta, \quad \frac{1}{2}(\beta - \gamma) \\ \frac{1}{2}\beta, \quad 1 + \frac{1}{2}(\gamma + \beta) \end{array} ; -\frac{x}{1-x} \right].$$

$$(2.18) \quad F_{1:0;2}^{2:0;2} \left[\begin{array}{c} \alpha, \gamma : -; \quad 1 + \frac{1}{2}\gamma, \frac{1}{2}(\gamma - \beta) \\ \beta : -; \quad \frac{1}{2}\gamma, 1 + \frac{1}{2}(\gamma + \beta) \end{array} ; x, -x \right] \\ = (1-x)^{-\alpha} {}_2F_1 \left[\begin{array}{c} \alpha, \quad \frac{1}{2}(\beta - \gamma) \\ 1 + \frac{1}{2}(\gamma + \beta) \end{array} ; -\frac{x}{1-x} \right].$$

$$(2.19) \quad F_{1:0;2}^{2:0;2} \left[\begin{array}{c} \alpha, \gamma : -; \quad 1 + \frac{1}{2}\gamma, \frac{1}{2}(\gamma - \beta - 1) \\ \beta : -; \quad \frac{1}{2}\gamma, \frac{1}{2}(3 + \gamma + \beta) \end{array} ; x, -x \right] \\ = (1-x)^{-\alpha} {}_3F_2 \left[\begin{array}{c} \alpha, \quad 1 + \frac{1}{2}\beta, \quad \frac{1}{2}(\beta - \gamma - 1) \\ \frac{1}{2}\beta, \quad \frac{1}{2}(3 + \gamma + \beta) \end{array} ; -\frac{x}{1-x} \right].$$

$$(2.20) \quad F_{1:0;1}^{1:1;2} \left[\begin{array}{c} \alpha : \epsilon; \quad \beta - \epsilon, \gamma \\ \beta : -; \quad \delta \end{array} ; x, x \right] \\ = (1-x)^{-\alpha} {}_3F_2 \left[\begin{array}{c} \alpha, \quad \beta - \epsilon, \quad \delta - \gamma \\ \beta, \quad \delta \end{array} ; -\frac{x}{1-x} \right].$$

$$(2.21) \quad F_{1:0;2}^{1:1;3} \left[\begin{array}{c} \alpha : \epsilon; \beta - \epsilon, \gamma, 1 + \frac{1}{2}\gamma \\ \beta : -; \delta, \frac{1}{2}\gamma \end{array} ; x, x \right] \\ = (1-x)^{-\alpha} {}_4F_3 \left[\begin{array}{c} \alpha, \beta - \epsilon, \delta - \gamma - 1, \delta - \gamma + 2 \\ \beta, \delta, \gamma - \delta + 1 \end{array} ; -\frac{x}{1-x} \right].$$

3. Reduction formulas for the Kampé de Fériet function

In this section, the following twenty one interesting reduction formulas have been established.

$$(3.1) \quad F_{2:0;0}^{2:1;1} \left[\begin{array}{c} \alpha, \lambda : \epsilon; \beta - \epsilon \\ \beta, \mu : -; - \end{array} ; 1, 1 \right] = \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \alpha)\Gamma(\mu - \lambda)}.$$

$$(3.2) \quad F_{2:0;1}^{2:1;2} \left[\begin{array}{c} \alpha, \lambda : \epsilon; \beta - \epsilon, \gamma \\ \beta, \mu : -; \gamma + \beta \end{array} ; 1, 1 \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu + \beta - \alpha - \epsilon - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu + \beta - \alpha - \epsilon)} {}_3F_2 \left[\begin{array}{c} \beta - \epsilon, \gamma + \beta - \alpha, \lambda \\ \gamma + \beta, \mu + \beta - \alpha - \epsilon \end{array} ; 1 \right].$$

$$(3.3) \quad F_{2:0;1}^{2:1;2} \left[\begin{array}{c} \alpha, \lambda : \epsilon; \beta - \epsilon, 1 + \frac{1}{2}\alpha \\ \beta, \mu : -; \frac{1}{2}\alpha \end{array} ; 1, 1 \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu + \beta - \alpha - \epsilon - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu + \beta - \alpha - \epsilon)} {}_3F_2 \left[\begin{array}{c} \beta - \epsilon, 1 + \frac{1}{2}\beta, \lambda \\ \frac{1}{2}\beta, \mu + \beta - \alpha - \epsilon \end{array} ; 1 \right].$$

$$(3.4) \quad F_{2:0;1}^{2:1;2} \left[\begin{array}{c} \alpha, \lambda : \epsilon; \beta - \epsilon, \frac{1}{2}(\alpha - \beta) \\ \beta, \mu : -; 1 + \frac{1}{2}(\alpha + \beta) \end{array} ; 1, 1 \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu + \beta - \alpha - \epsilon - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu + \beta - \alpha - \epsilon)} \\ \times {}_4F_3 \left[\begin{array}{c} \beta - \epsilon, 1 + \frac{1}{2}\beta, \frac{1}{2}(\beta - \alpha), \lambda \\ \frac{1}{2}\beta, 1 + \frac{1}{2}(\beta + \alpha), \mu + \beta - \alpha - \epsilon \end{array} ; 1 \right].$$

$$(3.5) \quad F_{2:0;2}^{2:1;3} \left[\begin{array}{c} \alpha, \lambda : \epsilon; \beta - \epsilon, 1 + \frac{1}{2}\alpha, \frac{1}{2}(\alpha - \beta) \\ \beta, \mu : -; \frac{1}{2}\alpha, 1 + \frac{1}{2}(\alpha + \beta) \end{array} ; 1, 1 \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu + \beta - \alpha - \epsilon - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu + \beta - \alpha - \epsilon)} {}_3F_2 \left[\begin{array}{c} \beta - \epsilon, \frac{1}{2}(\beta - \alpha), \lambda \\ 1 + \frac{1}{2}(\alpha + \beta), \mu + \beta - \alpha - \epsilon \end{array} ; 1 \right].$$

$$(3.6) \quad F_{2:0;2}^{2:1;3} \left[\begin{array}{c} \alpha, \lambda : \epsilon; \quad \beta - \epsilon, 1 + \frac{1}{2}\alpha, \frac{1}{2}(\alpha - \beta + 1) \\ \beta, \mu : -; \quad \frac{1}{2}\alpha, \frac{1}{2}(\alpha + \beta + 3) \end{array} ; 1, 1 \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu + \beta - \alpha - \epsilon - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu + \beta - \alpha - \epsilon)} \\ \times {}_4F_3 \left[\begin{array}{c} \beta - \epsilon, \quad 1 + \frac{1}{2}\beta, \quad \frac{1}{2}(\beta - \alpha + 1), \quad \lambda \\ \frac{1}{2}\beta, \quad \frac{1}{2}(\alpha + \beta + 3), \quad \mu + \beta - \alpha - \epsilon \end{array} ; 1 \right].$$

$$(3.7) \quad F_{1:1;2}^{1:2;3} \left[\begin{array}{c} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \beta - \alpha, \gamma \\ \beta : \mu; \quad \delta, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu + \beta - \alpha - \epsilon - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu + \beta - \alpha - \epsilon)} \\ \times {}_4F_3 \left[\begin{array}{c} \beta - \alpha, \quad \beta - \epsilon, \quad \delta - \gamma, \quad \lambda \\ \beta, \quad \delta, \quad \mu + \beta - \alpha - \epsilon \end{array} ; 1 \right].$$

$$(3.8) \quad F_{1:1;3}^{1:2;4} \left[\begin{array}{c} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \beta - \alpha, \gamma, 1 + \frac{1}{2}\gamma \\ \beta : \mu; \quad \delta, \frac{1}{2}\gamma, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu + \beta - \alpha - \epsilon - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu + \beta - \alpha - \epsilon)} \\ \times {}_5F_4 \left[\begin{array}{c} \beta - \alpha, \quad \beta - \epsilon, \quad \delta - \gamma - 1, \quad \gamma - \delta + 2, \quad \lambda \\ \beta, \quad \delta, \quad \gamma - \delta + 1, \quad \mu + \beta - \alpha - \epsilon \end{array} ; 1 \right].$$

$$(3.9) \quad F_{1:1;1}^{1:2;2} \left[\begin{array}{c} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \gamma \\ \beta : \mu; \quad 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} {}_3F_2 \left[\begin{array}{c} \beta - \epsilon, \quad \alpha + \gamma, \quad \lambda \\ \beta, \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right].$$

$$(3.10) \quad F_{1:1;2}^{1:2;3} \left[\begin{array}{c} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \gamma, \delta \\ \beta : \mu; \quad \alpha + \gamma + \delta, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\ = {}_4F_3 \left[\begin{array}{c} \beta - \epsilon, \quad \alpha + \gamma, \quad \alpha + \delta, \quad \lambda \\ \beta, \quad \alpha + \gamma + \delta, \quad 1 - \mu + \lambda \end{array} ; 1 \right].$$

$$\begin{aligned}
(3.11) \quad & F_{1:1;2}^{1:2;3} \left[\begin{array}{c} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \gamma, 1 + \frac{1}{2}\gamma \\ \beta : \mu; \quad \frac{1}{2}\gamma, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\
&= \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} \\
&\quad \times {}_4F_3 \left[\begin{array}{c} \beta - \epsilon, \quad \alpha + \gamma, \quad 1 + \frac{1}{2}(\alpha + \gamma), \quad \lambda \\ \beta, \quad \frac{1}{2}(\alpha + \gamma), \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right]. \\
(3.12) \quad & F_{1:1;2}^{1:2;3} \left[\begin{array}{c} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \gamma, -\frac{1}{2}\gamma \\ \beta : \mu; \quad 1 + \frac{1}{2}\alpha + \gamma, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\
&= \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} \\
&\quad \times {}_5F_4 \left[\begin{array}{c} \alpha + \gamma, \quad 1 + \frac{1}{2}(\alpha + \gamma), \quad \frac{1}{2}\alpha, \quad \beta - \epsilon, \quad \lambda \\ \frac{1}{2}(\alpha + \gamma), \quad 1 + \frac{1}{2}\alpha + \gamma, \quad \beta, \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right]. \\
(3.13) \quad & F_{1:1;3}^{1:2;4} \left[\begin{array}{c} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \gamma, 1 + \frac{1}{2}\gamma, -\frac{1}{2}\gamma \\ \beta : \beta; \quad \frac{1}{2}\gamma, 1 + \frac{1}{2}\alpha + \gamma, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\
&= \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} \\
&\quad \times {}_4F_3 \left[\begin{array}{c} \beta - \epsilon, \quad \alpha + \gamma, \quad \frac{1}{2}\alpha, \quad \lambda \\ \beta, \quad 1 + \frac{1}{2}\alpha + \gamma, \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right]. \\
(3.14) \quad & F_{1:1;3}^{1:2;4} \left[\begin{array}{c} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \gamma, 1 + \frac{1}{2}\gamma, -\frac{\alpha+1}{2} \\ \beta : \mu; \quad \frac{1}{2}\gamma, \frac{3}{2} + \frac{1}{2}\alpha + \gamma, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\
&= \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} \\
&\quad \times {}_5F_4 \left[\begin{array}{c} \alpha + \gamma, \quad 1 + \frac{1}{2}(\alpha + \gamma), \quad \frac{1}{2}(\alpha - 1), \quad \beta - \epsilon, \quad \lambda \\ \frac{1}{2}(\alpha + \gamma), \quad \frac{3}{2} + \frac{1}{2}\alpha + \gamma, \quad \beta, \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right]. \\
(3.15) \quad & F_{2:0;1}^{3:0;1} \left[\begin{array}{c} \alpha, \gamma, \lambda : -; \quad \epsilon \\ \beta, \mu : -; \quad \beta + \epsilon \end{array} ; 1, -1 \right] \\
&= \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} {}_3F_2 \left[\begin{array}{c} \alpha, \quad \beta + \epsilon - \gamma, \quad \lambda \\ \beta + \epsilon, \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right].
\end{aligned}$$

$$(3.16) \quad F_{2:0;1}^{3:0;1} \left[\begin{matrix} \alpha, \gamma, \lambda : -; & 1 + \frac{1}{2}\gamma & \\ & & ; 1, -1 \end{matrix} \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} {}_3F_2 \left[\begin{matrix} \alpha, & 1 + \frac{1}{2}\beta, & \lambda \\ \frac{1}{2}\beta, & 1 - \mu + \alpha + \lambda & \end{matrix} ; 1 \right].$$

$$(3.17) \quad F_{2:0;1}^{3:0;1} \left[\begin{matrix} \alpha, \gamma, \lambda : -; & \frac{1}{2}(\gamma - \beta) & \\ & & ; 1, -1 \end{matrix} \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} \\ \times {}_4F_3 \left[\begin{matrix} \alpha, & 1 + \frac{1}{2}\beta, & \frac{1}{2}(\beta - \gamma), & \lambda \\ \frac{1}{2}\beta, & 1 + \frac{1}{2}(\gamma + \beta) & 1 - \mu + \alpha + \lambda & \end{matrix} ; 1 \right].$$

$$(3.18) \quad F_{2:0;2}^{3:0;2} \left[\begin{matrix} \alpha, \gamma, \lambda : -; & 1 + \frac{1}{2}\gamma, \frac{1}{2}(\gamma - \beta - 1) & \\ & & ; 1, -1 \end{matrix} \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} {}_3F_2 \left[\begin{matrix} \alpha, & \frac{1}{2}(\beta - \gamma), & \lambda \\ 1 + \frac{1}{2}(\beta + \gamma), & 1 - \mu + \alpha + \lambda & \end{matrix} ; 1 \right].$$

$$(3.19) \quad F_{2:0;2}^{3:0;2} \left[\begin{matrix} \alpha, \gamma, \lambda : -; & 1 + \frac{1}{2}\gamma, \frac{1}{2}(\gamma - \beta - 1) & \\ & & ; 1, -1 \end{matrix} \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} \\ \times {}_4F_3 \left[\begin{matrix} \alpha, & 1 + \frac{1}{2}\beta, & \frac{1}{2}(\beta - \gamma - 1), & \lambda \\ \frac{1}{2}\beta, & \frac{1}{2}(3 + \gamma + \beta) & 1 - \mu + \alpha + \lambda & \end{matrix} ; 1 \right].$$

$$(3.20) \quad F_{2:0;1}^{2:1;2} \left[\begin{matrix} \alpha, \lambda : \epsilon; & \beta - \epsilon, \gamma & \\ & & ; 1, 1 \end{matrix} \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} \\ \times {}_4F_3 \left[\begin{matrix} \alpha, & \beta - \epsilon, & \delta - \gamma, & \lambda \\ \beta, & \delta & 1 - \mu + \alpha + \lambda & \end{matrix} ; 1 \right].$$

$$\begin{aligned}
(3.21) \quad & F_{2:0;2}^{2:1;3} \left[\begin{array}{c} \alpha, \lambda : \quad \epsilon; \quad \beta - \epsilon, \gamma, 1 + \frac{1}{2}\gamma \\ \beta, \mu : \quad -; \quad \delta, \frac{1}{2}\gamma \end{array} ; 1, 1 \right] \\
&= \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} \\
&\quad \times {}_5F_4 \left[\begin{array}{c} \alpha, \quad \beta - \epsilon, \quad \delta - \gamma - 1, \quad \delta - \gamma + 2 \quad \lambda \\ \beta, \quad \delta \quad \gamma - \delta + 1 \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right].
\end{aligned}$$

Proof. The derivation of the reduction formulas (3.1) to (3.21) are quite straight forward. We shall establish only one of them, say, for example, (3.2) and rest can be proven on a similar fashion.

In order to establish the result (3.2), we proceed as follows. Multiply both sides of the equation (2.2) by $x^{\lambda-1}(1-x)^{\mu-\lambda-1}$ and integrating with respect to x between the interval $[0, 1]$, we have

$$\begin{aligned}
(3.22) \quad & \int_0^1 x^{\lambda-1}(1-x)^{\mu-\lambda-1} F_{1:0;1}^{1:1;2} \left[\begin{array}{c} \alpha : \quad \epsilon; \quad \beta - \epsilon, \gamma \\ \beta : \quad -; \quad \gamma + \beta \end{array} ; x, x \right] dx \\
&= \int_0^1 x^{\lambda-1}(1-x)^{\mu-\lambda-\alpha-\epsilon+\beta-1} {}_2F_1 \left[\begin{array}{c} \beta - \epsilon, \quad \gamma + \beta - \alpha \\ \gamma + \beta \end{array} ; x \right] dx.
\end{aligned}$$

Now, let us denote the left-hand side of (3.22) by I_1 and expressing the Kampé de Fériet function by its definition, changing the order of integration and summation, which is easily seen to be justified due to the uniform convergence of the series involved in the process, we have

$$I_1 = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(\alpha)_{m+n}(\epsilon)_m(\beta - \epsilon)_n(\gamma)_n}{(\beta)_{m+n}(\gamma + \beta)_n m! n!} \int_0^1 x^{\lambda+m+n-1}(1-x)^{\mu-\lambda-1} dx.$$

Evaluating the beta integral, we have after some algebra

$$I_1 = \frac{\Gamma(\lambda)\Gamma(\mu - \lambda)}{\Gamma(\mu)} \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(\alpha)_{m+n}(\lambda)_{m+n}(\epsilon)_m(\beta - \epsilon)_n(\gamma)_n}{(\beta)_{m+n}(\mu)_{m+n}(\gamma + \beta)_n m! n!}.$$

Summing up the series with the help of the definition of Kampé de Fériet function, we have

$$(3.23) \quad I_1 = \frac{\Gamma(\lambda)\Gamma(\mu - \lambda)}{\Gamma(\mu)} F_{2:0;1}^{2:1;2} \left[\begin{array}{c} \alpha, \lambda \quad \epsilon; \quad \beta - \epsilon, \gamma \\ \beta, \mu : \quad -; \quad \gamma + \beta \end{array} ; 1, 1 \right].$$

Now, denoting the right-hand side of (3.22) by I_2 , expressing ${}_2F_1$ as a series and proceeding the same as above and after a little simplification, we have

$$I_2 = \frac{\Gamma(\lambda)\Gamma(\mu + \beta - \alpha - \epsilon - \lambda)}{\Gamma(\mu + \beta - \alpha - \epsilon)} \sum_{n=0}^{\infty} \frac{(\beta - \epsilon)_n(\gamma + \beta - \alpha)_n(\lambda)_n}{(\gamma + \beta)_n(\mu + \beta - \alpha - \epsilon)_n n!}.$$

Summing up the series, we have

$$(3.24) \quad I_2 = \frac{\Gamma(\lambda)\Gamma(\mu + \beta - \alpha - \epsilon - \lambda)}{\Gamma(\mu + \beta - \alpha - \epsilon)} {}_3F_2 \left[\begin{matrix} \beta - \epsilon, & \gamma + \beta - \alpha, & \lambda \\ \gamma + \beta, & \mu + \beta - \alpha - \epsilon \end{matrix} ; 1 \right].$$

Finally, substituting the values of (3.23) and (3.24) in (3.22), we arrive at the desired reduction formula (3.2). This completes the proof of (3.2). \square

4. Transformation formulas for the Kampé de Fériet function

In this section, we shall establish the following twenty interesting transformation formulas for the Kampé de Fériet function.

$$(4.1) \quad \begin{aligned} & F_{2;0;1}^{2;1;2} \left[\begin{matrix} \alpha, \lambda : \epsilon; \beta - \epsilon, \gamma \\ \beta, \mu : -; \gamma + \beta \end{matrix} ; 1, 1 \right] \\ &= F_{1;0;1}^{1;1;2} \left[\begin{matrix} \lambda : \alpha + \epsilon - \beta; \beta - \epsilon, \gamma + \beta - \alpha \\ \mu : -; \gamma + \beta \end{matrix} ; 1, 1 \right]. \end{aligned}$$

$$(4.2) \quad \begin{aligned} & F_{2;0;1}^{2;1;2} \left[\begin{matrix} \alpha, \lambda : \epsilon; \beta - \epsilon, 1 + \frac{1}{2}\alpha \\ \beta, \mu : -; \frac{1}{2}\alpha \end{matrix} ; 1, 1 \right] \\ &= F_{1;0;1}^{1;1;2} \left[\begin{matrix} \lambda : \alpha + \epsilon - \beta; \beta - \epsilon, 1 + \frac{1}{2}\beta \\ \mu : -; \frac{1}{2}\beta \end{matrix} ; 1, 1 \right]. \end{aligned}$$

$$(4.3) \quad \begin{aligned} & F_{2;0;1}^{2;1;2} \left[\begin{matrix} \alpha, \lambda : \epsilon; \beta - \epsilon, \frac{1}{2}(\alpha - \beta) \\ \beta, \mu : -; 1 + \frac{1}{2}(\alpha + \beta) \end{matrix} ; 1, 1 \right] \\ &= F_{1;0;2}^{1;1;3} \left[\begin{matrix} \lambda : \alpha + \epsilon - \beta; \beta - \epsilon, 1 + \frac{1}{2}\beta, \frac{1}{2}(\beta - \alpha) \\ \mu : -; \frac{1}{2}\beta, 1 + \frac{1}{2}(\alpha + \beta) \end{matrix} ; 1, 1 \right]. \end{aligned}$$

$$(4.4) \quad \begin{aligned} & F_{2;0;2}^{2;1;3} \left[\begin{matrix} \alpha, \lambda : \epsilon; \beta - \epsilon, 1 + \frac{1}{2}\alpha, \frac{1}{2}(\alpha - \beta) \\ \beta, \mu : -; \frac{1}{2}\alpha, 1 + \frac{1}{2}(\alpha + \beta) \end{matrix} ; 1, 1 \right] \\ &= F_{1;0;1}^{1;1;2} \left[\begin{matrix} \lambda : \alpha + \epsilon - \beta; \beta - \epsilon, \frac{1}{2}(\beta - \alpha) \\ \mu : -; 1 + \frac{1}{2}(\alpha + \beta) \end{matrix} ; 1, 1 \right]. \end{aligned}$$

$$(4.5) \quad F_{2;0;2}^{2;1;3} \left[\begin{matrix} \alpha, \lambda : \epsilon; \beta - \epsilon, 1 + \frac{1}{2}\alpha, \frac{1}{2}(\alpha - \beta + 1) \\ \beta, \mu : -; \frac{1}{2}\alpha, \frac{1}{2}(\alpha + \beta + 3) \end{matrix} ; 1, 1 \right]$$

$$= F_{1:0;2}^{1:1;3} \left[\begin{array}{l} \lambda : \alpha + \epsilon - \beta; \quad \beta - \epsilon, 1 + \frac{1}{2}\beta, \frac{1}{2}(\beta - \alpha - 1) \\ \mu : \quad -; \quad \frac{1}{2}\beta, \frac{1}{2}(\alpha + \beta + 3) \end{array} ; 1, 1 \right].$$

$$(4.6) \quad F_{1:1;2}^{1:2;3} \left[\begin{array}{l} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \beta - \alpha, \gamma \\ \beta : \quad \mu; \quad \delta, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\ = F_{1:0;2}^{1:1;3} \left[\begin{array}{l} \lambda : \alpha + \epsilon - \beta; \quad \beta - \alpha, \beta - \epsilon, \delta - \gamma \\ \mu : \quad -; \quad \beta, \delta \end{array} ; 1, 1 \right].$$

$$(4.7) \quad F_{1:1;3}^{1:2;4} \left[\begin{array}{l} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \beta - \alpha, \gamma, 1 + \frac{1}{2}\gamma \\ \beta : \quad \mu; \quad \delta, \frac{1}{2}\gamma, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\ = F_{1:0;3}^{1:1;4} \left[\begin{array}{l} \lambda : \alpha + \epsilon - \beta; \quad \beta - \alpha, \beta - \epsilon, \delta - \gamma - 1, \gamma - \delta + 2 \\ \mu : \quad -; \quad \beta, \delta, \gamma - \delta + 1 \end{array} ; 1, 1 \right].$$

$$(4.8) \quad F_{1:1;1}^{1:2;2} \left[\begin{array}{l} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \gamma \\ \beta : \quad \mu; \quad 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\ = F_{0:1;2}^{1:1;2} \left[\begin{array}{l} \lambda : \alpha; \quad \beta - \epsilon, \alpha + \gamma \\ - : \quad \mu; \quad \beta, 1 - \mu - \lambda \end{array} ; 1, 1 \right].$$

$$(4.9) \quad F_{1:1;2}^{1:2;3} \left[\begin{array}{l} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \gamma, \delta \\ \beta : \quad \mu; \quad \alpha + \gamma + \delta, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\ = F_{0:1;2}^{1:1;3} \left[\begin{array}{l} \lambda : \alpha; \quad \beta - \epsilon, \alpha + \gamma, \alpha + \delta \\ - : \quad \mu; \quad \beta, \alpha + \gamma + \delta, 1 - \mu + \lambda \end{array} ; 1, 1 \right].$$

$$(4.10) \quad F_{1:1;2}^{1:2;3} \left[\begin{array}{l} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \gamma, 1 + \frac{1}{2}\gamma \\ \beta : \quad \mu; \quad \frac{1}{2}\gamma, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\ = F_{0:1;3}^{1:1;3} \left[\begin{array}{l} \lambda : \alpha; \quad \beta - \epsilon, \alpha + \gamma, 1 + \frac{1}{2}(\alpha + \gamma) \\ - : \quad \mu; \quad \beta, \frac{1}{2}(\alpha + \gamma), 1 - \mu + \lambda \end{array} ; 1, 1 \right].$$

$$(4.11) \quad F_{1:1;2}^{1:2;3} \left[\begin{array}{l} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \gamma, -\frac{1}{2}\gamma \\ \beta : \quad \mu; \quad 1 + \frac{1}{2}\alpha + \gamma, 1 - \mu + \lambda \end{array} ; 1, 1 \right]$$

$$\begin{aligned}
&= F_{0:1;4}^{1:1;4} \left[\begin{array}{l} \lambda : \alpha; \quad \beta - \epsilon, \alpha + \gamma, 1 + \frac{1}{2}(\alpha + \gamma), \frac{1}{2}\alpha \\ - : \mu; \quad \beta, \frac{1}{2}(\alpha + \gamma), 1 + \frac{1}{2}\alpha + \gamma, 1 - \mu + \lambda \end{array} ; 1, 1 \right]. \\
(4.12) \quad &F_{1:1;3}^{1:2;4} \left[\begin{array}{l} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \gamma, 1 + \frac{1}{2}\gamma, -\frac{1}{2}\gamma \\ \beta : \beta; \quad \frac{1}{2}\gamma, 1 + \frac{1}{2}\alpha + \gamma, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\
&= F_{0:1;3}^{1:1;3} \left[\begin{array}{l} \lambda : \alpha; \quad \beta - \epsilon, \alpha + \gamma, \frac{1}{2}\alpha \\ - : \mu; \quad \beta, 1 + \frac{1}{2}\alpha + \gamma, 1 - \mu + \lambda \end{array} ; 1, 1 \right].
\end{aligned}$$

$$\begin{aligned}
(4.13) \quad &F_{1:1;3}^{1:2;4} \left[\begin{array}{l} \lambda : \alpha, \epsilon; \quad \beta - \epsilon, \gamma, 1 + \frac{1}{2}\gamma, -\frac{\alpha+1}{2} \\ \beta : \mu; \quad \frac{1}{2}\gamma, \frac{3}{2} + \frac{1}{2}\alpha + \gamma, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\
&= F_{0:1;4}^{1:1;4} \left[\begin{array}{l} \lambda : \alpha; \quad \beta - \epsilon, \alpha + \gamma, 1 + \frac{1}{2}(\alpha + \gamma), \frac{1}{2}(\alpha - 1) \\ - : \mu; \quad \beta, \frac{1}{2}(\alpha + \gamma), \frac{3}{2} + \frac{1}{2}\alpha + \gamma, 1 - \mu + \lambda \end{array} ; 1, 1 \right].
\end{aligned}$$

$$\begin{aligned}
(4.14) \quad &F_{2:0;1}^{3:0;1} \left[\begin{array}{l} \alpha, \gamma, \lambda : -; \quad \epsilon \\ \beta, \mu : -; \quad \beta + \epsilon \end{array} ; 1, -1 \right] \\
&= F_{0:1;2}^{1:1;2} \left[\begin{array}{l} \lambda : \alpha; \quad \alpha, \beta + \epsilon - \gamma \\ - : \mu; \quad \beta + \epsilon, \gamma - \mu + \lambda \end{array} ; 1, 1 \right].
\end{aligned}$$

$$\begin{aligned}
(4.15) \quad &F_{2:0;1}^{3:0;1} \left[\begin{array}{l} \alpha, \gamma, \lambda : -; \quad 1 + \frac{1}{2}\gamma \\ \beta, \mu : -; \quad \frac{1}{2}\gamma \end{array} ; 1, -1 \right] \\
&= F_{0:1;2}^{1:1;2} \left[\begin{array}{l} \lambda : \alpha; \quad \alpha, 1 + \frac{1}{2}\beta \\ - : \mu; \quad \frac{1}{2}\beta, 1 - \mu + \lambda \end{array} ; 1, 1 \right].
\end{aligned}$$

$$\begin{aligned}
(4.16) \quad &F_{2:0;1}^{3:0;1} \left[\begin{array}{l} \alpha, \gamma, \lambda : -; \quad \frac{1}{2}(\gamma - \beta) \\ \beta, \mu : -; \quad 1 + \frac{1}{2}(\gamma + \beta) \end{array} ; 1, -1 \right] \\
&= F_{0:1;3}^{1:1;3} \left[\begin{array}{l} \lambda : \alpha; \quad \alpha, 1 + \frac{1}{2}\beta, \frac{1}{2}(\beta - \gamma) \\ - : \mu; \quad \frac{1}{2}\beta, 1 + \frac{1}{2}(\gamma + \beta), 1 - \mu + \lambda \end{array} ; 1, 1 \right].
\end{aligned}$$

$$(4.17) \quad F_{2:0;2}^{3:0;2} \left[\begin{array}{l} \alpha, \gamma, \lambda : -; \quad 1 + \frac{1}{2}\gamma, \frac{1}{2}(\gamma - \beta - 1) \\ \beta, \mu : -; \quad \frac{1}{2}\gamma, \frac{1}{2}(3 + \gamma + \beta) \end{array} ; 1, -1 \right]$$

$$\begin{aligned}
&= F_{0:1;2}^{1:1;2} \left[\begin{array}{c} \lambda : \alpha; \quad \alpha, \frac{1}{2}(\beta - \gamma) \\ - : \mu; \quad 1 + \frac{1}{2}(\gamma + \beta), 1 - \mu + \lambda \end{array} ; 1, 1 \right]. \\
(4.18) \quad &F_{2:0;2}^{3:0;2} \left[\begin{array}{c} \alpha, \gamma, \lambda : -; \quad 1 + \frac{1}{2}\gamma, \frac{1}{2}(\gamma - \beta - 1) \\ \beta, \mu : -; \quad \frac{1}{2}\gamma, \frac{1}{2}(3 + \gamma + \beta) \end{array} ; 1, -1 \right] \\
&= F_{0:1;3}^{1:1;3} \left[\begin{array}{c} \lambda : \alpha; \quad \alpha, 1 + \frac{1}{2}\beta, \frac{1}{2}(\beta - \mu - 1) \\ - : \mu; \quad \frac{1}{2}\beta, 1 + \frac{1}{2}(2 + \gamma + \beta), 1 - \mu + \lambda \end{array} ; 1, 1 \right].
\end{aligned}$$

$$\begin{aligned}
(4.19) \quad &F_{2:0;1}^{2:1;2} \left[\begin{array}{c} \alpha, \lambda : \epsilon; \quad \beta - \epsilon, \gamma \\ \beta, \mu : -; \quad \delta \end{array} ; 1, 1 \right] \\
&= F_{0:1;3}^{1:1;3} \left[\begin{array}{c} \lambda : \alpha; \quad \alpha, \beta - \epsilon, \delta - \gamma \\ - : \mu; \quad \beta, \delta, 1 - \mu + \lambda \end{array} ; 1, 1 \right].
\end{aligned}$$

$$\begin{aligned}
(4.20) \quad &F_{2:0;2}^{2:1;3} \left[\begin{array}{c} \alpha, \lambda : \epsilon; \quad \beta - \epsilon, \gamma, 1 + \frac{1}{2}\gamma \\ \beta, \mu : -; \quad \delta, \frac{1}{2}\gamma \end{array} ; 1, 1 \right] \\
&= F_{0:1;4}^{1:1;4} \left[\begin{array}{c} \lambda : \alpha; \quad \alpha, \beta - \epsilon, \delta - \gamma - 1, \delta - \gamma + 2 \\ - : \mu; \quad \beta, \delta, \gamma - \delta + 1, 1 - \mu + \lambda \end{array} ; 1, 1 \right].
\end{aligned}$$

Proof. The derivation of the reduction formulas (4.1) to (4.20) are quite straight forward. We shall establish only one of them, say, for example, (4.1) and rest can be proven on a similar fashion.

In order to establish the result (4.1), we proceed as follows. In equation (2.2), if we replace $(1-x)^{\beta-\alpha-\epsilon}$ to ${}_1F_0 \left[\begin{array}{c} \alpha + \epsilon - \beta \\ - \end{array} ; x \right]$, then it takes the following form:

$$\begin{aligned}
(4.21) \quad &F_{1:0;1}^{1:1;2} \left[\begin{array}{c} \alpha : \epsilon; \quad \beta - \epsilon, \gamma \\ \beta : -; \quad \gamma + \beta \end{array} ; x, x \right] \\
&= {}_1F_0 \left[\begin{array}{c} \alpha + \epsilon - \beta \\ - \end{array} ; x \right] {}_2F_1 \left[\begin{array}{c} \beta - \epsilon, \quad \gamma + \beta - \alpha \\ \gamma + \beta \end{array} ; x \right].
\end{aligned}$$

Now, multiply both sides of (4.21) by $x^{\lambda-1}(1-x)^{\mu-\lambda-1}$ and integrating with respect to x between the interval $[0, 1]$, we have

$$(4.22) \quad \int_0^1 x^{\lambda-1}(1-x)^{\mu-\lambda-1} F_{1:0;1}^{1:1;2} \left[\begin{array}{c} \alpha : \epsilon; \quad \beta - \epsilon, \gamma \\ \beta : -; \quad \gamma + \beta \end{array} ; x, x \right] dx$$

$$(5.11) \quad F_{0:1;4}^{1:1;4} \left[\begin{array}{c} \lambda : \alpha; \quad \beta - \epsilon, \alpha + \gamma, 1 + \frac{1}{2}(\alpha + \gamma), \frac{1}{2}\alpha \\ - : \mu; \quad \beta, \frac{1}{2}(\alpha + \gamma), 1 + \frac{1}{2}\alpha + \gamma, 1 - \mu - \lambda \end{array} ; 1, 1 \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} \\ \times {}_5F_4 \left[\begin{array}{c} \alpha + \gamma, \quad 1 + \frac{1}{2}(\alpha + \gamma), \quad \frac{1}{2}\alpha, \quad \beta - \epsilon, \quad \lambda \\ \frac{1}{2}(\alpha + \gamma), \quad 1 + \frac{1}{2}\alpha + \gamma, \quad \beta, \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right].$$

$$(5.12) \quad F_{0:1;3}^{1:1;3} \left[\begin{array}{c} \lambda : \alpha; \quad \beta - \epsilon, \alpha + \gamma, \frac{1}{2}\alpha \\ - : \mu; \quad \beta, 1 + \frac{1}{2}\alpha + \gamma, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} \\ \times {}_4F_3 \left[\begin{array}{c} \beta - \epsilon, \quad \alpha + \gamma, \quad \frac{1}{2}\alpha, \quad \lambda \\ \beta, \quad 1 + \frac{1}{2}\alpha + \gamma, \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right].$$

$$(5.13) \quad F_{0:1;3}^{1:1;3} \left[\begin{array}{c} \lambda : \alpha; \quad \beta - \epsilon, \alpha + \gamma, 1 + \frac{1}{2}(\alpha + \gamma), \frac{1}{2}(\alpha - 1) \\ - : \mu; \quad \beta, \frac{1}{2}(\alpha + \gamma), \frac{3}{2} + \frac{1}{2}\alpha + \gamma, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} \\ \times {}_5F_4 \left[\begin{array}{c} \alpha + \gamma, \quad 1 + \frac{1}{2}(\alpha + \gamma), \quad \frac{1}{2}(\alpha - 1), \quad \beta - \epsilon, \quad \lambda \\ \frac{1}{2}(\alpha + \gamma), \quad \frac{3}{2} + \frac{1}{2}\alpha + \gamma, \quad \beta, \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right].$$

$$(5.14) \quad F_{0:1;2}^{1:1;2} \left[\begin{array}{c} \lambda : \alpha; \quad \alpha, \beta + \epsilon - \gamma \\ - : \mu; \quad \beta + \epsilon, \gamma - \mu + \lambda \end{array} ; 1, -1 \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} {}_3F_2 \left[\begin{array}{c} \alpha, \quad \beta + \epsilon - \gamma, \quad \lambda \\ \beta + \epsilon, \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right].$$

$$(5.15) \quad F_{0:1;2}^{1:1;2} \left[\begin{array}{c} \lambda : \alpha; \quad \alpha, 1 + \frac{1}{2}\beta \\ - : \mu; \quad \frac{1}{2}\beta, 1 - \mu + \lambda \end{array} ; 1, -1 \right] \\ = \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} {}_3F_2 \left[\begin{array}{c} \alpha, \quad 1 + \frac{1}{2}\beta, \quad \lambda \\ \frac{1}{2}\beta, \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right].$$

$$\begin{aligned}
(5.16) \quad & F_{0:1;3}^{1:1;3} \left[\begin{array}{c} \lambda : \alpha; \quad \alpha, 1 + \frac{1}{2}\beta, \frac{1}{2}(\beta - \gamma) \\ - : \mu; \quad \frac{1}{2}\beta, 1 + \frac{1}{2}(\beta + \gamma), 1 - \mu + \lambda \end{array} ; 1, -1 \right] \\
&= \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} \\
&\quad \times {}_4F_3 \left[\begin{array}{c} \alpha, \quad 1 + \frac{1}{2}\beta, \quad \frac{1}{2}(\beta - \gamma), \quad \lambda \\ \frac{1}{2}\beta, \quad 1 + \frac{1}{2}(\gamma + \beta), \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right]. \\
(5.17) \quad & F_{0:1;2}^{1:1;2} \left[\begin{array}{c} \lambda : \alpha; \quad \alpha, \frac{1}{2}(\beta - \gamma) \\ - : \mu; \quad 1 + \frac{1}{2}(\beta + \gamma), 1 - \mu + \lambda \end{array} ; 1, -1 \right] \\
&= \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} {}_3F_2 \left[\begin{array}{c} \alpha, \quad \frac{1}{2}(\beta - \gamma), \quad \lambda \\ 1 + \frac{1}{2}(\beta + \gamma), \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right]. \\
(5.18) \quad & F_{0:1;3}^{1:1;3} \left[\begin{array}{c} \lambda : \alpha; \quad \alpha, 1 + \frac{1}{2}\beta, \frac{1}{2}(\beta - \mu - 1) \\ - : \mu; \quad \frac{1}{2}\beta, \frac{1}{2}(2 + \beta + \gamma), 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\
&= \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} \\
&\quad \times {}_4F_3 \left[\begin{array}{c} \alpha, \quad 1 + \frac{1}{2}\beta, \quad \frac{1}{2}(\beta - \gamma - 1), \quad \lambda \\ \frac{1}{2}\beta, \quad \frac{1}{2}(3 + \gamma + \beta), \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right]. \\
(5.19) \quad & F_{0:1;3}^{1:1;3} \left[\begin{array}{c} \lambda : \alpha; \quad \alpha, \beta - \epsilon, \delta - \gamma \\ - : \mu; \quad \beta, \delta, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\
&= \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} \\
&\quad \times {}_4F_3 \left[\begin{array}{c} \alpha, \quad \beta - \epsilon, \quad \delta - \gamma, \quad \lambda \\ \beta, \quad \delta, \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right]. \\
(5.20) \quad & F_{0:1;4}^{1:1;4} \left[\begin{array}{c} \lambda : \alpha; \quad \alpha, \beta - \epsilon, \delta - \gamma - 1, \delta - \gamma + 2 \\ - : \mu; \quad \beta, \delta, \gamma - \delta + 1, 1 - \mu + \lambda \end{array} ; 1, 1 \right] \\
&= \frac{\Gamma(\mu)\Gamma(\mu - \alpha - \lambda)}{\Gamma(\mu - \lambda)\Gamma(\mu - \alpha)} \\
&\quad \times {}_5F_4 \left[\begin{array}{c} \alpha, \quad \beta - \epsilon, \quad \delta - \gamma - 1, \quad \delta - \gamma + 2, \quad \lambda \\ \beta, \quad \delta, \quad \gamma - \delta + 1, \quad 1 - \mu + \alpha + \lambda \end{array} ; 1 \right].
\end{aligned}$$

Proof. The derivation of the reduction formulas (5.1) to (5.20) are quite straight forward. In order to establish the reduction formula (5.1), we observe that the left-hand side of the results (3.2) and (4.1) are the same, so equating the results (3.2) and (4.1), we immediately get the reduction formula (5.1).

In exactly the same manner, on equating the results (3.3) to (3.21) to the respectively results (4.2) to (4.20), we arrive at the reduction formulas (5.2) to (5.20). \square

6. Concluding remark

In this paper, we have established a large number of reduction and transformation formulas for the Kampé de Fériet. The results are achieved with the help of the results recently obtained by Liu and Wang.

It is believed that the results established in this paper may be useful in the area of applied mathematics, engineering and mathematical physics.

Acknowledgment. The authors are grateful to the referee for carefully reading the manuscript and pointing out certain corrections.

References

- [1] M. Abramowitz and I. A. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, National Bureau of Standards Applied Mathematics Series, 55, For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC, 1964.
- [2] G. E. Andrews, R. Askey, and R. Roy, *Special Functions*, Encyclopedia of Mathematics and its Applications, 71, Cambridge University Press, Cambridge, 1999. <https://doi.org/10.1017/CB09781107325937>
- [3] P. Appell, *Sur les fonctions hypérogéométriques de plusieurs variables*, Gauthier-Villars, Paris, 1925.
- [4] P. Appell and J. Kampé de Fériet, *Fonctions hypergéométriques et hypersphériques: polynomes d'Hermite*, Gauthier-Villars, Paris, 1926.
- [5] W. N. Bailey, *Generalized hypergeometric series*, Cambridge Tracts in Mathematics and Mathematical Physics, No. 32, Stechert-Hafner, Inc., New York, 1964.
- [6] J. L. Burchnall and T. W. Chaundy, *Expansions of Appell's double hypergeometric functions*, Quart. J. Math. Oxford Ser. **11** (1940), 249–270. <https://doi.org/10.1093/qmath/os-11.1.249>
- [7] J. L. Burchnall and T. W. Chaundy, *Expansions of Appell's double hypergeometric functions. II*, Quart. J. Math. Oxford Ser. **12** (1941), 112–128. <https://doi.org/10.1093/qmath/os-12.1.112>
- [8] R. G. Buschman and H. M. Srivastava, *Series identities and reducibility of Kampé de Fériet functions*, Math. Proc. Cambridge Philos. Soc. **91** (1982), no. 3, 435–440. <https://doi.org/10.1017/S0305004100059478>
- [9] L. Carlitz, *Summation of a double hypergeometric series*, Matematiche (Catania) **22** (1967), 138–142.
- [10] W.-C. C. Chan, K. Chen, C. Chyan, and H. M. Srivastava, *Some multiple hypergeometric transformations and associated reduction formulas*, J. Math. Anal. Appl. **294** (2004), no. 2, 418–437. <https://doi.org/10.1016/j.jmaa.2004.02.008>
- [11] K.-Y. Chen and H. M. Srivastava, *Series identities and associated families of generating functions*, J. Math. Anal. Appl. **311** (2005), no. 2, 582–599. <https://doi.org/10.1016/j.jmaa.2005.03.030>

- [12] J. Choi and A. K. Rathie, *On the reducibility of Kampé de Fériet function*, Honam Math. J. **36** (2014), no. 2, 345–355. <https://doi.org/10.5831/HMJ.2014.36.2.345>
- [13] J. Choi and A. K. Rathie, *General summation formulas for the Kampé de Fériet function*, Montes Taures J. Pure Appl. Math. **1** (2019), no. 1, 107–128.
- [14] J. Choi and A. K. Rathie, *Further summation formulas for the Kampé de Fériet function*, arxiv:1912.10441.
- [15] J. Choi and A. K. Rathie, *Reducibility of Kampé de Fériet function*, Appl. Math. Sci. **9** (2015), no. 85, 4219–4232.
- [16] J. Choi and A. K. Rathie, *Reducibility of certain Kampé de Fériet function with an application to generating relations for products of two Laguerre polynomials*, Filomat **30** (2016), no. 7, 2059–2066. <https://doi.org/10.2298/FIL1607059C>
- [17] J. Choi, A. K. Rathie, and H. M. Srivastava, *Certain hypergeometric identities deducible by using the beta integral method*, Bull. Korean Math. Soc. **50** (2013), no. 5, 1673–1681. <https://doi.org/10.4134/BKMS.2013.50.5.1673>
- [18] D. Cvijović and A. R. Miller, *A reduction formula for the Kampé de Fériet function*, Appl. Math. Lett. **23** (2010), no. 7, 769–771. <https://doi.org/10.1016/j.aml.2010.03.006>
- [19] V. L. Deshpande, *Some infinite summation formulae involving Kampé de Fériet function*, Defence Sci. J. **21** (1971), 125–130.
- [20] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, *Higher transcendental functions. Vol. I*, McGraw-Hill Book Company, Inc., New York, 1953.
- [21] H. Exton, *On the reducibility of the Kampé de Fériet function*, J. Comput. Appl. Math. **83** (1997), no. 1, 119–121. [https://doi.org/10.1016/S0377-0427\(97\)86597-1](https://doi.org/10.1016/S0377-0427(97)86597-1)
- [22] P. Humbert, *The confluent hypergeometric functions of two variables*, Proc. Royal Soc. of Edinburgh **41** (1922), 73–96.
- [23] R. N. Jain, *Sum of a double hypergeometric series*, Matematiche (Catania) **21** (1966), 300–301.
- [24] J. Kampé de Fériet, *Les fonctions hypergéométriques d'ordre supérieure à deux variables*, CR Acad. Sci. Paris **173** (1921), 401–404.
- [25] P. W. Karlsson, *Some reduction formulae for double power series and Kampé de Fériet functions*, Nederl. Akad. Wetensch. Indag. Math. **46** (1984), no. 1, 31–36.
- [26] Y. S. Kim, *On certain reducibility of Kampé de Fériet function*, Honam Math. J. **31** (2009), no. 2, 167–176. <https://doi.org/10.5831/HMJ.2009.31.2.167>
- [27] Y. S. Kim, M. A. Rakha, and A. K. Rathie, *Extensions of certain classical summation theorems for the series ${}_2F_1$, ${}_3F_2$, and ${}_4F_3$ with applications in Ramanujan's summations*, Int. J. Math. Math. Sci. **2010** (2000), Art. ID 309503, 26 pp. <https://doi.org/10.1155/2010/309503>
- [28] C. Krattenthaler and K. Srinivasa Rao, *Automatic generation of hypergeometric identities by the beta integral method*, J. Comput. Appl. Math. **160** (2003), no. 1-2, 159–173. [https://doi.org/10.1016/S0377-0427\(03\)00629-0](https://doi.org/10.1016/S0377-0427(03)00629-0)
- [29] E. D. Krupnikov, *A register of computer-oriented reduction identities for the Kampé de Fériet function*, Novosibirsk, Russia, 1996.
- [30] J.-L. Lavoie, F. Grondin, and A. K. Rathie, *Generalizations of Watson's theorem on the sum of a ${}_3F_2$* , Indian J. Math. **34** (1992), no. 1, 23–32.
- [31] J.-L. Lavoie, F. Grondin, and A. K. Rathie, *Generalizations of Whipple's theorem on the sum of a ${}_3F_2$* , J. Comput. Appl. Math. **72** (1996), no. 2, 293–300. [https://doi.org/10.1016/0377-0427\(95\)00279-0](https://doi.org/10.1016/0377-0427(95)00279-0)
- [32] J.-L. Lavoie, F. Grondin, A. K. Rathie, and K. Arora, *Generalizations of Dixon's theorem on the sum of a ${}_3F_2$* , Math. Comp. **62** (1994), no. 205, 267–276. <https://doi.org/10.2307/2153407>

- [33] H. Liu and W. Wang, *Transformation and summation formulae for Kampé de Fériet series*, J. Math. Anal. Appl. **409** (2014), no. 1, 100–110. <https://doi.org/10.1016/j.jmaa.2013.06.068>
- [34] F. M. Ragab, *Expansions of Kampé de Fériet's double hypergeometric function of higher order*, J. Reine Angew. Math. **212** (1963), 113–119. <https://doi.org/10.1515/crll.1963.212.113>
- [35] E. D. Rainville, *Special Functions*, The Macmillan Company, New York, (1960), Reprinted by Chelsea Publishing Company, New York, (1971).
- [36] M. A. Rakha, M. M. Awad, and A. K. Rathie, *On a reducibility of the Kampé de Fériet function*, Math. Methods Appl. Sci. **38** (2015), no. 12, 2600–2605. <https://doi.org/10.1002/mma.3245>
- [37] M. A. Rakha and A. K. Rathie, *Generalizations of classical summation theorems for the series ${}_2F_1$ and ${}_3F_2$ with applications*, Integral Transforms Spec. Funct. **22** (2011), no. 11, 823–840. <https://doi.org/10.1080/10652469.2010.549487>
- [38] S. Saran, *Reducibility of generalised Kampé de Fériet function*, Ganita **31** (1980), no. 1-2, 89–98.
- [39] O. Shanker and S. Saran, *Reducibility of Kampé de Fériet function*, Ganita **21** (1970), no. 1, 9–16.
- [40] B. L. Sharma, *Sum of a double series*, Proc. Amer. Math. Soc. **52** (1975), 136–138. <https://doi.org/10.2307/2040117>
- [41] R. P. Singal, *Transformation formulae for the modified Kampé de Fériet function*, Math. Student **40A** (1972), 327–330.
- [42] S. N. Singh and S. P. Singh, *Transformation of Kampé de Fériet function*, Indian J. Pure Appl. Math. **13** (1982), no. 10, 1163–1166.
- [43] L. J. Slater, *Generalized Hypergeometric Functions*, Cambridge University Press, Cambridge, 1966.
- [44] H. M. Srivastava and J. Choi, *Zeta and q-Zeta functions and associated series and integrals*, Elsevier, Inc., Amsterdam, 2012. <https://doi.org/10.1016/B978-0-12-385218-2.00001-3>
- [45] H. M. Srivastava and M. C. Daoust, *A note on the convergence of Kampé de Fériet's double hypergeometric series*, Math. Nachr. **53** (1972), 151–159. <https://doi.org/10.1002/mana.19720530114>
- [46] H. M. Srivastava and P. W. Karlsson, *Multiple Gaussian hypergeometric series*, Ellis Horwood Series: Mathematics and its Applications, Ellis Horwood Ltd., Chichester, 1985.
- [47] H. M. Srivastava and R. Panda, *An integral representation for the product of two Jacobi polynomials*, J. London Math. Soc. (2) **12** (1975/76), no. 4, 419–425. <https://doi.org/10.1112/jlms/s2-12.4.419>
- [48] R. Vidūnas, *Specialization of Appell's functions to univariate hypergeometric functions*, J. Math. Anal. Appl. **355** (2009), no. 1, 145–163. <https://doi.org/10.1016/j.jmaa.2009.01.047>
- [49] A. D. Wadhwa, *A theorem on Kampé de Fériet function*, Math. Education **4** (1970), A137–A139.
- [50] A. D. Wadhwa, *Fourier series for Kampé de Fériet function*, Defence Sci. J. **21** (1971), 179–186.

MEDHAT A. RAKHA
DEPARTMENT OF MATHEMATICS
FACULTY OF SCIENCE
SUEZ CANAL UNIVERSITY
EL-SHEIK ZAYED 41522, ISMAILIA - EGYPT
Email address: medhat_rakha@science.suez.edu.eg

ARJUN K. RATHIE
DEPARTMENT OF MATHEMATICS
VEDANT COLLEGE OF ENGINEERING & TECHNOLOGY
RAJASTHAN TECHNICAL UNIVERSITY
TULSI, JAKHMUND, BUNDI-323021, RAJASTHAN STATE, INDIA
Email address: arjunkumarrathie@gmail.com