

MIAO-TAM EQUATION ON ALMOST COKÄHLER MANIFOLDS

TARAK MANDAL

ABSTRACT. In the present paper, we have studied Miao-Tam equation on three dimensional almost coKähler manifolds. We have also proved that there does not exist non-trivial solution of Miao-Tam equation on the said manifolds if the dimension is greater than three. Also we give an example to verify the deduced results.

1. Introduction

Recently, the study of differentiable manifolds endowed with certain structures, namely almost contact and almost complex structures has become a subject of growing interest due to their applications in relativity, cosmology, string theory etc. An odd dimensional differentiable manifold M equipped with a $(1, 1)$ tensor field ϕ , a vector field ξ , a 1-form η satisfying

$$(1) \quad \phi^2 X = -X + \eta(X)\xi, \quad \eta(\xi) = 1$$

for any $X \in \chi(M)$, the set of all vector fields on M , is known as an almost contact manifold [4].

If the 2-form Φ given by $\Phi(X, Y) = g(\phi X, Y)$ for any $X, Y \in \chi(M)$ and the 1-form η both are closed, then the almost contact manifold is called an almost coKähler manifold [3]. Due to additional properties that Φ and η are closed almost coKähler manifolds show some special properties which are not found generally in almost contact manifolds. Therefore almost coKähler manifolds need special attention. In 1967, Blair [3] introduced the notion of almost coKähler manifolds. The almost coKähler manifolds are odd dimensional analogues of the almost Kähler manifolds [17]. So many examples of almost coKähler manifolds have been constructed by various authors. For instance, the Riemannian product of a real line and an almost Kähler manifold admits an almost coKähler manifold [19, 23, 24]. Many authors such as Blair [5], De and Sardar [11], De, Majhi and Suh [10], Dacko [8], Dacko and Olszak [9],

Received June 27, 2021; Revised October 14, 2021; Accepted November 12, 2021.

2010 *Mathematics Subject Classification.* Primary 53C15, 53D25.

Key words and phrases. Almost coKähler manifold, critical equation, Miao-Tam equation, Einstein manifold.

Goldberg and Yano [14], Olszak [23, 24], and Wang [27, 28] have been studied almost coKähler manifolds.

In [18], Miao-Tam equation has been studied on normal almost contact manifold. Due to distinguishing nature of almost coKähler manifolds we feel the necessity of investigation of Miao-Tam equation on almost coKähler manifolds. Miao-Tam equation on a compact Riemannian manifold M^n of dimension greater than two is given by

$$(2) \quad -(\Delta_g \lambda)g + \nabla_g^2 \lambda - \lambda S = g,$$

where $\lambda : M^n \rightarrow R$ is smooth on the manifold M^n and $\lambda = 0$ on the boundary ∂M^n . Δ_g, ∇_g^2 are the negative Laplacian and Hessian operator, respectively, with respect to the metric g . For details see ([2, 20]). In the paper [21], the authors studied conformally flat Riemannian manifolds satisfying Miao-Tam equation. Miao-Tam equation on different manifolds has been studied by several other authors ([1, 7, 12, 13, 16, 25, 26]).

The present paper is organized as follows: After introduction, we give some preliminaries in Section 2. In Section 3, we have studied three dimensional (κ, μ) -almost coKähler manifolds satisfying Miao-Tam equation. In the same section we have proved that there does not exist a non-trivial solution of the Miao-Tam equation if the dimension of the (κ, μ) -almost coKähler manifold is greater than three. In the last section, we give an example to verify the deduced results.

2. Preliminaries

An almost contact manifold satisfying (1) is called an almost contact metric manifold if it admits a Riemannian metric g satisfying

$$g(\phi X, \phi Y) = g(X, Y) - \eta(X)\eta(Y).$$

As a consequence of (1) and the above equation, we have

$$\phi \xi = 0, \quad g(X, \xi) = \eta(X), \quad \eta(\phi X) = 0,$$

$$g(\phi X, Y) = -g(X, \phi Y)$$

for any vector fields $X, Y \in \chi(M)$ ([11, 15]).

For an almost coKähler manifold the 2-form Φ and the 1-form η are closed. That is, $d\Phi = 0$ and $d\eta = 0$.

On an almost coKähler manifold, we also have ([11, 15])

$$(3) \quad h\xi = 0, \quad tr(h) = 0, \quad h\phi = -\phi h,$$

$$(4) \quad \nabla_X \xi = h\phi X,$$

$$\phi l\phi - l = 2h^2$$

for any vector field $X \in \chi(M)$. Here the operators h and l are defined by $h = \frac{1}{2}\mathcal{L}_\xi\phi$ and $l = R(\cdot, \xi)\xi$, respectively, where R is the Riemannian curvature tensor and \mathcal{L} is the Lie differentiation operator. 'tr' denotes trace. The operators h and l both are symmetric.

For an almost coKähler manifold the 1-form η satisfies

$$(\nabla_X\eta)Y - (\nabla_Y\eta)X = 0$$

for any vector fields $X, Y \in \chi(M)$.

In [6], Blair et al. introduced the idea of (κ, μ) -nullity distribution in the context of contact geometry. The vector field ξ is said to belong to (κ, μ) -nullity distribution if

$$(5) \quad R(X, Y)\xi = \kappa[\eta(Y)X - \eta(X)Y] + \mu[\eta(Y)hX - \eta(X)hY]$$

for any vector fields $X, Y \in \chi(M)$ and κ, μ being real numbers. If κ, μ both are smooth functions on the manifold, then the nullity distribution is called generalized (κ, μ) -nullity distribution.

If the Reeb vector field ξ of an almost coKähler manifold belongs to (κ, μ) -nullity distribution, then it is called a (κ, μ) -almost coKähler manifold. For details see ([11, 15]). On a (κ, μ) -almost coKähler manifold, one obtains

$$(6) \quad h^2X = \kappa\phi^2X,$$

$$(7) \quad S(X, \xi) = 2n\kappa\eta(X),$$

$$(8) \quad Q\xi = 2n\kappa\xi$$

for any vector field $X \in \chi(M)$, where S and Q are the Ricci tensor of type (0,2) and Ricci operator, respectively.

Now let us recall two known lemmas:

Lemma 2.1 ([27]). *Let M^{2n+1} be a (κ, μ) -almost coKähler manifold of dimension greater than 3 with $\kappa < 0$. Then the Ricci operator Q is given by*

$$(9) \quad QX = \mu hX + 2n\kappa\eta(X)\xi$$

for all vector field X on M^{2n+1} , where κ is a non-zero constant and μ is a smooth function satisfying $d\mu \wedge \eta = 0$. Moreover, the scalar curvature of M^{2n+1} is $2n\kappa$.

Lemma 2.2 ([13]). *Let (M^{2n+1}, g) be a Riemannian manifold of dimension $(2n + 1)$ satisfying the Miao-Tam equation. Then the curvature tensor R can be expressed as*

$$(10) \quad \begin{aligned} R(X, Y)D\lambda &= (X\lambda)QY - (Y\lambda)QX + \lambda((\nabla_XQ)Y - (\nabla_YQ)X) \\ &+ (Xf)Y - (Yf)X \end{aligned}$$

for any vector fields X, Y on M , where $f = -\frac{r\lambda+1}{2n}$ and D is the gradient operator.

Moreover,

$$(11) \quad \nabla_X D\lambda = \lambda QX + fX$$

for any vector field X .

3. Almost coKähler manifolds satisfying Miao-Tam equation

In the present section, we study (κ, μ) -almost coKähler manifolds with $\kappa < 0$ which satisfy Miao-Tam equation.

Theorem 3.1. *If a (κ, μ) -almost coKähler manifold of dimension three satisfy Miao-Tam equation, then either $\mu^2 = -\frac{(6\kappa-r)^2}{4\kappa}$ or the potential function is constant, i.e., the manifold is Einstein.*

Proof. Taking inner product of (10) with the vector field ξ and using (7), we get

$$(12) \quad \begin{aligned} g(R(X, Y)D\lambda, \xi) &= 2\kappa(X\lambda)\eta(Y) - 2\kappa(Y\lambda)\eta(X) \\ &\quad + \lambda[g((\nabla_X Q)\xi, Y) - g((\nabla_Y Q)\xi, X)] \\ &\quad + (Xf)\eta(Y) - (Yf)\eta(X). \end{aligned}$$

Since any Riemannian metric g satisfying the Miao-Tam equation must have constant scalar curvature [20], r is constant. Thus we have $(Xf) = -\frac{r}{2}(X\lambda)$ and $(Yf) = -\frac{r}{2}(Y\lambda)$. Therefore from (12), we get

$$(13) \quad \begin{aligned} g(R(X, Y)D\lambda, \xi) &= (2\kappa - \frac{r}{2})[(X\lambda)\eta(Y) - (Y\lambda)\eta(X)] \\ &\quad + \lambda[g((\nabla_X Q)\xi, Y) - g((\nabla_Y Q)\xi, X)]. \end{aligned}$$

Taking covariant derivative of (8) along the vector field X , we obtain

$$(14) \quad (\nabla_X Q)\xi = 2\kappa h\phi X - Qh\phi X.$$

Using (14) in (13), we get

$$\begin{aligned} g(R(X, Y)D\lambda, \xi) &= (2\kappa - \frac{r}{2})[(X\lambda)\eta(Y) - (Y\lambda)\eta(X)] \\ &\quad - \lambda[S(h\phi X, Y) - S(X, h\phi Y)]. \end{aligned}$$

Putting $X = \xi$ in the above equation and using (7), we get

$$(15) \quad g(R(\xi, Y)D\lambda, \xi) = (2\kappa - \frac{r}{2})[(\xi\lambda)\eta(Y) - (Y\lambda)].$$

Putting $X = \xi$ in (5) and then taking inner product with $D\lambda$, we get

$$(16) \quad g(R(\xi, Y)\xi, D\lambda) = \kappa[(\xi\lambda)\eta(Y) - (Y\lambda)] - \mu(hY\lambda).$$

Since $g(R(X, Y)Z, W) = -g(R(X, Y)W, Z)$ for all vector fields X, Y, Z and W , from (15) and (16), we get

$$(17) \quad (3\kappa - \frac{r}{2})[(\xi\lambda)\eta(Y) - (Y\lambda)] = \mu(hY\lambda).$$

Replacing Y by hY in (17), we obtain

$$(18) \quad (hY\lambda) = -\frac{2\kappa\mu}{6\kappa-r}[(\xi\lambda)\eta(Y) - (Y\lambda)].$$

Therefore, from (17) and (18), we obtain

$$\frac{(6\kappa-r)^2 + 4\kappa\mu^2}{2(6\kappa-r)}[(\xi\lambda)\eta(Y) - (Y\lambda)] = 0.$$

Thus, there arise two cases

Case-I: $(6\kappa-r)^2 + 4\kappa\mu^2 = 0$, which gives $\mu^2 = -\frac{(6\kappa-r)^2}{4\kappa}$.

Case-II: $(Y\lambda) = (\xi\lambda)\eta(Y)$, which gives

$$(19) \quad D\lambda = (\xi\lambda)\xi.$$

Taking differentiation of (19) covariantly along the vector field X and using (4), we get

$$(20) \quad \nabla_X D\lambda = X(\xi\lambda)\xi + (\xi\lambda)h\phi X.$$

Comparing (11) and (20), we get

$$X(\xi\lambda)\xi + (\xi\lambda)h\phi X = \lambda QX + fX.$$

Tracing over X in the above equation and using $tr(h\phi) = 0$, we obtain

$$(21) \quad \xi(\xi\lambda) = r\lambda + 3f,$$

where r is the scalar curvature.

Putting $X = \xi$ in (11) and taking inner product with ξ , we get

$$(22) \quad \xi(\xi\lambda) = \lambda S(\xi, \xi) + f.$$

From (7), we get

$$(23) \quad S(\xi, \xi) = 2\kappa.$$

From (22) and (23), we obtain

$$(24) \quad \xi(\xi\lambda) = 2\kappa\lambda + f.$$

Comparing (21) and (24), we get

$$\lambda = -\frac{1}{2\kappa},$$

which is a non-zero constant. \square

Theorem 3.2. *There does not exist a non-trivial solution of Miao-Tam equation on a (κ, μ) -almost coKähler manifold M^{2n+1} of dimension greater than three with $\kappa < 0$.*

Proof. Taking inner product of (10) with the vector field ξ and using (7), we get

$$(25) \quad \begin{aligned} g(R(X, Y)D\lambda, \xi) &= 2n\kappa(X\lambda)\eta(Y) - 2n\kappa(Y\lambda)\eta(X) \\ &+ \lambda[g((\nabla_X Q)\xi, Y) - g((\nabla_Y Q)\xi, X)] \\ &+ (Xf)\eta(Y) - (Yf)\eta(X). \end{aligned}$$

Since r is constant, we have $(Xf) = -\frac{r}{2n}(X\lambda)$ and $(Yf) = -\frac{r}{2n}(Y\lambda)$. Therefore from (25), we get

$$(26) \quad \begin{aligned} g(R(X, Y)D\lambda, \xi) &= (2n\kappa - \frac{r}{2n})[(X\lambda)\eta(Y) - (Y\lambda)\eta(X)] \\ &+ \lambda[g((\nabla_X Q)\xi, Y) - g((\nabla_Y Q)\xi, X)]. \end{aligned}$$

Taking covariant derivative of equation (8) along the vector field X , we obtain

$$(27) \quad (\nabla_X Q)\xi = 2n\kappa h\phi X - Qh\phi X.$$

Using (27) in (26), we get

$$(28) \quad \begin{aligned} g(R(X, Y)D\lambda, \xi) &= (2n\kappa - \frac{r}{2n})[(X\lambda)\eta(Y) - (Y\lambda)\eta(X)] \\ &- \lambda[S(h\phi X, Y) - S(X, h\phi Y)]. \end{aligned}$$

Replacing X by ϕX and Y by ϕY in (28), we obtain

$$(29) \quad g(R(\phi X, \phi Y)D\lambda, \xi) = -\lambda[S(h\phi^2 X, \phi Y) - S(\phi X, h\phi^2 Y)].$$

From (3), (6) and (9), we obtain

$$(30) \quad S(h\phi^2 X, \phi Y) = \mu\kappa g(X, \phi Y).$$

Therefore, from (29) and (30), we get

$$(31) \quad g(R(\phi X, \phi Y)D\lambda, \xi) = -2\mu\kappa\lambda g(X, \phi Y).$$

Also, replacing X by ϕX and Y by ϕY in (5) and then taking inner product with $D\lambda$, we obtain

$$(32) \quad g(R(\phi X, \phi Y)\xi, D\lambda) = 0.$$

Since $g(R(X, Y)Z, W) = -g(R(X, Y)W, Z)$ for all vector fields X, Y, Z and W , from (31) and (32), we get

$$(33) \quad \mu\kappa\lambda g(X, \phi Y) = 0.$$

Since $\kappa < 0$, from (33), we get

$$\mu\lambda = 0.$$

Let us assume that $\lambda \neq 0$ in some open subset Ω of the manifold. Then $\mu = 0$ on Ω .

Now, putting $X = \xi$ in (5) and then taking inner product with $D\lambda$, we get

$$(34) \quad g(R(\xi, Y)\xi, D\lambda) = \kappa[(\xi\lambda)\eta(Y) - (Y\lambda)].$$

Again, putting $X = \xi$ in (28) and using (7), we get

$$(35) \quad g(R(\xi, Y)D\lambda, \xi) = (2n\kappa - \frac{r}{2n})[(\xi\lambda)\eta(Y) - (Y\lambda)].$$

Since $g(R(X, Y)Z, W) = -g(R(X, Y)W, Z)$ for all vector fields X, Y, Z and W , from (34) and (35), we get

$$(36) \quad 2n\kappa[(\xi\lambda)\xi - (Y\lambda)] = 0,$$

since $r = 2n\kappa$.

Therefore, from (36), we get

$$(37) \quad D\lambda = (\xi\lambda)\xi,$$

since $\kappa < 0$.

Taking covariant derivative of (37) and using (4), we get

$$(38) \quad \nabla_X D\lambda = X(\xi\lambda)\xi + (\xi\lambda)h\phi X.$$

Also, from (9) and (11), we get

$$(39) \quad \nabla_X D\lambda = 2n\kappa\lambda\eta(X)\xi - (\kappa\lambda + \frac{1}{2n})X.$$

Comparing (38) and (39), we get

$$(40) \quad X(\xi\lambda)\xi + (\xi\lambda)h\phi X = 2n\kappa\lambda\eta(X)\xi - (\kappa\lambda + \frac{1}{2n})X.$$

Tracing over X in (40) and using $tr(h\phi) = 0$, we obtain

$$(41) \quad \xi(\xi\lambda) = -(\kappa\lambda + \frac{2n+1}{2n}).$$

Again, putting $X = \xi$ in (39) and then taking inner product with ξ , we get

$$(42) \quad \xi(\xi\lambda) = 2n\kappa\lambda - (\kappa\lambda + \frac{1}{2n}).$$

From (41) and (42), we get

$$(43) \quad 2n\kappa\lambda + 1 = 0.$$

Again, operating (40) by ϕ and using (3) and (43), we get

$$(\xi\lambda) = 0.$$

Therefore, from (37), λ is a constant.

Thus, from (2), we get

$$(44) \quad S(X, Y) = -\frac{1}{\lambda}g(X, Y).$$

Putting $X = Y = e_i$ in (44), where $\{e_i\}$ is the orthonormal basis of the tangent space of the manifold and summing over i , $1 \leq i \leq 2n+1$, we obtain

$$\lambda = -\frac{2n+1}{2n\kappa},$$

since $r = 2n\kappa$.

Thus

$$(45) \quad S(X, Y) = \frac{2n\kappa}{2n+1}g(X, Y).$$

Putting $X = Y = \xi$ in (45) and using (7), we get $\kappa = 0$, which is a contradiction. \square

4. Example

In the paper [22], the authors have constructed an example of almost coKähler manifold of dimension three. Now we construct an example of almost coKähler manifold of dimension five following this example.

Let us consider the manifold $M = \{x, y, u, v, z \in \mathbb{R}^5 : z \neq 0\}$ of dimension 5, where $\{x, y, u, v, z\}$ are standard co-ordinates in \mathbb{R}^5 . We choose the vector fields

$$\begin{aligned} e_1 &= e^{\frac{z}{2}} \frac{\partial}{\partial x} - e^{-\frac{z}{2}} \frac{\partial}{\partial y}, & e_2 &= e^{\frac{z}{2}} \frac{\partial}{\partial x} + e^{-\frac{z}{2}} \frac{\partial}{\partial y}, \\ e_3 &= e^{\frac{z}{2}} \frac{\partial}{\partial u} - e^{-\frac{z}{2}} \frac{\partial}{\partial v}, & e_4 &= e^{\frac{z}{2}} \frac{\partial}{\partial u} + e^{-\frac{z}{2}} \frac{\partial}{\partial v}, & e_5 &= \frac{\partial}{\partial z}, \end{aligned}$$

which are linearly independent at each point of M . We get the following by direct computations

$$\begin{aligned} [e_1, e_2] &= 0, & [e_1, e_3] &= 0, & [e_1, e_4] &= 0, & [e_1, e_5] &= -\frac{1}{2}e_2, \\ [e_2, e_3] &= 0, & [e_2, e_4] &= 0, & [e_2, e_5] &= -\frac{1}{2}e_1, \\ [e_3, e_4] &= 0, & [e_3, e_5] &= -\frac{1}{2}e_4, & [e_4, e_5] &= -\frac{1}{2}e_3. \end{aligned}$$

Let the metric tensor g be defined by

$$g(e_1, e_1) = g(e_2, e_2) = g(e_3, e_3) = g(e_4, e_4) = g(e_5, e_5) = 1$$

and $g(e_i, e_j) = 0$ for all $i \neq j$; $i, j = 1, 2, 3, 4, 5$.

The 1-form η is defined by $\eta(X) = g(X, e_5)$ for all X on M . Let ϕ be the (1, 1)-tensor field defined by

$$\phi(e_1) = -e_2, \quad \phi(e_2) = e_1, \quad \phi(e_3) = -e_4, \quad \phi(e_4) = e_3, \quad \phi(e_5) = 0.$$

Then we find that

$$\begin{aligned} \eta(e_5) &= 1, & \phi^2 X &= -X + \eta(X)e_5, \\ g(\phi X, \phi Y) &= g(X, Y) - \eta(X)\eta(Y) \end{aligned}$$

for any vector fields X, Y on M . Thus (ϕ, e_5, η, g) defines an almost contact metric structure.

For the Levi-Civita connection ∇ with respect to the metric g on M , we can write

$$\begin{aligned} 2g(\nabla_X Y, Z) &= Xg(Y, Z) + Yg(Z, X) - Zg(X, Y) \\ &\quad - g(X, [Y, Z]) - g(Y, [X, Z]) + g(Z, [X, Y]), \end{aligned}$$

which is known as Koszul's formula.

By Koszul's formula, we get the following expressions:

$$\begin{aligned}
\nabla_{e_1}e_1 &= 0, & \nabla_{e_1}e_2 &= \frac{1}{2}e_5, & \nabla_{e_1}e_3 &= 0, & \nabla_{e_1}e_4 &= 0, \\
\nabla_{e_1}e_5 &= -\frac{1}{2}e_2, & \nabla_{e_2}e_2 &= 0, & \nabla_{e_2}e_1 &= \frac{1}{2}e_5, & \nabla_{e_2}e_3 &= 0, \\
\nabla_{e_2}e_4 &= 0, & \nabla_{e_2}e_5 &= -\frac{1}{2}e_1, & \nabla_{e_3}e_1 &= 0, & \nabla_{e_3}e_2 &= 0, \\
\nabla_{e_3}e_3 &= 0, & \nabla_{e_3}e_4 &= \frac{1}{2}e_5, & \nabla_{e_3}e_5 &= -\frac{1}{2}e_4, & \nabla_{e_4}e_1 &= 0, \\
\nabla_{e_4}e_2 &= 0, & \nabla_{e_4}e_3 &= \frac{1}{2}e_5, & \nabla_{e_4}e_4 &= 0, & \nabla_{e_4}e_5 &= -\frac{1}{2}e_3, \\
\nabla_{e_5}e_1 &= 0, & \nabla_{e_5}e_2 &= 0, & \nabla_{e_5}e_3 &= 0, \\
\nabla_{e_5}e_4 &= 0, & \nabla_{e_5}e_5 &= 0.
\end{aligned}$$

From the above expressions of ∇ , we conclude that the given manifold is an almost coKähler manifold with $he_1 = -\frac{1}{2}e_1$, $he_2 = \frac{1}{2}e_2$, $he_3 = -\frac{1}{2}e_3$, $he_4 = \frac{1}{2}e_4$ and $he_5 = 0$.

Using the formula $R(X, Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]}Z$, we get

$$\begin{aligned}
R(e_1, e_2)e_1 &= -\frac{1}{4}e_2, & R(e_1, e_2)e_2 &= \frac{1}{4}e_1, & R(e_2, e_5)e_5 &= -\frac{1}{4}e_2, \\
R(e_5, e_2)e_2 &= -\frac{1}{4}e_5, & R(e_1, e_5)e_5 &= -\frac{1}{4}e_1, & R(e_5, e_1)e_1 &= -\frac{1}{4}e_5, \\
R(e_3, e_4)e_3 &= -\frac{1}{4}e_4, & R(e_3, e_4)e_4 &= \frac{1}{4}e_3, & R(e_3, e_5)e_5 &= -\frac{1}{4}e_3, \\
R(e_5, e_3)e_3 &= -\frac{1}{4}e_5, & R(e_4, e_5)e_5 &= -\frac{1}{4}e_4, & R(e_5, e_4)e_4 &= -\frac{1}{4}e_5
\end{aligned}$$

and the remaining $R(e_i, e_j)e_k = 0$, $i, j, k = 1, 2, 3, 4, 5$.

From above, we conclude that the manifold is a (κ, μ) -almost coKähler manifold with $\kappa = -\frac{1}{4}$ and $\mu = 0$.

From the expressions of curvature tensor, we get

$$\begin{aligned}
S(e_1, e_1) &= 0, & S(e_2, e_2) &= 0, & S(e_3, e_3) &= 0, \\
S(e_4, e_4) &= 0, & S(e_5, e_5) &= -1
\end{aligned}$$

and $S(e_i, e_j) = 0$, for all $i \neq j$; $i, j = 1, 2, 3, 4, 5$.

Let r be the scalar curvature. Then from above, we get

$$r = S(e_1, e_1) + S(e_2, e_2) + S(e_3, e_3) + S(e_4, e_4) + S(e_5, e_5) = -1.$$

Let $\lambda = e^z + b$, where b is a constant. Therefore, $D\lambda = e^z e_5 = (\lambda - b)e_5$.

Thus we get the following:

$$\begin{aligned}
\nabla_{e_1}D\lambda &= -\frac{1}{2}(\lambda - b)e_2, & \nabla_{e_2}D\lambda &= -\frac{1}{2}(\lambda - b)e_1, \\
\nabla_{e_3}D\lambda &= -\frac{1}{2}(\lambda - b)e_4, & \nabla_{e_4}D\lambda &= -\frac{1}{2}(\lambda - b)e_3, & \nabla_{e_5}D\lambda &= (\lambda - b)e_5.
\end{aligned}$$

From the above expressions, we get

$$\Delta_g \lambda = (\lambda - b).$$

Thus,

$$\begin{aligned} -(\Delta_g \lambda)g(e_1, e_1) + g(\nabla_{e_1} D\lambda, e_1) - \lambda S(e_1, e_1) &= -(\lambda - b), \\ -(\Delta_g \lambda)g(e_2, e_2) + g(\nabla_{e_2} D\lambda, e_2) - \lambda S(e_2, e_2) &= -(\lambda - b), \\ -(\Delta_g \lambda)g(e_3, e_3) + g(\nabla_{e_3} D\lambda, e_3) - \lambda S(e_3, e_3) &= -(\lambda - b), \\ -(\Delta_g \lambda)g(e_4, e_4) + g(\nabla_{e_4} D\lambda, e_4) - \lambda S(e_4, e_4) &= -(\lambda - b), \\ -(\Delta_g \lambda)g(e_5, e_5) + g(\nabla_{e_5} D\lambda, e_5) - \lambda S(e_5, e_5) &= \lambda. \end{aligned}$$

From the last five expressions, we conclude that there does not exist any solution of the Miao-Tam equation on the given manifold and this verifies Theorem 3.2.

Acknowledgement. The author is thankful to the referee and the Editor for their valuable suggestions towards the improvement of the paper.

References

- [1] A. Barros and E. Ribeiro, Jr., *Critical point equation on four-dimensional compact manifolds*, Math. Nachr. **287** (2014), no. 14-15, 1618–1623. <https://doi.org/10.1002/mana.201300149>
- [2] A. L. Besse, *Einstein manifolds*, reprint of the 1987 edition, Classics in Mathematics, Springer-Verlag, Berlin, 2008.
- [3] D. E. Blair, *The theory of quasi-Sasakian structures*, J. Differential Geometry **1** (1967), 331–345. <http://projecteuclid.org/euclid.jdg/1214428097>
- [4] D. E. Blair, *Contact manifolds in Riemannian geometry*, Lecture Notes in Mathematics, Vol. 509, Springer-Verlag, Berlin, 1976.
- [5] D. E. Blair, *Riemannian geometry of contact and symplectic manifolds*, Progress in Mathematics, 203, Birkhäuser Boston, Inc., Boston, MA, 2002. <https://doi.org/10.1007/978-1-4757-3604-5>
- [6] D. E. Blair, T. Koufogiorgos, and B. J. Papantoniou, *Contact metric manifolds satisfying a nullity condition*, Israel J. Math. **91** (1995), no. 1-3, 189–214. <https://doi.org/10.1007/BF02761646>
- [7] X. Chen, *On almost f -cosymplectic manifolds satisfying the Miao-Tam equation*, J. Geom. **111** (2020), no. 2, Paper No. 28, 12 pp. <https://doi.org/10.1007/s00022-020-00542-7>
- [8] P. Dacko, *On almost cosymplectic manifolds with the structure vector field ξ belonging to the k -nullity distribution*, Balkan J. Geom. Appl. **5** (2000), no. 2, 47–60.
- [9] P. Dacko and Z. Olszak, *On conformally flat almost cosymplectic manifolds with Kählerian leaves*, Rend. Sem. Mat. Univ. Politec. Torino **56** (1998), no. 1, 89–103 (2000).
- [10] U. C. De, P. Majhi, and Y. J. Suh, *Semisymmetric properties of almost coKähler 3-manifolds*, Bull. Korean Math. Soc. **56** (2019), no. 1, 219–228. <https://doi.org/10.4134/BKMS.b180180>
- [11] U. C. De and A. Sardar, *Classification of (k, μ) -almost co-Kähler manifolds with vanishing Bach tensor and divergence free Cotton tensor*, Commun. Korean Math. Soc. **35** (2020), no. 4, 1245–1254. <https://doi.org/10.4134/CKMS.c200091>
- [12] A. Ghosh and D. S. Patra, *The critical point equation and contact geometry*, J. Geom. **108** (2017), no. 1, 185–194. <https://doi.org/10.1007/s00022-016-0333-3>

- [13] A. Ghosh and D. S. Patra, *Certain almost Kenmotsu metrics satisfying the Miao-Tam equation*, Publ. Math. Debrecen **93** (2018), no. 1-2, 107–123.
- [14] S. I. Goldberg and K. Yano, *Integrability of almost cosymplectic structures*, Pacific J. Math. **31** (1969), 373–382. <http://projecteuclid.org/euclid.pjm/1102977874>
- [15] D. Kar and P. Majhi, *Beta-almost Ricci solitons on almost coKähler manifolds*, Korean J. Math. **27** (2019), no. 3, 691–705. <https://doi.org/10.11568/kjm.2019.27.3.691>
- [16] D. Kar and P. Majhi, *Almost coKähler manifolds satisfying Miao-Tam equation*, J. Geom. **110** (2019), no. 1, Paper No. 4, 10 pp. <https://doi.org/10.1007/s00022-018-0460-0>
- [17] H. Li, *Topology of co-symplectic/co-Kähler manifolds*, Asian J. Math. **12** (2008), no. 4, 527–543. <https://doi.org/10.4310/AJM.2008.v12.n4.a7>
- [18] T. Mandal, *Miao-Tam equation on normal almost contact metric manifolds*, Differ. Geom. Dyn. Syst. **23** (2021), 135–143.
- [19] J. C. Marrero and E. Padron, *New examples of compact cosymplectic solvmanifolds*, Arch. Math. (Brno) **34** (1998), no. 3, 337–345.
- [20] P. Miao and L.-F. Tam, *On the volume functional of compact manifolds with boundary with constant scalar curvature*, Calc. Var. Partial Differential Equations **36** (2009), no. 2, 141–171. <https://doi.org/10.1007/s00526-008-0221-2>
- [21] P. Miao and L.-F. Tam, *Einstein and conformally flat critical metrics of the volume functional*, Trans. Amer. Math. Soc. **363** (2011), no. 6, 2907–2937. <https://doi.org/10.1090/S0002-9947-2011-05195-0>
- [22] D. M. Naik, V. Venkatesha, and H. A. Kumara, *Ricci solitons and certain related metrics on almost co-Kähler manifolds*, Zh. Mat. Fiz. Anal. Geom. **16** (2020), no. 4, 402–417.
- [23] Z. Olszak, *On almost cosymplectic manifolds*, Kodai Math. J. **4** (1981), no. 2, 239–250. <http://projecteuclid.org/euclid.kmj/1138036371>
- [24] Z. Olszak, *Almost cosymplectic manifolds with Kählerian leaves*, Tensor (N.S.), **46** (1987), 117–124.
- [25] D. S. Patra and A. Ghosh, *Certain contact metrics satisfying the Miao-Tam critical condition*, Ann. Polon. Math. **116** (2016), no. 3, 263–271. <https://doi.org/10.4064/ap3704-11-2015>
- [26] A. Sarkar and G. G. Biswas, *Critical point equation on K-paracontact manifolds*, Balkan J. Geom. Appl. **25** (2020), no. 1, 117–126.
- [27] Y. Wang, *A generalization of the Goldberg conjecture for coKähler manifolds*, Mediterr. J. Math. **13** (2016), no. 5, 2679–2690. <https://doi.org/10.1007/s00009-015-0646-8>
- [28] Y. Wang, *Almost co-Kähler manifolds satisfying some symmetry conditions*, Turkish J. Math. **40** (2016), no. 4, 740–752.

TARAK MANDAL
DEPARTMENT OF MATHEMATICS
JANGIPUR COLLEGE
JANGIPUR, MURSHIDABAD 742213, WEST BENGAL, INDIA
Email address: mathtarak@gmail.com